

Evaluation of maltose-based cationic liposomes with different hydrophobic tails for plasmid DNA delivery

Bo Li[†], Liangliang Deng[†], Meiyan Liu, Youlin Zeng^{*}

(National & Local Joint Engineering Laboratory for New Petro-chemical Materials and Fine Utilization of Resources, Hunan Normal University, Changsha, Hunan, 410081, P. R. China)

Contents

Fig. S1	S1
NMR spectrums of intermediates and lipids IX a (Malt-DiC12MA)	S2
NMR spectrums of intermediates and lipids IX b (Malt-DiC14MA)	S22
NMR spectrums of intermediates and lipids IX c (Malt-DiC16MA)	S30




Fig. S1 Fluorescence microscopic images (100×) of cellular uptake of Malt-DiC12MA Lipids/DNA complexes at the N/P ratio of 8:1 in HEK293 after 4 h of gene transfection. The upper column (A) represents single fluorescent images, the down column (B) represents merger images. Bar: 20 μm. (Green: Dio label cytomembrane, Red: Cy3-labeled pDNA, Blue: Hoechst 33342 stained cell nuclei).

* Corresponding author. Tel.: +86 13975805056; fax: +86 73188872531

E-mail address: youlinzengcn@gmail.com (Youlin Zeng).

Project supported by the Nation Nature Sciences Foundation of China General Program: 21272064)

[†]These authors have contributed equally to this work.

