

Supplementary material

Copper-catalyzed synthesis of unsymmetrical diorganyl chalcogenides (Te/Se/S) from boronic acids under solvent-free conditions

**Sumbal Saba^{1,2}, Giancarlo V. Botteselle³, Marcelo Godoi⁴, Tiago E. A. Frizon⁵, Fábio Z. Galetto¹,
Jamal Rafique^{1,*} and Antonio L. Braga^{1,*}**

¹ Departamento de Química, Universidade Federal de Santa Catarina, UFSC, Florianópolis 88040-900, SC, Brazil



² Department of chemistry, Shaheed Benazir Bhutto Women University, Peshawar, 25000, KPK, Pakistan



³ Laboratório de Pesquisa Química, CECE, Universidade Estadual do Oeste do Paraná UNIOESTE, 85819-110, PR, Brazil



⁴ Escola de Química e Alimentos, Universidade Federal do Rio Grande-Campus Santo Antônio da Patrulha, RS, Brazil



⁵ Universidade Federal de Santa Catarina, UFSC, Campus Araranguá, 88905-120, SC, Brazil

*E-mail: jamal.chm@gmail.com (J.R.); braga.antonio@ufsc.br (A.L.B.)









—148.5
—142.4
—139.6
—131.1
—130.1
—130.0
—129.1
—122.5
—117.0
—113.2

