

## Supplementary Materials

### Purified phlorizin from *Docynia Indica* (Wall.) Decne by HSCCC, compared with whole extract, phlorizin and non-phlorizin fragment ameliorate obesity, insulin resistance and improves intestinal barrier function in high-fat-diet-fed mice

Xiao-yu Zhang<sup>1†</sup>, Kang Yi<sup>1†</sup>, Jiang Chen<sup>2</sup>, Rui-ping Li<sup>1</sup>, Jie Xie<sup>1</sup>, Yan Jin<sup>1</sup>, Xue-ran Mei<sup>3</sup>, Yao-jun Li<sup>1</sup>, Gang Liu<sup>1</sup>, Zhan-guo Wang<sup>4\*</sup>

1 College of Life Sciences, Sichuan Normal University, Longquan, Chengdu 610101, China

2 State Key Laboratory for Quality Research in Chinese Medicine, University of Macau, Macao SAR, China

3 College of Life Sciences, Sichuan University, Wuhou, Chengdu 610065, China

4 School of Medicine and Nursing, Chengdu University, Longquan, Chengdu 610106, China

\* Correspondence: wangzhanguo@cdu.edu.cn; Tel: +86 28 84617082; Fax: +86 28 84617082,

† These authors contributed equally to this work.

#### Supplementary Table

**Table 2** The ingredients of the normal chow diet (NCD).

Macronutrient	gm%	kcal%
Protein	20.4	20
Carbohydrate	65.2	70
Fat	5.3	10
Total	---	100
Ingredient	gm	kcal
Casein, 30 mesh	200	800
L-cystine	3	12
Corn starch	315	1260
Maltodextrin 10	35	140
Sucrose	350	1400
Microcrystalline cellulose	50	0
Soybean oil	25	225
Lard	20	180
V10001 mixed vitamin.	10	40
Total	1008	4057

**Table 2** The ingredients of the high fat diet (HFD).

<b>Macronutrient</b>	<b>gm%</b>	<b>kcal%</b>
Protein	24	20
Carbohydrate	41	35
Fat	24	45
Total	---	100
<b>Ingredient</b>	<b>gm</b>	<b>kcal</b>
Casein, 30 mesh	200	800
L-cystine	3	12
Corn starch	72.8	291
Maltodextrin 10	100	400
Sucrose	172.8	691
Microcrystalline cellulose	50	0
Soybean oil	25	225
Lard	177.5	1598
V10001 mixed vitamin.	10	40
Total	811.1	4057



© 2018 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).