

Cytisine-Pterocarpin Derived Compounds - Biomimetic Synthesis and apoptosis-inducing activity in Human Breast Cancer Cells

Ting-Ting Peng^a, Xuan-Rong Sun^b, Ren-Hao Liu^a, Lu-Xia Hua^a, Dong-Ping Cheng^a, Bin Mao^b,
Xing-Nuo Li^{a*}

^aCollege of Pharmaceutical Science, Zhejiang University of Technology, Hangzhou, 310014,
P.R. China.

^bCollaborative Innovation Center of Yangtze River Delta Region Green Pharmaceuticals,
Zhejiang University of Technology, Hangzhou, 310014, P.R. China.


t

Table of contents


I. Spectra (NMR) of 3	S2
II. Spectra (NMR & Mass) and HPLC chromatograms of 4	S3

I. NMR (^1H & ^{13}C) of 3

^1H NMR spectrum of 3




^{13}C NMR spectrum of 3




II. NMR (^1H & ^{13}C), ESI-MS and HPLC chromatograms of 4


^1H NMR spectrum of 4




^{13}C NMR spectrum of 4



ESI-MS spectrum of 4



HPLC chromatogram of 4



— Channel:2998; Processed Channel: PDA 310 nm

Processed Channel Descr.:PDA 310.0 nm

	Processed Channel Descr.	RT	Area	% Area	Height
1	PDA 310.0 nm	17.384	5712720	99.50	290778
2	PDA 310.0 nm	18.335	28600	0.50	2519