

Article

# Functional Characterisation of New Sesquiterpene Synthase from the Malaysian Herbal Plant, *Polygonum Minus*

Nor Azizun Rusdi <sup>1,2</sup>, Hoe-Han Goh <sup>1</sup> , Suriana Sabri <sup>3,4</sup> , Ahmad Bazli Ramzi <sup>1</sup>,  
Normah Mohd Noor <sup>1</sup> and Syarul Nataqain Baharum <sup>1,\*</sup> 

<sup>1</sup> Institute of Systems Biology (INBIOSIS), Universiti Kebangsaan Malaysia, Bangi 43600 UKM, Selangor, Malaysia; azizun@ums.edu.my (N.A.R.); gohgh@ukm.edu.my (H-H.G.); bazliramzi@ukm.edu.my (A.B.R.); normah@ukm.edu.my (N.M.N.)

<sup>2</sup> Institutes for Tropical Biology and Conservation, Universiti Malaysia Sabah, Jalan UMS, Kota Kinabalu 88400, Sabah, Malaysia

<sup>3</sup> Enzyme and Microbial Technology Research Center, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, UPM Serdang 43400, Malaysia; suriana@upm.edu.my

<sup>4</sup> Department of Microbiology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, UPM Serdang 43400, Malaysia

\* Correspondence: nataqain@ukm.edu.my; Tel.: +603-892-145-50; Fax: +603-892-133-98

Received: 23 March 2018; Accepted: 31 May 2018; Published: 4 June 2018



**Abstract:** *Polygonum minus* (syn. *Persicaria minor*) is a herbal plant that is well known for producing sesquiterpenes, which contribute to its flavour and fragrance. This study describes the cloning and functional characterisation of *PmSTPS1* and *PmSTPS2*, two sesquiterpene synthase genes that were identified from *P. minus* transcriptome data mining. The full-length sequences of the *PmSTPS1* and *PmSTPS2* genes were expressed in the *E. coli* pQE-2 expression vector. The sizes of *PmSTPS1* and *PmSTPS2* were 1098 bp and 1967 bp, respectively, with open reading frames (ORF) of 1047 and 1695 bp and encoding polypeptides of 348 and 564 amino acids, respectively. The proteins consist of three conserved motifs, namely, Asp-rich substrate binding (DDxxD), metal binding residues (NSE/DTE), and cytoplasmic ER retention (RxR), as well as the terpene synthase family N-terminal domain and C-terminal metal-binding domain. From the in vitro enzyme assays, using the farnesyl pyrophosphate (FPP) substrate, the *PmSTPS1* enzyme produced multiple acyclic sesquiterpenes of  $\beta$ -farnesene,  $\alpha$ -farnesene, and farnesol, while the *PmSTPS2* enzyme produced an additional nerolidol as a final product. The results confirmed the roles of *PmSTPS1* and *PmSTPS2* in the biosynthesis pathway of *P. minus*, to produce aromatic sesquiterpenes.

**Keywords:** farnesol; nerolidol; *Polygonum minus*; sesquiterpene synthase

## 1. Introduction

Over the last 25 years, nearly 65,000 chemical structures of terpenoids have been discovered, making terpenoids the class of natural products with the greatest structural diversity [1,2]. Terpenoids are involved in a variety of important functions in regulating plant growth (especially for terpenoid lactones) and play an ecological role in attracting pollinators [3]. Terpenoids are grouped into different classes based on the number of 5-carbon building blocks [4–7]. All terpenoids are derived from the common phosphorylated five-carbon (C5) building units, isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) [8]. There are two major pathways involved in the biosynthesis of terpenoids, namely, the mevalonate (MVA) pathway, which is primarily found in eukaryotes, and the methylerythritol phosphate (MEP) pathway (non-mevalonate pathway), which is primarily found in prokaryotes and plant chloroplasts [9–11]. For sesquiterpene biosynthesis, IPP and DMAPP undergo condensation to form farnesyl pyrophosphate

(FPP), which is catalysed by the enzyme FPP synthase. The resulting linear FPP undergoes electrophilic cyclisation and rearrangement to form acyclic and non-acyclic sesquiterpenes, based on the sesquiterpene synthase enzyme reactions [12,13].

In particular, monoterpenes and sesquiterpenes are commonly present in plant essential oils, which are widely utilised for commercial purposes, such as fragrances, cosmetics, pharmaceuticals, medicine, biofuel precursors, and industrial materials [14,15]. To date, over 7000 sesquiterpenes (C<sub>15</sub>) with different hydrocarbon skeletons and stereo-chemically structures have been reported [16]. Studies on naturally derived 15-carbon terpenoids in aromatic herbal plants as anti-malarial, anti-microbial, and insecticidal agents have also increased over the last several years [17]. Hence, many sesquiterpenes are found as major components of fruits and vegetables, floral scents, and essential oils of herbs. In previous studies, a number of genes encoding the sesquiterpene enzymes, which control the key steps of secondary metabolic pathways, have been extensively cloned and characterized from a number of herbal plant species. For example, squalene synthase from *Tripterygium wilfordii* [18], germacrene A synthase from *Achillea millefolium* [19], drimenol synthase from *Valeriana officinalis* [20], E-E-farnesol synthase and  $\alpha$ -bisabolene synthase from *Ginkgo biloba* [21],  $\alpha$ -humulene synthase from *Zingiber zerumbet* Smith [22], (+)-epi- $\alpha$ -bisabolol synthase from *Lippia dulcis* [23],  $\beta$ -caryophyllene synthase from *Ocimum basilicum* L. [24],  $\alpha$ -humulene synthase from *Zingiber zerumbate* Smith [25], and Germacrene D synthase from *Zingiber officinale* [26].

*Polygonum minus* (syn. *Persicaria minor*) is a herbal plant that originated from Southeast Asia and belongs to the Polygonaceae family. In Malaysia, *P. minus* is locally known as a 'kesum', and is commonly used as a food additive and flavouring agent. Many studies have been carried out because of the popularity of the *P. minus* as a potential medicinal plant with high antioxidant and antimicrobial activities and strong anti-inflammatory properties [27–30]. *P. minus* is an economically important herbal plant because of its essential oils, which cause it to emit a strong scent from a simple mixture of terpenoid hydrocarbons (monoterpenoids and sesquiterpenoids), including  $\beta$ -farnesene,  $\alpha$ -farnesene, nerolidol, farnesol, caryophyllene,  $\alpha$ -bergamotene, and drimenol [31,32]. Other characteristic components that have been identified from *P. minus* include aldehydes (decanal and dodecanal), esters, and organic acids [33]. Moreover, recent studies have revealed that the sesquiterpenes compounds are the main contributors to the characteristic fragrance of this plant [34]. These studies have shown the potential for developing *P. minus* as a resource to produce natural products. While many structurally diverse secondary metabolites, especially terpenoid compounds, have been identified in *P. minus*, its biosynthesis of major constituents remains unclear because of the limited genomic information that is available for this plant. Therefore, this plant is investigated for the isolation and characterisation of novel sesquiterpene synthase genes. Additionally, there is not much work on sesquiterpene synthase from *P. minus*, especially at the genetic level, as it is still very scarce.

In previous studies, several works on sesquiterpene synthases were identified. The first *P. minus* putative sesquiterpene synthase gene was cloned and expressed in *E. coli* systems [35]. Song et al. [36] successfully overexpressed a sesquiterpene synthase, *PmSTS*, in metabolically engineered gram-positive bacteria, *Lactococcus lactis*, with the MVA pathway. Then, a structural study was performed that demonstrated an active catalytic site with the same gene sequence [37]. The encoded enzyme was named  $\beta$ -sesquiphellandrene synthase, based on the principal product that was formed. In a subsequent study, the influence of jasmonic acid treatment on the expression level of the *Persicaria minor* sesquiterpene synthase (*PmSS*) gene was reported [38]. Until very recently, the purification and overexpression of *P. minor* sesquiterpene synthase encoded as *PmSTS* recombinant protein in pET28b vector, using the *E. coli* BL21 (DE3) strain, were reported [39]. All of the studies on *P. minus* that have been mentioned above were based on the same sesquiterpene synthase. In order to understand the terpenoid metabolism of *P. minus*, the potential gene for terpene synthases must be isolated and studied. Therefore, in this study, we described the cDNAs isolation and characterisation of two new sesquiterpene synthases (STPS) that were responsible for the formation of two key aromatic compounds, which made substantial contributions to the flavour and fragrance of *P. minus* essential oil. The results could have provided a foundation for the further exploration of gene function in *P. minus*, and helped to reveal the regulation of terpenoid biosynthesis.

## 2. Results

### 2.1. Screening and Isolation of Sesquiterpene Synthase Gene from *P. minus*

Two new candidates of sesquiterpene synthase genes, *PmSTPS1* (comp62410\_co\_seq6) and *PmSTPS2* (comp47018\_c0\_seq1), were successfully identified through the sequence analysis of *P. minus* transcriptome [40]. The 1098 bp *PmSTPS1* transcript contained an open reading frame (ORF) of 1047 bp, encoding 348 amino acids with a calculated molecular mass of 40.9 kDa and an isoelectric point (pI) of 6.15 (Figure S1). The ORF of *PmSTPS1* started from the nucleotide position at 25 and ended at position 1071. The deduced amino acid sequence of *PmSTPS1* (GenBank accession no. MG921605) showed no signal peptide. A ProtParam analysis of the predicted amino acid sequence of *PmSTPS1* revealed 47 negatively charged residues (Asp and Glu) and 52 positively charged residues (Arg and Lys), which represented the aliphatic index of this protein. This was a positive factor for the increased thermostability of the globular protein. The second sesquiterpene synthase transcript, *PmSTPS2* (GenBank accession No: MG921606), was 1974 bp long and had an ORF of 1695, which encoded a polypeptide of 564 amino acids (Figure S2). The ORF of *PmSTPS2* started from the nucleotide position at 97 and ended at position 1798. The calculated molecular mass of the mature protein was approximately 65.96 kDa, with a predicted pI of 5.75. A ProtParam analysis of the predicted amino acid sequence of *PmSTPS2* identified 83 negatively charged residues (Asp and Glu) and 70 positively charged residues (Arg and Lys).

Based on the BLASTx analysis, the predicted amino acid sequences of *PmSTPS1* (Table 1) and *PmSTPS2* (Table 2) had the closest hit to drimenol synthase from *Persicaria hydropiper*, with a 96% and 46% identity, respectively. The predicted amino acid sequence of *PmSTPS2* was consistent with those of other sesquiterpene synthases encoding proteins of 550–580 amino acids, with molecular weights of 60–70 kDa. Conversely, the length of *PmSTPS1*, with only 348 amino acids, was much shorter than that of the other sesquiterpene synthases. Therefore, only *PmSTPS2* met the range of other reported plant terpene synthases [21,41–45].

The presence of the conserved domains in the *PmSTPS1* and *PmSTPS2* proteins was consistent with and similar to that of the other terpene synthase features. The terpene synthase family N-terminal domain (PF01397) and the synthase family C-terminal metal-binding domain (PF03936) contained highly conserved aspartate-rich motifs (DDxxD), which were essential for enzyme-substrate binding and catalytic function. The first aspartate-rich motif played a role in the determination of the chain length for the resulting prenyl pyrophosphate.

Based on the multiple sequence alignment of *PmSTPS1* (Figure 1), several conserved motifs that were found in typical terpene synthases were identified, including the DDxxD (residue 100–104) and NSE/DTE (residue 245–253) motifs. The DDxxD and NSE/DTE motifs flanked the entrance of the active site. In addition to these motifs, there was a highly conserved arginine-rich RxR motif, which was involved in the complexing of the diphosphate group, after the ionisation of FPP [18,39]. The RxR motif was located at 45 amino acids, upstream of the first DDxxD motif. For *PmSTPS2*, the conserved arginine-rich (RxR) region at amino acid position 278–281 was conserved in all of the terpene synthases [19]. Moreover, the aspartate-rich motif of DDxxD, which might have been the Mg<sup>2+</sup> binding site, was located at position 314–318 of the amino acid sequence. Another metal binding motif, the NSE/DTE motif, was detected at amino acid position 461–467 (Figure 1). The less conserved motif NSE/DTE, apparently evolved from a second motif that was conserved in prenyl transferase. In general, these motifs were located on the opposite sides of the active site [6,7]. The metal binding residues appeared as NSE in most microbial and fungal cyclases and as DTE in most plant cyclases.

**Table 1.** BLASTx analysis of *PmSTPS1* with the NCBI protein database.

Description	Organism	Score	E-value	Identity (%)	Accession
Drimenol synthase	<i>Persicaria hydropiper</i>	678	0.0	96	AHF2284.1
Putative sesquiterpene synthase	<i>Persicaria minor</i>	664	0.0	95	ALO69830.1
Delta-cadinene synthase isozyme A	<i>Theobroma cacao</i>	393	$5 \times 10^{-129}$	54	XP_007021123.1
Predicted: probable sesquiterpene synthase	<i>Beta vulgaris</i> subsp. <i>vulgaris</i>	389	$3 \times 10^{-127}$	54	XP_010694277.1
Predicted: probable terpene synthase 2	<i>Rocinus communis</i>	381	$3 \times 10^{-124}$	52	XP_002523635.1

Table 1. Cont.

Description	Organism	Score	E-value	Identity (%)	Accession
Probable sesquiterpene synthase	<i>Santalum murrayanum</i>	379	$1 \times 10^{-123}$	54	F6M8H1
(-)-germacrene D synthase-like isoform X2	<i>Citrus sinensis</i>	379	$1 \times 10^{-123}$	52	XP_015384843.1

Table 2. BLASTx analysis of *PmSTPS2* with the NCBI protein database.

Description	Organism	Score	E-value	Identity (%)	Accession
Drimenol synthase	<i>Persicaria hydropiper</i>	492	$3 \times 10^{-164}$	46	AHF2284.1
Probable sesquiterpene synthase	<i>Beta vulgaris</i> subsp. <i>vulgaris</i>	492	$3 \times 10^{-164}$	43	XP_010694277.1
Valencene synthase-like	<i>Vitis vinifera</i>	489	$2 \times 10^{-163}$	44	NP_001268028.1
Germacrene A synthase	<i>Vitis vinifera</i>	489	$2 \times 10^{-163}$	44	ADR66821.1
Putative sesquiterpene synthase	<i>Persicaria minor</i>	484	$3 \times 10^{-161}$	45	ALO69830.1
Predicted: (-)-germacrene D synthase	<i>Vitis vinifera</i>	479	$4 \times 10^{-159}$	42	XP_002282488.1
(E)-beta-caryophyllene synthase	<i>Vitis vinifera</i>	477	$1 \times 10^{-158}$	45	ADR74194.1

```

MG921606 -----MSSQTVVLASPPVAAQTKFDVTRPIAKFHPCVWGNHFLNYTPSDEEV--VTERKM
XP_007021123.1 -----MSSLASSSLPVSTNDSMSNENRRSTKFPHSIWGDLFLSFPKVVN---DITTLQI
F6M8H7.1 -----MENQKMPISSSVFN--LKDLNMI SRPIANFPFSIWGDRFINYTCEDEN--DQTKQER
XP_002282488.1 -----MSVPLSVSVTPILSQRIDPEVARHEATYHPNFWGDRFLHYNPDDDFCGTHACKEQ
ADR74194.1 -----MSIQVSECP--VQIPKPEHRPMAEFHPSIWGDQFIAYTPEDEV--SRACKEK
NP_001268028.1 -----MSIQVSTCPL--VQIPKPEHRPMAEFHPSIWGDQFIAYTPEDED--TRACKEK
ADR66821.1 -----MSIQVSTCPL--VQIPKPEHRPMAEFHPSIWGDQFIAYTPEDED--TRACKEK
XP_010694277.1 -----MQTANSNCNRP LAHYHPSLWGDHFLNYTPPEDEL--TQRQRE
MG921605 -----
AHF22834.1 -----MSTAVNVPSAVRPADKRP IASFHPSFWGDYFLKYVPCDQV--TQAKMED
ALO69830.1 -----MSTAVNVPSAVRPADKRP IASFHPSFWGDYFLKYVPCDQV--TQAKMED

MG921606 ELESLEKLEVRKELKI--A-DRPLESMKLDIAIERLGVSYHFEKEIEESLQKAYDAYKKN
XP_007021123.1 HYEELKQEVRRMLRVP--TDNQLSQKRLRIETIKRLGVSYHFEREIEDVLQNIYDQ----
F6M8H7.1 QVEELKEQVRRELAAT--V-DKPLQQLNIIDATQRLGIAYLFENEIEESLKHIIYLHTYVE
XP_002282488.1 QIQELKBEVVRKSLAAT--A-GNPSQLLIDSDIQRGLAYHFEREIEEALKAMYQTYTL-
ADR74194.1 QVEDLKAQVRRRELMMAA--A-GNPSQLLNFIDAVQRLGVAYHFEREIEESLQHIYDRFHD-
NP_001268028.1 QVEDLKAQVRRRELMMAA--A-GNPSQLLNFIDAVQRLGVAYHFEREIEESLQHIYDRFHD-
ADR66821.1 QVEDLKAQVRRRELMMAA--A-GNPAQLLNFIDAVQRLGVAYHFEREIEESLQHIYDRFHD-
XP_010694277.1 EVKELKBEVVRKLLGFAEDR---KQRLVVIDVVEQLGVAYHFEABIEDNLQQFYNSYQD
MG921605 -----
AHF22834.1 EVKKEVEDVKKELRKLAKAVGKPLELNFIDVVERLGVGYRLEQBIEDLVQAI FDNKFG
ALO69830.1 EVKKEVEDVKKELRKLAKAVGKPLELNFIDVVERLGVGYRLEQBIEDLVQTI FDNKFG

MG921606 DQ--DSSLIDLYHTSLSFRILRQHGYVVS-SEIFSKFIDEHGSFIGSLKSDIMGLLSLYDA
XP_007021123.1 --DY-KDDNLEVTSLRFRLLRREHGFNVQ-CETFNKFKDDKGNFKVSLKSDVKGLLELYEA
F6M8H7.1 NNCFFGSDLYSVALWFRLLRQNGYKVS-CDVFNKFRDNEGNFKNNLMEDAKGILELYEA
XP_002282488.1 --VDD-NDHLTTVSLFRLLRQEGYHIP-SDVFKKFMDEGGNFKESLVGDLPGMLALYEA
ADR74194.1 --ADDTNDDLYNIALRFRLLRQGGYNIS-CGIFNKFKDEEGSFKEDLISNIQGMGLLYEA
NP_001268028.1 --ADDTEDDLYNIALQFRLLRQGGYNIS-CGIFNKFKDEKGSFKEDLISNIQGMGLLYEA
ADR66821.1 --ADDTEDDLYNIALQFRLLRQGGYNIS-CGIFNKFKDEKGSFKEDLISNVQGMGLLYEA
XP_010694277.1 DEI--YENDLHYVSLRFRLLRQHGFIAS-SDVYFKFNKTDGHFKESLKNVGLLNLYEA
MG921605 -----
AHF22834.1 --V--DEFDLYHTSLWFRLLRQHGFIAS-SDVYFKFNKTDGHFKESLKNVGLLNLYEA
ALO69830.1 --V--HEFDLYHTSLWFRLLRQHGFIAS-SDVYFKFNKTDGHFKESLKNVGLLNLYEA

MG921606 AYLRNGDITIMEKALEFSVVHLKVVAM-EKDHPLAPQICRALVYPIHKGVVLI ESRYIS
XP_007021123.1 AHIRTHGEHILEEALAFITTHLESAKTSLCEYPI SVLVSHARKRPIRKGLRLEARRFIS
F6M8H7.1 THVS IHGSEMLDDALEFTKTRLESVVS-HLNYPLAQVRRHALYQPLHRGLPRLEAVYFRR
XP_002282488.1 AHLMVHGEDILDEALGFTTAHLQSMAI-DSDNPLTKQVIRALKRPIRKGLRPRVEARHYIT
ADR74194.1 AHLRVHGEDILDEALAFITTHLAKAVE-SLGCPLAEQVAHALKRPPIRKGLRLEARRWYIS
NP_001268028.1 AHLRVHGEDILDEALAFITTHLAKATVE-SLGYPLAEQVAHALKRPPIRKGLRLEARRWYIS
ADR66821.1 AHLRVHGEDILDEALAFITTHLAKATVE-SLGYPLAEQVAHALKRPPIRKGLRLEARRWYIS
XP_010694277.1 SYLRVHGENILDEGVEFTTTQLKSLVSNL-SDPLATQVTQALKPLHKGVRRLLSRHYIS
MG921605 -----MPLRESRHIFIT
AHF22834.1 SHVRTHGDDTLDEALVFTTTHLKA VVTNQP NHPLVQVTHALMQPYHKGMPRLSRHFIA
ALO69830.1 SHVRTHGDDTLDEALVFTTTHLKA VVTNQP NHPLVQVTHALMQPYHKGMPRLSRHFIA
: : : : :
MG921606 FYEKQPFHMRSLRYSKLEFNVQRSLHKEELKDLTMWRSFEFPKTFPFI--RDRVVEAYL
XP_007021123.1 IYQEDGSHDKTLLKFAKLDVSLVQNLHKEELSKISRWWKDLDFKRLPFA-RDRIVEGYF
F6M8H7.1 IYEAHSHINKALLKAKLDENLQSFHKKELSDIARWWWKSLDFAAKPFA-RDRIVEGYF
XP_002282488.1 IYQEDDGHNESLKLAKLDYNNLQSLHKEELSEITKWWKGLDFATKLPFA-RDRIVEGYF
ADR74194.1 IYQDEAFHDKTLLKFAKLDVSLVQNLHKEELSNLARWWWKLDFAATKLPFA-RDRIVEGYF
NP_001268028.1 LYQDEASHDKTLLKFAKLDVSLVQNLHKEELSNLARWWWKLDFAATKLPFA-RDRIVEGYF
ADR66821.1 LYQDEASHDKTLLKFAKLDVSLVQNLHKEELSNLARWWWKLDFAATKLPFA-RDRIVEGYF
XP_010694277.1 TYEAHPLHDETLKFAKLDVSLVQNLHKEELSDIARWWWKSLDFAAKPFA-RDRIAEVYF
MG921605 FYEKDPRHDKTLLKFAKLDVSLVQNLHKEELKDKTRWWWKDLDMHAKMPPSRDRVPEGYF
AHF22834.1 FYEKDPYHDKTLLKFAKLDVSLVQNLHKEELKDLRWWWKDLDMHAKMPPSRDRVPEGYF
ALO69830.1 FYEKDPYHDKTLLKFAKLDVSLVQNLHKEELKDLRWWWKDLDMHAKMPPSRDRVPEGYF
* : . * . ** .** : . * * * : ** : : : * ** * :
RxR
MG921606 NSVSVFHDPKFSVARVYTKAFLLLTVDIYDSTYGTIEEALFTKAIERWDKSTIDELP

```

Figure 1. Cont.

```

XP_007021123.1 WILGVYFEPQYSFARQILTKAIVMASTMDDDIYDAYGTFEELQLFTKAIERWDNFCMDRLP
F6M8H7.1 WVLGVYFEPQYSLARKIIIKVFTMIISTIDDIYDAYGTLEDELKLFKAMQRWDVGSLLDQLP
XP_002282488.1 WILGVYFEPQYLLARRILMKVFGVLSIVDDIYDAYGTFEELKLFTEAIERWDASSIDQLP
ADR74194.1 WIVGVYFEPQYLWAIIRILTKIIVMTTVIDDIYDAYGTFEELIKHFTEAIERWDINSIDHLP
NP_001268028.1 WTLGVYFEPQYSRARRILTKLFMSASIIDDIYDAYGTLEELQPFTEAIERWDIKSIDHLP
ADR66821.1 WTLGVYFEPQYSRARRILTKLFAMASIIDDIYDAYGTLEELQPFTEAIERWDIKSIDHLP
XP_010694277.1 WMLGTFYEPQYALGREIFTKLYKMTSMVDDIYDAYGLFEELEVYTEAVQGWKSNMKNLPL
MG921605 WTLGPFYEPQFALCRKFFLQVFKVTSIVDDIYDAYGTIDELTFTKVVWRWDRSCLDQLP
AHF22834.1 WTLGPFYEPQFALCRKFFLQVFKVTSIVDDIYDAYGTIDELTFTKAAERWDRSCLDELPL
ALO69830.1 WTLGPFYEPQFALSRKFFLQVFKVTSIVDDIYDAYGTIDELTFTKAAERWDRSCLDELPL
      . . . : : * * * * * : : * * * * *
      DDxxD
MG921606 SYMKLFYEIQYDSYKQPEDDVAITLGIKNYVQYAVKEVITLSKAYLQEARWCKMKYVPTFD
XP_007021123.1 AYMKLFYKALLDVYEEEMVMTKQVKSIRVVKYAKEAMKQLSQAYFVEAKWYHENYVPTVE
F6M8H7.1 EYMKPCYKSIDLVYNEIEEEMANQSSLFMRHYAKEVMKTIVEGYMDEAKWCHEKYVPTFQ
XP_002282488.1 DYMKVCYQALLDVYEEEMEMTKQGLKLYRVHYAQAAALKRQVQAYLLEAKWIKQYIIPMD
ADR74194.1 EYMKLFYVALLDVYKEIEEEMEKERHQRVYHYADAMKNQVRAVFAEAKWHEGHIPTME
NP_001268028.1 EYMKLFYVTLDDLKYKIDQELEKYGNQYRVYVYAKEVLSQVRAVFAEAKWSHEGYIPTIE
ADR66821.1 EYMKLFYVTLDDLKYKIDQELEKYGNQYRVYVYAKEVLSQVRAVFAEAKWSHEGYIPTIE
XP_010694277.1 DYMKLYTEALLDTPDGFQSLANEGRSQLVSYVREMMKAQCRGYPQEAKWCHERYVPTFD
MG921605 DYMKLSYASLIDTFEEFERDLAPQGRSWSVKYAREEMIQMCRVVYQEAKWCHERYSPICD
AHF22834.1 EYMKVSYASLIDTFEEFERDLAPQGRSWSVKYAREEMIQMCRVVYQEAKWCHERYSPICD
ALO69830.1 DYMKVSYASLIDTFEEFERDLAPQGRSWSVKYAREEMIQMCRVVYQEAKWCHERYSPICD
      * * * * * : : . : : * * * * * : : * * * * *
      EYFQNA-IKSSATIMLVVVSLLFNVDETDAKKTFSVSKNSKPKQRATCIIGRMDLVLV
MG921606 EYMTNA-LVSSGYIMVTVTS--FVGMGNVVTKETPQWASNNPRIVRASSIITARINDDIVS
XP_007021123.1 EYMSVA-LVTSGYFLLTTS--YLGMEIASKAEFDLWFSHPVIEASESVGRINDDMRS
F6M8H7.1 EYMSNA-LVSACSMLETTTS--FVGMGDIVTKEAFDWFVSDPKMIRASNVICRIADDIVS
XP_002282488.1 EYMSVA-LVSSGYLLATSS--FVGMGIVSKEAFDVIIDPKIITASTVITARINDDMTS
ADR74194.1 EYMLVA-LVTSGSCILATWS--FVGMGEMTKEAFDVIIDPKIITASTVITARINDDMTS
NP_001268028.1 EYMLVA-LVTSGSCILATWS--FVGMGEMTKEAFDVIIDPKIITASTVITARINDDMTS
ADR66821.1 EYMLVA-LVTSGSCILATWS--FVGMGEMTKEAFDVIIDPKIITASTVITARINDDMTS
XP_010694277.1 EYLNDAAITTAGYTIMSGTT--FLGMGEFATKAVFEWARETPKPKASCIIIGRIIGMAS
MG921605 EYLEKASIVSFGYNLGTVVC--FLGMGDVATKEAFEWARGNPKVIRAACIIIGRIDDIGS
ADR66821.1 EYLEKASIVSFGYNLGTVVC--FLGMGDVATKEAFEWARGNPKVIRAACIIIGRIDDIGS
AHF22834.1 EYLEKASIVSFGYNLGTVVC--FLGMGDVATKEAFEWARGNPKVIRAACIIIGRIDDIGS
ALO69830.1 EYLEKASIVSFGYNLGTVVC--FLGMGDVATKEAFEWARGNPKVIRAACIIIGRIDDIGS
      * * * * * : : . : : * * * * * : : * * * * *
      NSE/DTE
MG921606 YEWKDRHRDHHVGGVDCYIKQHSVSEKAFEAINMMVENAWKEINEEFESTTTIAKPLLP
XP_007021123.1 HKFEQQRGHVSAIECYMKQHGVSSEKACNELNKQIENAWKIDINQELVRRPPAGVMPVLLT
F6M8H7.1 HKFEQQRGHVSAIECYMKQYGVTEEAHDEFKRRLVKAWKIDINEELRPP--YRVPKPLLT
XP_002282488.1 HEFEQQRGHVSAIECYMKQYGVSEKAADEFFKQVESAWKDNNEEFLQP--TAVVPVLLT
ADR74194.1 HKFEQQRGHVSAIECYMKQYGVSEKAADEFFKQVESAWKDNNEEFLQP--TAVVPVLLT
NP_001268028.1 HKFEQQRGHVSAIECYMKQYGVSEKAADEFFKQVESAWKDNNEEFLQP--TAVVPVLLT
ADR66821.1 HKFEQQRGHVSAIECYMKQYGVSEKAADEFFKQVESAWKDNNEEFLQP--TAVVPVLLT
XP_010694277.1 YKTERKREHVASAVECYMKQYGVSEKAADEFFKQVESAWKDNNEEFLQP--TAVVPVLLT
MG921605 HHFEQQRGHVSAIECYMKQYGVSEKAADEFFKQVESAWKDNNEEFLQP--TAVVPVLLT
AHF22834.1 HHFEQQRGHVSAIECYMKQYGVSEKAADEFFKQVESAWKDNNEEFLQP--TAVVPVLLT
ALO69830.1 HHFEQQRGHVSAIECYMKQYGVSEKAADEFFKQVESAWKDNNEEFLQP--TAVVPVLLT
      . . . * * * : : : * * * : : . : : * * * : : . : : * * *
      YIYNFCRIMDVYK--NENGYTIVSRMKDITKLVFIEPLSI*
MG921606 RILNLRVMDVLYK--EGDGYTHVGEAVKDGITSLMIDPIPI-
XP_007021123.1 RILNLRVMDVLYK--EGDGYTHVGEAVKDGITSLMIDPIPI-
F6M8H7.1 RILNLRVMDVLYK--EGDGYTHVGEAVKDGITSLMIDPIPI-
XP_002282488.1 RVLNLSRVMDVLYK--DEDEYTLVGLPLMKDLVAGLIDPVP-
ADR74194.1 RVLNLSRVMDVLYK--EGDGYTHVGEAVKDGITSLMIDPIPI-
NP_001268028.1 RVLNLSRVMDVLYK--EGDGYTHVGEAVKDGITSLMIDPIPI-
ADR66821.1 RVLNLSRVMDVLYK--EGDGYTHVGEAVKDGITSLMIDPIPI-
XP_010694277.1 RILNLSRVIEVYRLEGGDYSIVSQMVKVLELVLIHPVIV-
MG921605 RILNLSRVIEVYRLEGGDYSIVSQMVKVLELVLIHPVIV-
AHF22834.1 RILNLSRVIEVYRLEGGDYSIVSQMVKVLELVLIHPVIV-
ALO69830.1 RILNLSRVIEVYRLEGGDYSIVSQMVKVLELVLIHPVIV-
      : * * : : * * : : * * : : * * : : * * : : * *


```

**Figure 1.** Comparison of the deduced amino acid sequence of *PmSTPS1* and *PmSTPS2*, with other terpene synthase sequences of the highest sequence similarity. *Polygonum minus* sesquiterpene synthase 1 (MG921605); *Polygonum minus* sesquiterpene synthase 2 (MG921606); *Persicaria hydropiper* drimenol synthase (AHF22834.1); *Persicaria minor* putative sesquiterpene synthase (ALO69830.1); *Theobroma cacao* Delta-cadinene synthase isozyme A (XP\_00702113.1); *Beta vulgaris* subs. *vulgaris* probable sesquiterpene synthase (XP\_010694277.1); *Santalum murrayanum* probable sesquiterpene synthase (F6M8H1); *Vitis vinifera* (–)-Germacrene D synthase (XP\_002282488.1); *Vitis vinifera* Valencene synthase (NP\_001268028.1); and *Vitis vinifera* Germacrene A synthase (ADR61821.1). Amino acid residues conserved in all of the genes are marked with asterisk [\*]. Amino acid residues conserved in four or five genes are indicated by double dots [:]. The universally conserved DDxxD, RxR motifs, and NSE/DTE are highlighted in boxes.

2.2. Phylogenetic Analysis of *P. minus* Sesquiterpene Synthase (*PmSTPS*)

The *PmSTPS1* and *PmSTPS2* amino acid sequences were aligned and compared with other flowering plant terpene synthase sequences, using Clustal Omega (Figure 2), and they showed a low sequence similarity (42.94%). The phylogenetic analysis showed a particularly close relationship between the *PmSTPS1* and *PmSTPS2* amino acid sequences. The *PmSTPS1* was clustered in the same clade with sesquiterpene synthase (*PmSTS*) from *Persicaria minor* and drimenol synthase from *Persicaria hydropiper*. The results showed that the

*PmSTPS1* from *P. minus* was grouped into a single clade with a 43.04% identity, which suggested a monophyletic origin of the gene. Additionally, *PmSTPS2* was placed in the same clade with (+) delta-cadinene synthase from *Ricinus communis*. Moreover, *PmSTPS1* and *PmSTPS2* were grouped together with the terpene synthases from the *Santalum* and *Vitis vinifera* species. Multiple sequence alignments of *PmSTPS1* and *PmSTPS2* amino acid sequences, with sesquiterpenes from other plants species, showed a high sequence similarity (42–96%).



**Figure 2.** Phylogenetic tree of *PmSTPS1* and *PmSTPS2* protein sequences with amino acid sequences, with selected terpene synthases from other plants. The alignment was performed using the Clustal Omega algorithm. The tree was built using the neighbor-joining method and 1000 replicates for bootstrapping. The numbers indicated are the actual bootstrap values of the branches.

### 2.3. Expression of *PmSTPS1* and *PmSTPS2* in *E. coli*

For the analysis of the protein expression, recombinant bacterial strains harbouring pQE2 in *E. coli* M15 with *PmSTPS1* and *PmSTPS2* were compared with those harbouring the control empty pQE2 vector. The cells were harvested at different times (1, 3, and 5 h) post-induction. After sonication and centrifugation of the bacteria, soluble and insoluble crude fractions were separated with 10% SDS-PAGE. The SDS-PAGE analysis (Figure S3) showed unclear corresponding protein bands at the expected size at different post-induction times, as well as in the control sample. However, Western Blotting (Figure S4) confirmed the correct size of recombinant proteins. There was no band was observed in the control sample as expected. Correct protein sizes of 40.9 and 65.9 kDa were obtained for *PmSTPS1* and *PmSTPS2*, respectively. From these findings, the recombinant *PmSTPS1* and *PmSTPS2* proteins from *P. minus* were successfully expressed in *E. coli*, and the activities of these enzymes were further investigated via enzymatic assays.

### 2.4. Identification of *PmSTPS1* and *PmSTPS2* Assay Products

A functional characterisation of the *PmSTPS1* and *PmSTPS2* genes was performed by an in vitro enzyme assay of the recombinant proteins. In this crude protein assay, the *E. coli* strain harbouring empty pQE-2 vector was used as the control strain. A GC-MS analysis showed that *PmSTPS1* and *PmSTPS2* produced  $\beta$ -farnesene,  $\alpha$ -farnesene, and farnesol as the final products. Additionally, *PmSTPS2* also produced nerolidol. For *PmSTPS1*, the products formed were  $\beta$ -farnesene (14.49 min),  $\alpha$ -farnesene (15.79 min), and farnesol (16.98 min). Additionally, the principle products from *PmSTPS2* enzyme were  $\beta$ -farnesene (14.49 min),  $\alpha$ -farnesene (15.83 min), farnesol (16.84 min), and nerolidol (17.25 min). Based on the GC-MS analysis, both extracts from *PmSTPS1* and *PmSTPS2* showed multiple peaks for corresponding sesquiterpene products, compared with no peaks observed in the control sample, which did not exhibit any major products, although exogenous substrates were added (Figure 3). These findings therefore demonstrated the successful production of sesquiterpenes in the recombinant *E. coli* strains overexpressing *PmSTPS1* and *PmSTPS2* enzymes, respectively.





Figure 3. Cont.



**Figure 3.** GC-MS chromatogram of products formed by crude *PmSTPS1* and *PmSTPS2* protein at different retention times (RT). (a) The chromatogram of control used consisted of M15 *E. coli* host harboring empty pQE-2 taqzyme plasmid; (b) *PmSTPS1* (c); and *PmSTPS2*.

All of the peaks were further confirmed by comparison with NIST and Wiley libraries, mass spectra, authentic sesquiterpene standards, and control (Figure S5). Interestingly, although the sizes of *PmSTPS1* and *PmSTPS2* were different, the two enzymes were capable of producing similar sesquiterpene products,  $\beta$ -farnesene,  $\alpha$ -farnesene, and farnesol (Figure 3), but they did so at different levels. *PmSTPS1* successfully converted the precursor FPP to produce 9.50%  $\beta$ -farnesene as the main product, followed by 8.86%  $\alpha$ -farnesene, and 5.08% farnesol. *PmSTPS2* showed syntheses nerolidol (48.33%) as a major product, followed by farnesol (15.30%),  $\beta$ -farnesene (5.07%), and  $\alpha$ -farnesene (2.76%). Although the (E,E)-farnesyl pyrophosphate (FPP) substrate was added to the enzymatic assay, the control pQE-2 sample did not produce any significant products, indicating that endogenous metabolites did not affect the protein expression and analysis in this study.

### 3. Discussion

In this study, we provided the first cloning and functional characterisation of *PmSTPS1* encoding a putative  $\beta$ -farnesene synthase, and *PmSTPS2* encoding a putative nerolidol synthase in *P. minus*. A sequence comparison between the *PmSTPS1* and *PmSTPS2* indicated that the two enzymes had different protein and nucleotide sequences. However, both of the enzymes were structurally similar to other plant sesquiterpene synthases and contained all of the conserved motifs, including DDxxD, RXR, and NSE/EDTA, which were important for terpene synthase functionality [5,45,46]. Based on the phylogenetic analysis, *PmSTPS1* and *PmSTPS2* clustered on the same group with four distinct sesquiterpene synthases.

In addition, the unexpected band in the upper layer (Figure S4) might have been caused by protease aggregation, according [47]. As shown in several recent studies, the protein sizes of a few plants sesquiterpene synthases showed a nearly similar molecular weight, with *PmSTPS2* within the range of 60–70 kDa [24,48,49]. In general, terpene synthase could be classified into monoterpene, sesquiterpene, and diterpene synthase, with 550–860 amino acids encoding a 50–100 kDa protein [4,50]. As a result of the absence of the signal peptide sequence with 50–70 amino acids, the size of sesquiterpene synthases were typically smaller than those of the monoterpenes and diterpenes [51]. Ee et al. [37] also reported that the protein size of the *P. minus*  $\beta$ -sesquiphelandrene synthase was 65.1 kDa. Additionally, studies on  $\beta$ -caryophyllene synthase, that were encoded by OkBCS (GenBank accession no. KP226502) from *Ocimum kilimandscharicum* Gürke, showed a molecular weight of 63.6 kDa [24]. Until recently, no short sesquiterpene synthase sequence was characterised as a *PmSTPS1*. However, the short-chain length of this enzyme could be associated with several prenyltransferase (PT) enzymes. Based on previous findings, two prenyltransferases, Santalum farnesyl diphosphate synthase (SaFDS) and Hedychium farnesyl pyrophosphate synthase (HcFPPs), comprising 1029 and 1068 bp nucleotide sequences and encoding polypeptides of 343 and 356 amino acids, respectively, were reported [3,52].

Many terpene synthases (TPSs) had the ability to synthesise one or multiple products from a single substrate, regardless of whether it was farnesyl pyrophosphate (FPP) or geranyl pyrophosphate (GPP) [53,54]. In addition, the sesquiterpene synthases from different plant species produced more than one product [25,55,56]. Interestingly, although the sizes of *PmSTPS1* and *PmSTPS2* were different, the two enzymes produced similar sesquiterpene products ( $\beta$ -farnesene,  $\alpha$ -farnesene, and farnesol, Figure 3), albeit at different percentages. Both *PmSTPS1* and *PmSTPS2* could catalyse the formation of  $\beta$ -farnesene,  $\alpha$ -farnesene, nerolidol, and farnesol, whose functions were different from those of the previous STPS enzymes characterised in *P. minus* [36–39]. In addition, the enzymes demonstrated an inherent capacity for TPS enzymes to evolve different products and substrate specificities [57]. Moreover, the main factor of sesquiterpene diversity was the large number of different sesquiterpene synthases expressed in plants and the ability of some sesquiterpene synthases to form multiple products from a single FPP substrate [58].

Based on previous chemical profiling studies of *P. minus* essential oils from hydro-distillation extraction, low levels of nerolidol and farnesol were detected at 0.24% and 0.14%, respectively [34]. A similar finding was reported, which indicated that the percentage of  $\beta$ -farnesene and  $\alpha$ -farnesene compounds were also found at 0.92% and 0.82%, respectively [31,32,34]. The percentage of terpenes that were obtained directly from the GC-MS analysis of *P. minus* leaf essential oil was lower compared with the products that were produced by the enzymatic assay of crude protein *PmSTPS1* and *PmSTPS2*. The variation in composition could have been because of the variable amounts of sesquiterpenes that were produced in the plants, depending on environmental factors. Nevertheless, the sesquiterpenoids that were produced by in vitro assay potentially contributed to the plant fragrance, because most of the acyclic sesquiterpenes compounds, namely, farnesene, nerolidol, and farnesol, were previously reported in various plant essential oils [21,59,60]. These compounds could be potentially commercialised as fragrances, flavouring agents, or pharmaceutical products. In addition, farnesene is an important compound for diesel and jet fuels [61]. Understanding the physiological and ecological roles of plant volatile sesquiterpenes has been challenging. Several sesquiterpenes compounds might have acted as defense chemicals against biological stresses. For instance, E,E- $\alpha$ -farnesene was reported to have potential for use as an alarm pheromone in the control of aphid pests [62]. Furthermore, the existence of farnesol in *P. minus* essential oil could have been related to the biosynthetic pathway of juvenile hormone (JH) III [63]. Nerolidol was not only used in cosmetics and non-cosmetic products [64], but was also proven to possess pharmacological and biological activities [65,66]. Therefore, the advantages of nerolidol have made it a promising drug candidate for industrial production [67,68].

## 4. Materials and Methods

### 4.1. Plant Material

*P. minus* plants were grown in an experimental plot at Universiti Kebangsaan Malaysia (UKM) under natural light and environmental conditions. The samples were originally collected from Ulu Yam, Selangor, Malaysia (UY; 3°16'14.63" N, 101°41'11.32" E), and were identified using ITS sequences [69]. The voucher specimens were deposited at the UKM herbarium. Leaf samples from *P. minus* plants were harvested in the morning, between 8 to 9 am, frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$  for RNA extraction.

### 4.2. RNA Isolation and cDNA Synthesis

Total RNA was isolated and extracted as it was previously reported [70]. The quantity, purity, and integrity of the RNA were determined using standard methods. Three micrograms of RNA were reverse transcribed into cDNA using the Onetaq<sup>®</sup> One-step RT-PCR kit (New England Biolabs, Ipswich, MA, USA), according to manufacturer's instructions.

#### 4.3. Candidate Gene Selection and Isolation of Full-Length *PmSTPS1* and *PmSTPS2*

The candidate gene selection was achieved by mining the *P. minus* transcriptome data [40] for transcripts that were related to the sesquiterpene biosynthetic pathway. The assembled transcripts were classified as sesquiterpene synthase, based on homology search, and the terpene synthases were selected and fully sequenced prior to further analysis. Two new *P. minus* sesquiterpene synthase (*PmSTPS*) candidate genes were identified. The predicted ORFs for *PmSTPS1* (GenBank accession no: MG921605) and *PmSTPS2* (GenBank accession no: MG921606) were amplified by PCR, using the Q5 High-Fidelity DNA Polymerase (NewEngland Biolabs, Ipswich, MA, USA). The cDNA-gene specific PCR primers were *PmSTPS1\_F* (5'-AAAGGTACCATGCCAA GGCTCG-3') and *PmSTPS1\_R* (5'-TTTGTCGACAATCGGAATGGGAT-3'); *PmSTPS2\_F* (5'-AAAGGTACCATGTCAT

CCCCAAA-3'), and *PmSTPS2\_R* (5'-TTTAAGCTTAATGGAGAGAGGTT-3') were synthesized to amplify the 5' end and 3' end, respectively.

The PCR reaction mixture contained 1× reaction buffer, 2 mM MgCl<sub>2</sub>, 0.2 mM dNTPs, 5 units of Taq polymerase (Promega, Madison, WI, USA), 0.5 μM of forward and reverse primers, and 20 ng of template cDNA. The reaction was performed under the following conditions: pre-denaturation at 98 °C for 30 s, followed by 32 cycles of 98 °C for 10 s, 60 °C for 10 s, and 72 °C for 20 s, with a final extension at 72 °C for 2 min. The amplicons were digested with *KpnI/SalI* and *KpnI/HindIII* before being cloned into pUC19 and pUC57-Kan cloning vectors, respectively. The ligation mixtures were transformed into *E. coli* top 10 competent cells, before sub-cloning into the pQE-2 plasmid. The positive transformants were screened on LB agar that were supplemented with 100 μg/mL ampicillin, 20 μg/mL X-gal, and 0.1 mM IPTG (ThermoFisher Scientific, Waltham, MA, USA). The recombinant plasmids pQE-2:*PmSTPS1* and pQE-2:*PmSTPS2* were then transformed into the *E. coli* M15 competent cells and the transformants were selected on LB agar that was supplemented with 50 μg/mL kanamycin. The positive transformants were confirmed by colony PCR, and the gene sequences were verified via DNA sequencing (First BASE Laboratories, Seri Kembangan, Selangor, Malaysia).

#### 4.4. Full-Length cDNA Sequence Analysis and Phylogenetic Tree Construction

The ORF for *PmSTPS1* and *PmSTPS2* were predicted using the ORF finder program (<http://www.ncbi.nlm.nih.gov>) and were subjected to BLASTX and BLASTP analyses. Multiple sequence alignment was achieved using the Clustal Omega pairwise alignment algorithm. Verification of the cDNA sequence, including the amino acid sequence, theoretical isoelectric point (pI), and predicted molecular weight (MW) of the analyses, was performed using ExPASy Proteomic tools (<http://www.cn.expasy.org/tools/protscale.html>). The physical and chemical characteristics of all of the deduced amino acid sequences were analysed by the ProtParam tool (<http://web.expasy.org/program/>). The signal peptide targeting location of the deduced proteins was predicted using the SignalP method (<http://www.cbs.dtu.dk/services/SignalP>) and ChloroP program (<http://www.cbs.dtu.dk/services/ChloroP/>). A protein domain analysis was performed using the SMART (Simple Modular Architectural Research Tool) database (<http://smart.embl-heidelberg.de/>).

#### 4.5. Phylogenetic Analysis

Phylogenetic and molecular evolutionary analyses of the amino acid sequences of the *PmSTPS1* and *PmSTPS2* from different plant species were constructed using the default parameters of PhyML software, which were available at Phylogeny.fr web services ([www.phylogeny.fr/version2/cgi/simplephylogeny.cgi](http://www.phylogeny.fr/version2/cgi/simplephylogeny.cgi)) [71]. PhyML was employed to construct a phylogenetic tree, by generating multiple alignments through the neighbour-joining computational method.

#### 4.6. Expression of *PmSTPS1* and *PmSTPS2* in *E. coli*

A single colony of recombinant *E. coli* M15 cells harbouring pQE-2:*PmSTPS1*, pQE2:*PmSTPS2*, and empty pQE-2 (as a negative control) were inoculated into 10 mL of an LB medium containing

kanamycin (50 µg/mL), and were grown overnight at 37 °C. Approximately 2 mL of the cultures were added to 200 mL of fresh LB, which contained 50 µg/mL kanamycin. The cultures were induced with 0.5 mM IPTG at OD<sub>600</sub> ~0.5. The cultures were incubated for 1, 3, and 5 h at 37 °C, and were then harvested by centrifugation at 4000× g for 30 min at 4 °C. Subsequently, the bacteria were resuspended in 100 mL of 25 mM sodium phosphate buffer, pH 7.5, containing 0.5 M Tris-HCl, 5% glycerol, 1 mM dithiothreitol (DTT), 10 mM MgCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>, pH 7.5, and 1 mM lysozyme (Sigma-Aldrich, St. Louis, MI, USA) [20]. The cells were sonicated for 2 min at 5 s pulses, with 5 s between the pulses on ice, using the Sonic Dismembrator Model 100 (Fisher Scientific, Hampton, NH, USA). The cell lysate was then centrifuged at 10,000× g for 30 min at 4 °C.

#### 4.7. Enzyme Assay

A standard assay was done according to a previous method, with slight modifications. Standard assays were performed in 2.5 mL glass GC vials containing 200 µg of crude protein mixed with 50 mM Tris (pH 7.5), 10 mM MgCl<sub>2</sub>, and 100 µM of (E,E)-farnesyl pyrophosphate (FPP) (Sigma-Aldrich, St. Louis, MI, USA). The reaction mixture with a total volume of 200 µL was vortexed, overlaid with 500 µL hexane, and incubated at 30 °C for 2 h. The hexane phase was concentrated to 200 µL by passing N<sub>2</sub> at the opening of the tube and was then further used for the GC-MS analysis.

#### 4.8. Detection of Sesquiterpenes Using GC-MS

The samples were analysed using a Clarus 600 GC-MS (PerkinElmer Inc., Waltham, MA, USA) that was equipped with a capillary column (Elite-5 30 m × 0.25 mm, film thickness 0.25 µm). The GC was operated at a flow rate of 2 mL/min, and the mass selector detector (MSD) was operated at 70 eV. Splitless injections (1.5 µL) were performed with an injector temperature of 250 °C. The GC system was programmed with an initial oven temperature of 50 °C (5 min hold), which was then increased to 180 °C at 10 °C/min (4 min hold), followed by a 100 °C/min ramp at 240 °C (1 min hold). A solvent delay of 8.5 min was allowed before the acquisition of the MS data. The MS system was operated in selected ion monitoring (SIM) mode to scan for the molecular ions at product peaks, which were quantified by the integration of peak areas with library search, using the NIST library [72].

### 5. Conclusions

In summary, two new sesquiterpene synthases, *PmSTPS1* and *PmSTPS2*, which were identified from *P. minus* leaf transcriptomics analysis, were cloned and characterised. Both of the enzymes produced industrially important acyclic sesquiterpenes, β-farnesene, α-farnesene, and farnesol. *PmSTPS2* also produced nerolidol as the major product from FPP conversion. This study demonstrated the production of *P. minus* characteristic fragrance-related sesquiterpenes, by both *PmSTPS1* and *PmSTP*, as well as the potential of further metabolic engineering in *E. coli*, using *PmSTPS2* for the microbial production of nerolidol.

**Supplementary Materials:** The following are available online, Figure S1: Nucleotide and predicted amino acid sequence of *PmSTPS1* and from *P. minus*; Figure S2: Nucleotide and predicted amino acid sequence of *PmSTPS2* from *P. minus*; Figure S3: SDS-PAGE analysis of recombinant pQE2-*PmSTPS1* and pQE2-*PmSTPS2* proteins is marked with red box and molecular mass markers are indicated; Figure S4: The expression analysis of *PmSTPS1* and *PmSTPS2* in *E. coli* M15 after 0.5 mM IPTG induction at 1, 3, and 5 h; and Figure S5: Mass spectra of major three sesquiterpenes produced by recombinant *PmSTPS1* and *PmSTPS2* in comparison with the mass spectra from authentic standards.

**Author Contributions:** N.A.R. and S.N.B. conceived and designed all of the experiments; N.A.R. performed the experiments and analysed the data; S.N.B. and H-H.G. contributed reagents/materials/analysis tools; N.A.R., S.N.B., H-H.G., S.S., A.B.R., and N.M.N. wrote and edited the paper. All of the authors reviewed the manuscript.

**Funding:** This research was funded by Research University Grants from Universiti Kebangsaan Malaysia [UKM-GUP-2013-028 and DLP 2013-024] awarded to Syarul Nataqain Baharum. We also acknowledge the Centre for Research and Instrumentation Management (CRIM), UKM for a Research Instrumentation Development Grant 2010 (PIP-CRIM). The publication fee was supported by DPP-2018-010 awarded to INBIOSIS, UKM.

**Acknowledgments:** The authors thank Kamarul Azlan Azizan and Syahmi Afiq Mustaza for their technical assistance and the reviewers for constructive comments.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. McCaskill, D.R.C. Prospects for the Bioengineering of Isoprenoid Biosynthesis. In *Biotechnology of Aroma Compounds. Advances in Biochemical Engineering/Biotechnology*; Springer: Berlin/Heidelberg, Germany, 1997; pp. 107–146.
2. Lange, B.M.; Ahkami, A. Metabolic Engineering of Plant Monoterpenes, Sesquiterpenes and Diterpenes—Current Status and Future Opportunities. *Plant Biotechnol. J.* **2013**, *11*, 169–196. [[CrossRef](#)] [[PubMed](#)]
3. Lan, J.B.; Yu, R.C.; Yu, Y.Y.; Fan, Y.P. Molecular Cloning and Expression of *Hedychium coronarium* Farnesyl Pyrophosphate Synthase Gene and Its Possible Involvement in the Biosynthesis of Floral and Wounding/herbivory Induced Leaf Volatile Sesquiterpenoids. *Gene* **2013**, *518*, 360–367. [[CrossRef](#)] [[PubMed](#)]
4. Bohlmann, J.; Meyer-Gauen, G.; Croteau, R. Plant Terpenoid Synthases: Molecular Biology and Phylogenetic Analysis. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 4126–4133. [[CrossRef](#)] [[PubMed](#)]
5. Degenhardt, J.; Köllner, T.G.; Gershenzon, J. Monoterpene and Sesquiterpene Synthases and the Origin of Terpene Skeletal Diversity in Plants. *Phytochemistry* **2009**, *70*, 1621–1637. [[CrossRef](#)] [[PubMed](#)]
6. Tholl, D.; Lee, S. Terpene Specialized Metabolism in *Arabidopsis thaliana*. *Arabidopsis Book* **2011**, *9*, e0143. [[CrossRef](#)] [[PubMed](#)]
7. Christianson, D.W. Structural and Chemical Biology of Terpenoid Cyclases. *Chem. Rev.* **2017**, *117*, 11570–11648. [[CrossRef](#)] [[PubMed](#)]
8. Kuzuyama, T. Mevalonate and Nonmevalonate Pathways for the Biosynthesis of Isoprene Units. *Biosci. Biotechnol. Biochem.* **2002**, *66*, 1619–1627. [[CrossRef](#)] [[PubMed](#)]
9. Eisenreich, W.; Bacher, A.; Arigoni, D.; Rohdich, F. Biosynthesis of Isoprenoids via the Non-Mevalonate Pathway. *Cell. Mol. Life Sci.* **2004**, *61*, 1401–1426. [[CrossRef](#)] [[PubMed](#)]
10. Goldstein, J.L.; Brown, M.S. Regulation of the Mevalonate Pathway. *Nature* **1993**, *295*, 517–524. [[CrossRef](#)] [[PubMed](#)]
11. Rohmer, M. The Mevalonate-Independent Methylerythritol 4-Phosphate (MEP) Pathway for Isoprenoid Biosynthesis, Including Carotenoids. *Pure Appl. Chem.* **1999**, *71*, 2279–2284. [[CrossRef](#)]
12. Chandran, S.S.; Kealey, J.T.; Reeves, C.D. Microbial Production of Isoprenoids. *Process Biochem.* **2011**, *46*, 1703–1710. [[CrossRef](#)]
13. Keasling, J.D. Synthetic Biology and the Development of Tools for Metabolic Engineering. *Metab. Eng.* **2012**, *14*, 189–195. [[CrossRef](#)] [[PubMed](#)]
14. Peralta-Yahya, P.P.; Ouellet, M.; Chan, R.; Mukhopadhyay, A.; Keasling, J.D.; Lee, T.S. Identification and Microbial Production of a Terpene-Based Advanced Biofuel. *Nat. Commun.* **2011**, *2*, 483. [[CrossRef](#)] [[PubMed](#)]
15. Schwab, W.; Davidovich-Rikanati, R.; Lewinsohn, E. Biosynthesis of Plant-Derived Flavor Compounds. *Plant J.* **2008**, *54*, 712–732. [[CrossRef](#)] [[PubMed](#)]
16. Srivastava, P.L.; Daramwar, P.P.; Krithika, R.; Pandreka, A.; Shankar, S.S.; Thulasiram, H.V. Functional Characterization of Novel Sesquiterpene Synthases from Indian *Sandalwood*, *Santalum album*. *Sci. Rep.* **2015**, *5*, 10095. [[CrossRef](#)] [[PubMed](#)]
17. Paddon, C.J.; Keasling, J.D. Semi-Synthetic Artemisinin: A Model for the use of Synthetic Biology in Pharmaceutical Development. *Nat. Rev. Microbiol.* **2014**, *12*, 355. [[CrossRef](#)] [[PubMed](#)]
18. Aschenbrenner, A.-K.; Kwon, M.; Conrad, J.; Ro, D.-K.; Spring, O. Identification and Characterization of Two Bisabolene Synthases from Linear Glandular Trichomes of Sunflower (*Helianthus annuus* L., Asteraceae). *Phytochemistry* **2016**, *124*, 29–37. [[CrossRef](#)] [[PubMed](#)]
19. Pazouki, L.; Memari, H.R.; Kännaste, A.; Bichele, R.; Niinemets, Ü. Germacrene A Synthase in Yarrow (*Achillea millefolium*) is an Enzyme with Mixed Substrate Specificity: Gene Cloning, Functional Characterization and Expression Analysis. *Front. Plant Sci.* **2015**, *6*, 111. [[CrossRef](#)] [[PubMed](#)]
20. Kwon, M.; Cochrane, S.A.; Vederas, J.C.; Ro, D.K. Molecular Cloning and Characterization of Drimenol Synthase from Valerian Plant (*Valeriana officinalis*). *FEBS Lett.* **2014**, *588*, 4597–4603. [[CrossRef](#)] [[PubMed](#)]

21. Parveen, I.; Wang, M.; Zhao, J.; Chittiboyina, A.G.; Tabanca, N.; Ali, A.; Baerson, S.R.; Techen, N.; Chappell, J.; Khan, I.A.; et al. Investigating Sesquiterpene Biosynthesis in *Ginkgo biloba*: Molecular Cloning and Functional Characterization of (E,E)-Farnesol and  $\alpha$ -Bisabolene Synthases. *Plant Mol. Biol.* **2015**, *89*, 451–462. [[CrossRef](#)] [[PubMed](#)]
22. Yu, F.; Okamoto, S.; Nakasone, K.; Adachi, K.; Matsuda, S.; Harada, H.; Misawa, N.; Utsumi, R. Molecular Cloning and Functional Characterization of  $\alpha$ -Humulene Synthase, a Possible Key Enzyme of Zerumbone Biosynthesis in Shampoo Ginger (*Zingiber zerumbet* Smith). *Planta* **2008**, *227*, 1291–1299. [[CrossRef](#)] [[PubMed](#)]
23. Attia, M.; Kim, S.; Ro, D. Catalyzing the First Step in the Biosynthesis of the Natural Sweetener, Hernandulcin, in *Lippia dulcis*. *Arch. Biochem. Biophys.* **2012**, *527*, 37–41. [[CrossRef](#)] [[PubMed](#)]
24. Jayaramaiah, R.H.; Anand, A.; Beedkar, S.D.; Dholakia, B.B.; Puneekar, S.A.; Kalunke, R.M.; Gade, W.N.; Thulasiram, H.V.; Giri, A.P. Functional Characterization and Transient Expression Manipulation of a New Sesquiterpene Synthase Involved in  $\beta$ -Caryophyllene Accumulation in *Ocimum*. *Biochem. Biophys. Res. Commun.* **2016**, *473*, 265–271. [[CrossRef](#)] [[PubMed](#)]
25. König, J.C.; Scheper, T.; Beutel, S. Heterologous Expression, Purification, and Biochemical Characterization of  $\alpha$ -Humulene Synthase from *Zingiber zerumbet* Smith. *Appl. Biochem. Biotechnol.* **2016**, *178*, 474–489.
26. Picaud, S.; Olsson, M.E.; Brodelius, M.; Brodelius, P.E. Cloning, Expression, Puri W Cation and Characterization of Recombinant (+)-Germacrene D Synthase from *Zingiber officinale*. *Arch. Biochem. Biophys.* **2006**, *452*, 17–28. [[CrossRef](#)] [[PubMed](#)]
27. Abubakar, M.A.; Zulkifli, R.M.; Nur, W.; Wan, A.; Husni, A.; Shariff, M.; Ahmad, N.; Nik, N.; Zakaria, Z.; Ahmad, F. Antibacterial Properties of *Persicaria minor* (Huds.) Ethanolic and Aqueous-Ethanolic Leaf Extracts. *J. Appl. Pharm. Sci.* **2015**, *5* (Suppl. 2), 50–56.
28. Christopher, P.V.; Parasuraman, S.; Christina, J.M.A.; Asmawi, M.Z.; Vikneswaran, M. Review on *Polygonum minus*. Huds, a Commonly Used Food Additive in Southeast Asia. *Pharmacogn. Res.* **2014**, *7*, 1. [[CrossRef](#)] [[PubMed](#)]
29. George, A.; Chinnappan, S.; Choudhary, Y.; Bommu, P.; Sridhar, M. Immunomodulatory Activity of an Aqueous Extract of *Polygonum minus* Huds on Swiss Albino Mice Using Carbon Clearance Assay. *Asian Pac. J. Trop. Dis.* **2014**, *4*, 398–400. [[CrossRef](#)]
30. Qader, S.W.; Abdulla, M.A.; Chua, L.S. Potential Bioactive Property of *Polygonum minus* Huds (Kesum) Review. *Sci. Res. Essays* **2012**, *7*, 90–93.
31. Ahmad, R.; Baharum, S.N.; Bunawan, H.; Lee, M.; Noor, N.M.; Rohani, E.R.; Ilias, N.; Zin, N.M. Volatile Profiling of Aromatic Traditional Medicinal Plant, *Polygonum minus* in Different Tissues and Its Biological Activities. *Molecules* **2014**, *19*, 19220–19242. [[CrossRef](#)] [[PubMed](#)]
32. Baharum, S.N.; Bunawan, H.; Ghani, A.; Aida, W.; Mustapha, W.; Noor, N.M. Analysis of the Chemical Composition of the Essential Oil of *Polygonum minus* Huds. Using Two-Dimensional Gas Chromatography-Time-of-Flight Mass Spectrometry (GC-TOF MS). *Molecules*. [[CrossRef](#)] [[PubMed](#)]
33. Yaacob, K.B.; Noor Salleh, A.; Joulain, D. Studies on *Polygonum* Spp. In Proceedings of the 12th International Congress of Flavours Fragrances and Essential Oils, Vienna, Austria, 4–8 October 1992; pp. 491–498.
34. Rusdi, N.A.; Goh, H.H.; Baharum, S.N. GC-MS/Olfactometric Characterisation and Aroma Extraction Dilution Analysis of Aroma Active Compounds in *Polygonum minus* Essential Oil. *Plant Omics* **2016**, *9*, 289–294. [[CrossRef](#)]
35. Tan, E.; Othman, R. Characterization of  $\alpha$ -Farnesene Synthase Gene from *Polygonum minus*. *Trans. Malaysian Soc. Plant Physiol.* **2012**, *20*, 155–157.
36. Song, A.A.-L.; Abdullah, J.O.; Abdullah, M.P.; Shafee, N.; Othman, R.; Tan, E.-F.; Noor, N.M.; Raha, A.R. Overexpressing 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase (HMGR) in the Lactococcal Mevalonate Pathway for Heterologous Plant Sesquiterpene Production. *PLoS ONE* **2012**, *7*, e52444. [[CrossRef](#)] [[PubMed](#)]
37. Ee, S.; Othman, R.; Shaharuddin, N.A.; Ismail, I.; Zainal, Z. Functional Characterization of Sesquiterpene Synthase from *Polygonum minus*. *Sci. World J.* **2014**, *2014*. [[CrossRef](#)] [[PubMed](#)]
38. Kiat, C.J.; Ashraf, M.F.; Naeem-Ul-Hassan, M.; Che Mohd Zain, C.R.; Zainal, Z.; Ismail, I. Molecular Cloning, Characterization and Expression Pattern Analysis of a Jasmonic Acid Responsive Sesquiterpene Synthase Gene from *Persicaria minor*. *Plant Omics* **2016**, *9*, 360–368. [[CrossRef](#)]
39. Ker, D.-S.; Pang, S.L.; Othman, N.F.; Kumaran, S.; Tan, E.F.; Krishnan, T.; Chan, K.G.; Othman, R.; Hassan, M.; Ng, C.L. Purification and Biochemical Characterization of Recombinant *Persicaria Minor*  $\beta$ -Sesquiphellandrene Synthase. *PeerJ* **2017**, *5*, e2961. [[CrossRef](#)] [[PubMed](#)]

40. Loke, K.; Rahnamaie-tajadod, R.; Yeoh, C.; Goh, H.; Mohamed-Hussein, Z.; Mohd, N.; Zainal, Z.; Ismail, I. Genomics Data RNA-Seq Analysis for Secondary Metabolite Pathway Gene Discovery in *Polygonum minus*. *GDATA* **2016**, *7*, 12–13.
41. Chen, X.; Wang, Y.; Sun, J.; Wang, J.; Xun, H.; Tang, F. Cloning, Expression and Functional Characterization of Two Sesquiterpene Synthase Genes from Moso Bamboo (*Phyllostachys edulis*). *Protein Expr. Purif.* **2016**, *120*, 1–6. [[CrossRef](#)] [[PubMed](#)]
42. Hsieh, H.L.; Ma, L.T.; Wang, S.Y.; Chu, F.H. Cloning and Expression of a Sesquiterpene Synthase Gene from *Taiwania cryptomerioides*. *Holzforchung* **2015**, *69*, 1041–1048. [[CrossRef](#)]
43. Nagegowda, D.A.; Gutensohn, M.; Wilkerson, C.G.; Dudareva, N. Two Nearly Identical Terpene Synthases Catalyze the Formation of Nerolidol and Linalool in Snapdragon Flowers. *Plant J.* **2008**, *55*, 224–239. [[CrossRef](#)] [[PubMed](#)]
44. Zhang, Y.; Li, Z.; Yu, X.; Fan, J.; Pickett, J.A.; Jones, H.D.; Zhou, J.; Michael, A.; Caulfield, J.; Napier, J.A.; et al. Molecular Characterization of Two Isoforms of a Farnesyl Pyrophosphate Synthase Gene in Wheat and Their Roles in Sesquiterpene Synthesis and Inducible Defence against Aphid Infestation. *New Phytol.* **2015**, *206*, 1101–1115. [[CrossRef](#)] [[PubMed](#)]
45. Yahyaa, M.; Matsuba, Y.; Brandt, W.; Doron-Faigenboim, A.; Bar, E.; McClain, A.; Davidovich-Rikanati, R.; Lewinsohn, E.; Pichersky, E.; Ibdah, M. Identification, Functional Characterization, and Evolution of Terpene Synthases from a Basal Dicot. *Plant Physiol.* **2015**, *169*, 1683–1697. [[CrossRef](#)] [[PubMed](#)]
46. Alquézar, B.; Rodríguez, A.; de la Peña, M.; Peña, L. Genomic Analysis of Terpene Synthase Family and Functional Characterization of Seven Sesquiterpene Synthases from *Citrus sinensis*. *Front. Plant Sci.* **2017**, *8*. [[CrossRef](#)] [[PubMed](#)]
47. Mahmood, T.; Yang, P.C. Western Blot: Technique, Theory, and Trouble Shooting. *N. Am. J. Med. Sci.* **2012**, *4*.
48. Jin, J.; Kim, M.J.M.J.; Dhandapani, S.; Tjhang, J.G.; Yin, J.-L.J.-L.; Wong, L.; Sarojam, R.; Chua, N.-H.N.-H.; Jang, I.-C. The Floral Transcriptome of Ylang Ylang (*Cananga odorata* Var. *fruticosa*) Uncovers Biosynthetic Pathways for Volatile Organic Compounds and a Multifunctional and Novel Sesquiterpene Synthase. *J. Exp. Bot.* **2015**, *66*, 3959–3975. [[CrossRef](#)] [[PubMed](#)]
49. Su, P.; Hu, T.; Liu, Y.; Tong, Y.; Guan, H.; Zhang, Y.; Zhou, J.; Huang, L.; Gao, W. Functional Characterization of NES and GES Responsible for the Biosynthesis of (E)-Nerolidol and (E,E)-Geranylinalool in *Tripterygium wilfordii*. *Sci. Rep.* **2017**, *7*, 40851. [[CrossRef](#)] [[PubMed](#)]
50. Keeling, C.I.; Bohlmann, J. Genes, Enzymes and Chemicals of Terpenoid Diversity in the Constitutive and Induced Defence of Conifers against Insects and Pathogens. *New Phytol.* **2006**, *170*, 657–675. [[CrossRef](#)] [[PubMed](#)]
51. Trapp, S.C.; Croteau, R. Genomic Organization of Plant Terpene Synthases and Molecular Evolutionary Implications. *Genetics* **2001**, *158*, 811–832. [[PubMed](#)]
52. Mercke, P.; Crock, J.; Croteau, R.; Brodelius, P.E. Cloning, Expression, and Characterization of Epi-Cedrol Synthase, a Sesquiterpene Cyclase from *Artemisia annua* L. *Arch. Biochem. Biophys.* **1999**, *369*, 213–222. [[CrossRef](#)] [[PubMed](#)]
53. Irmisch, S.; Krause, S.T.; Kunert, G.; Gershenzon, J.; Degenhardt, J.; Köllner, T.G. The Organ-Specific Expression of Terpene Synthase Genes Contributes to the Terpene Hydrocarbon Composition of Chamomile Essential Oils. *BMC Plant Biol.* **2012**, *12*, 84. [[CrossRef](#)] [[PubMed](#)]
54. Hsu, L.J.; Chu, F.H. Plasticity Residues Involved in Secondary Cyclization of Terpene Synthesis in *Taiwania cryptomerioides*. *Tree Genet. Genomes* **2015**, *11*, 796. [[CrossRef](#)]
55. Picaud, S.; Brodelius, M.; Brodelius, P.E. Expression, Purification and Characterization of Recombinant (E)- $\beta$ -Farnesene Synthase from *Artemisia annua*. *Phytochemistry* **2005**, *66*, 961–967. [[CrossRef](#)] [[PubMed](#)]
56. Xie, X.; Kirby, J.; Keasling, J.D. Functional Characterization of Four Sesquiterpene Synthases from *Ricinus communis* (Castor Bean). *Phytochemistry* **2012**, *78*, 20–28. [[CrossRef](#)] [[PubMed](#)]
57. Green, S.A.; Chen, X.; Nieuwenhuizen, N.J.; Matich, A.J.; Wang, M.Y.; Bunn, B.J.; Yauk, Y.K.; Atkinson, R.G. Identification, Functional Characterization, and Regulation of the Enzyme Responsible for Floral (E)-Nerolidol Biosynthesis in Kiwifruit (*Actinidia chinensis*). *J. Exp. Bot.* **2012**, *63*, 1951–1967. [[CrossRef](#)] [[PubMed](#)]
58. Rosano, G.L.; Ceccarelli, E.A. Recombinant Protein Expression in *Escherichia Coli*: Advances and Challenges. *Front. Microbiol.* **2014**, *5*, 172. [[CrossRef](#)] [[PubMed](#)]

59. Albertsen, L.; Chen, Y.; Bach, L.S.; Rattleff, S.; Maury, J.; Brix, S.; Nielsen, J.; Mortensen, U.H. Diversion of Flux toward Sesquiterpene Production in *Saccharomyces cerevisiae* by Fusion of Host and Heterologous Enzymes. *Appl. Environ. Microbiol.* **2011**, *77*, 1033–1040. [[CrossRef](#)] [[PubMed](#)]
60. Kitaoka, N.; Lu, X.; Yang, B.; Peters, R.J. The Application of Synthetic Biology to Elucidation of Plant Mono-, Sesqui-, and Diterpenoid Metabolism. *Mol. Plant* **2015**, *8*, 6–16. [[CrossRef](#)] [[PubMed](#)]
61. Tippmann, S.; Scalcinati, G.; Siewers, V.; Nielsen, J. Production of Farnesene and Santalene by *Saccharomyces cerevisiae* Using Fed-Batch Cultivations with R-Controlled Feed. *Biotechnol. Bioeng.* **2016**, *113*, 72–81. [[CrossRef](#)] [[PubMed](#)]
62. Šobotník, J.; Hanus, R.; Kalinová, B.; Piskorski, R.; Cvačka, J.; Bourguignon, T.; Roisin, Y. (E,E)- $\alpha$ -Farnesene, an Alarm Pheromone of the Termite *Prorhinotermes canalifrons*. *J. Chem. Ecol.* **2008**, *34*, 478–486. [[CrossRef](#)] [[PubMed](#)]
63. Ahmad-Sohdi, N.A.S.; Seman-Kamarulzaman, A.F.; Mohamed-Hussein, Z.A.; Hassan, M. Purification and Characterization of a Novel NAD (P)<sup>+</sup>-Farnesol Dehydrogenase from *Polygonum minus* Leaves. *PLoS ONE* **2015**, *10*, e0143310. [[CrossRef](#)] [[PubMed](#)]
64. Lapczynski, A.; Letizia, C.S.; Api, A.M. Fragrance Material Review on Cis-Nerolidol. *Food Chem. Toxicol.* **2008**, *46*, 245–246. [[CrossRef](#)] [[PubMed](#)]
65. Vinholes, J.; Gonçalves, P.; Martel, F.; Coimbra, M.A.; Rocha, S.M. Assessment of the Antioxidant and Antiproliferative Effects of Sesquiterpenic Compounds in in Vitro Caco-2 Cell Models. *Food Chem.* **2014**, *156*, 204–211. [[CrossRef](#)] [[PubMed](#)]
66. AbouLaila, M.; Sivakumar, T.; Yokoyama, N.; Igarashi, I. Inhibitory Effect of Terpene Nerolidol on the Growth of Babesia Parasites. *Parasitol. Int.* **2010**, *59*, 278–282. [[CrossRef](#)] [[PubMed](#)]
67. Chan, W.K.; Tan, L.T.H.; Chan, K.G.; Lee, L.H.; Goh, B.H. Nerolidol: A Sesquiterpene Alcohol with Multi-Faceted Pharmacological and Biological Activities. *Molecules* **2016**, *529*. [[CrossRef](#)] [[PubMed](#)]
68. Silva, M.P.N.; Oliveira, G.L.S.; De Carvalho, R.B.F.; De Sousa, D.P.; Freitas, R.M.; Pinto, P.L.S.; De Moraes, J. Antischistosomal Activity of the Terpene Nerolidol. *Molecules* **2014**, *19*, 3793–3803. [[CrossRef](#)] [[PubMed](#)]
69. Bunawan, H.; Choong, C.Y.; Md-Zain, B.M.; Baharum, S.N.; Noor, N.M. Molecular Systematics of *Polygonum minus* Huds. Based on ITS Sequences. *Int. J. Mol. Sci.* **2011**, *12*, 7626–7634. [[CrossRef](#)] [[PubMed](#)]
70. Abdul-Rahman, A.; Suleman, N.I.; Zakaria, W.A.; Goh, H.H.; Noor, N.M.; Aizat, W.M. RNA Extractions of Mangosteen (*Garcinia mangostana* L.) Pericarps for Sequencing. *Sains Malaysiana* **2017**, *46*, 1231–1240. [[CrossRef](#)]
71. Kong, S.L.; Abdullah, S.N.A.; Ho, C.L.; Amiruddin, M.D. Molecular Cloning, Gene Expression Profiling and in Silico Sequence Analysis of Vitamin E Biosynthetic Genes from the Oil Palm. *Plant Gene* **2016**, *5*, 100–108. [[CrossRef](#)]
72. Maille, P.E.O.; Chappell, J.; Noel, J.P. A Single-Vial Analytical and Quantitative Gas Chromatography—Mass Spectrometry Assay for Terpene Synthases. *Anal. Biochem.* **2004**, *335*, 210–217.

**Sample Availability:** Samples of the compounds  $\beta$ -farnesene and nerolidol are not available from the authors.



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).