

Supporting Information For:
Cross-coupling Reaction of Allylic Ethers with Aryl Grignard Reagents
Catalysed by a Nickel Pincer Complex

Toru Hashimoto,* Kei Funatsu, Atsufumi Ohtani, Erika Asano, and Yoshitaka Yamaguchi*

Department of Advanced Materials Chemistry, Graduate School of Engineering, Yokohama

National University, 79-5 Tokiwadai, Hodogaya-ku, Yokohama 240-8501, Japan

Contents

¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4a	S-2
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4b	S-3
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4c	S-4
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4d	S-5
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4e	S-6
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4f	S-7
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4g	S-8
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4h	S-9
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4i	S-10
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4j	S-11
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling products 4k and 5k	S-12
¹ H NMR and ¹³ C{ ¹ H} NMR spectra of coupling product 4l	S-13

^1H NMR and ^{13}C NMR spectra of coupling products























