Supplementary materials for

Structural optimization and structure–activity relationship of 4-thiazolidinone derivatives as novel inhibitors of human dihydroorotate dehydrogenase

Fanxun Zeng ^{1,§}, Lina Quan ^{2,§}, Guantian Yang ¹, Tiantian Qi ², Letian Zhang ¹, Shiliang Li ², Honglin Li ², Lili Zhu ^{2,} *, Xiaoyong Xu ^{1,*}

- ¹ Shanghai Key Laboratory of Chemical Biology, School of Pharmacy, East China University of Science and Technology, 130 Meilong Road, Shanghai 200237, PR China; 429646704@qq.com (F.Z.); 853565296@qq.com (G.Y.); 125702429@qq.com (L.Z); xyxu@ecust.edu.cn (X.X.)
- ² Shanghai Key Laboratory of New Drug Design, State Key Laboratory of Bioreactor Engineering, School of Pharmacy, East China University of Science & Technology, Shanghai 200237, PR China; Inquan1991@163.com (L.Q.); 785823121@qq.com (T.Q.); Slli403@163.com (S.L.); hlli@ecust.edu.cn (H.L.); zhulfl@ecust.edu.cn (L.Z.)

1. Shanghai Key Laboratory of Chemical Biology, School of Pharmacy, East China University of Science and Technology, 130 Meilong Road, Shanghai 200237, PR China, <u>xyxu@ecust.edu.cn</u> (X.X.); Tel.: +86-21-64252945 (X.X);

2. Shanghai Key Laboratory of New Drug Design, State Key Laboratory of Bioreactor Engineering, School of Pharmacy, East China University of Science & Technology, Shanghai 200237, PR China <u>zhulfl@ecust.edu.cn (L.Z.)</u>; Tel.: +86-21-64253379 (L.Z.)

§ These authors contributed equally to this work.

^{*} Correspondence:

The copies of spectrum for target compounds 7-46.(¹H NMR, ¹⁹F NMR and ¹³C NMR spectra were recorded on Bruker AM-400 (¹H at 400 MHz, ¹³C at 100 MHz, ¹⁹F at 376 MHz) spectrometer with CDCl₃ or DMSO-*d*₆ as the solvent and TMS as the internal standard.)



FigureS2.¹³C-NMR of compound 7.



FigureS4.¹³C-NMR of compound 8.



FigureS6.¹³C-NMR of compound 9.



FigureS8.¹⁹F-NMR of compound 10.



FigureS10.¹H-NMR of compound 11.



FigureS12.¹H-NMR of compound 12.



FigureS14.¹³C-NMR of compound 12.



FigureS16.¹⁹F-NMR of compound 13.



FigureS18.¹H-NMR of compound 14.



FigureS20.¹³C-NMR of compound 14.



FigureS22.¹⁹F-NMR of compound 15.



FigureS24.¹H-NMR of compound 16.



FigureS26.¹H-NMR of compound 17.



FigureS28.¹³C-NMR of compound 17.







FigureS30.¹⁹F-NMR of compound 18.



FigureS32.¹³C-NMR of compound 19.



FigureS34.¹³C-NMR of compound 20.



FigureS36.¹⁹F-NMR of compound 21.



FigureS38.¹H-NMR of compound 22.



FigureS40.¹H-NMR of compound 23.



FigureS42.¹H-NMR of compound 24.



FigureS44.¹H-NMR of compound 25.



FigureS46.¹H-NMR of compound 26.



FigureS48.¹H-NMR of compound 27.



FigureS50.¹H-NMR of compound 28.



FigureS52.¹H-NMR of compound 29.



FigureS54.¹H-NMR of compound 30.



FigureS56.¹³C-NMR of compound 30.



FigureS58.¹⁹F-NMR of compound 31.







FigureS61.¹³C-NMR of compound 32.























210 200 190 180 170 160 150 140 130 120 110 100 90 80 70 60 50 40 30 20 10 0 -10 f1 (ppm)

FigureS68.¹³C-NMR of compound 35.



FigureS70.13C-NMR of compound 36



FigureS72.¹³C-NMR of compound 37.



FigureS74.¹³C-NMR of compound 38.



FigureS76.¹³C-NMR of compound 39.



FigureS78.¹³C-NMR of compound 40.



FigureS80.¹³C-NMR of compound 41.



FigureS82.¹³C-NMR of compound 42.



FigureS84.¹³C-NMR of compound 43.



FigureS86.¹³C-NMR of compound 44.



FigureS88.¹³C-NMR of compound 45.



FigureS90.¹³C-NMR of compound 46.