

Eruca sativa Meal against Diabetic Neuropathic Pain: An H₂S-Mediated Effect of Glucoerucin

Elena Lucarini ¹, Eleonora Pagnotta ², Laura Micheli ¹, Carmen Parisio ¹, Lara Testai ^{3,4,5}, Alma Martelli ^{3,4,5}, Vincenzo Calderone ^{3,4,5}, Roberto Matteo ², Luca Lazzeri ², Lorenzo Di Cesare Mannelli ^{1,*} and Carla Ghelardini ¹.

¹ Department of Neuroscience, Psychology, Drug Research and Child Health—NEUROFARBA—Pharmacology and Toxicology Section, University of Florence, Florence 50139, Italy

² CREA-Council for Agricultural Research and Economics, Research Centre for Cereal and Industrial Crops, Bologna 40128, Italy

³ Department of Pharmacy, University of Pisa, Pisa 56126, Italy

⁴ Interdepartmental Research Centre “Nutraceuticals and Food for Health (NUTRAFOOD)”, University of Pisa, Pisa 56126, Italy

⁵ Interdepartmental Research Centre of Ageing Biology and Pathology, University of Pisa, Pisa 56126, Italy.

* Correspondence: lorenzo.mannelli@unifi.it

Supplementary Materials




Figure S1. Effect of acute administration of GSSG on STZ-induced neuropathic pain. The response to a mechanical and stimulus was evaluated by measuring the latency to pain-related behavior (withdrawal of the paw). GSSG (20 mg kg⁻¹) was orally and subcutaneous administered in STZ-treated animals and the tests were performed 30, 60, 90, and 120 min after the injection. ^^ $p < 0.01$ versus vehicle + vehicle-treated mice.