

SUPPORTING INFORMATION

Two new flavonoids from the leaves of *Baccharis oblongifolia* (Ruiz & Pav.) Pers. (Asteraceae)

Paulo R. F. Zampieri ¹, Cinthia I. Tamayose ¹, Oriana A. Fávero ², Paulete Romoff ², and Marcelo J. P. Ferreira

^{1,*}

¹ Departamento de Botânica, Instituto de Biociências, Universidade de São Paulo, 05508-090, São Paulo, Brazil; paulorfzampieri@usp.br (P.R.F.Z.); cinthiatamay@gmail.com (C.I.T.); marcelopena@ib.usp.br (M.J.P.F.)

² Universidade Presbiteriana Mackenzie, CEP 01302-907, São Paulo, Brazil; oriana.favero@mackenzie.br (O.A.F.); pauleteromoff@yahoo.com.br (P.R.)

* Correspondence: marcelopena@ib.usp.br; Tel.: +55-11-3091-7546 (M.J.P.F.)

List of Supporting Information

Figures S1–5. Spectral data of compound 1: oblongifolioside A [quercetin-3-O- β -[2''-O-(*E*)-caffeoyl]-rutinoside

Figures S6–11. Spectral data of compound 2: oblongifolioside B [kaempferol-3-O- β -[2''-O-(*E*)-caffeoyl]-rutinoside




Figure S1. ¹H NMR spectrum of compound 1: oblongifolioside A [quercetin-3-O-β-[2''-O-(E)-cafeoil]-rutinoside]




Figure S2. Expansion of ^1H NMR of compound 1: oblongifolioside A [quercetin-3-O-β-[2''-O-(E)-cafeoyl]-rutinoside




Figure S3. ¹³C NMR spectrum of compound 1: oblongifolioside A [quercetin-3-O-β-[2''-O-(E)-cafeoyl]-rutinoside]

Quercetina-3-O-β-[2''-O-(E)-cafeoil]-rutinosídeo



Figure S4. HSQC spectrum of compound 1: oblongifolioside A [quercetin-3-O-β-[2''-O-(E)-cafeoyl]-rutinoside

Quercetina-3-O-β-[2''-O-(E)-cafeoil]-rutinosídeo



Figure S5. HMBC spectrum of compound 1: oblongifolioside A [quercetin-3-O-β-[2''-O-(E)-cafeoil]-rutinoside

Caempferol-3-O-β-[2''-O-(E)-cafeoil]-rutinosídeo




Figure S6. ¹H NMR spectrum of compound 2: oblongifolioside B [kaempferol-3-O-β-[2''-O-(E)-cafeoyl]-rutinoside]




Figure S7. Expansion of ^1H NMR of compound 2: oblongifolioside B [kaempferol-3-O-β-[2''-O-(E)-caffeoyl]-rutinoside

Caempferol-3-O-β-[2''-O-(E)-cafeoil]-rutinosídeo




Figure S8. ¹³C NMR spectrum of compound 2: oblongifolioside B [kaempferol-3-O-β-[2''-O-(E)-cafeoil]-rutinoside]

Caempferol-3-O-β-[2''-O-(E)-cafeoil]-rutinosídeo




Figure S9. HSQC spectrum of compound 2: oblongifolioside B [kaempferol-3-O-β-[2''-O-(*E*)-cafeoyl]-rutinoside

Caempferol-3-O-β-[2''-O-(E)-cafeoil]-rutinosídeo




Figure S10. HMBC spectrum of compound 2: oblongifolioside B [kaempferol-3-O-β-[2''-O-(E)-cafeoil]-rutinoside




Figure S11. HR-ESI-MS spectrum of compound 2: oblongifolioside B [kaempferol-3-O- β -[2''-O-(*E*)-caffeoyl]-rutinoside