

Article



A three-step, gram-scale synthesis of Hydroxytyrosol, Hydroxytyrosol Acetate, and 3,4dihydroxyphenylglycol.

Amalia D. Kalampaliki¹, Vassiliki Giannouli¹, Alexios-Leandros Skaltsounis², Ioannis K. Kostakis^{1*}.

- ¹ Division of Pharmaceutical Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis-Zografou, Athens 15771, Greece
- ² Division of Pharmacognosy & Natural Product Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis-Zografou, Athens 15771, Greece
- * Correspondence: <u>ikkostakis@pharm.uoa.gr</u>

Contents

Figure S1: 1H-NMR and 13C NMR spectrum of compound 2.

Figure S2: ¹H-NMR and ¹³C NMR spectrum of compound 3.

Figure S3: ¹H-NMR and ¹³C NMR spectrum of compound 4.

Figure S4: ¹H-NMR and ¹³C NMR spectrum of compound 5.

Figure S5: ¹H-NMR and ¹³C NMR spectrum of compound 6.

Figure S6: ¹H-NMR and ¹³C NMR spectrum of compound 7.

Figure S7: ¹H-NMR and ¹³C NMR spectrum of compound 8.




Figure S1: ¹H-NMR and ¹³C NMR spectrum of compound 2.




Figure S2: ¹H-NMR and ¹³C NMR spectrum of compound 3.




Figure S3: ¹H-NMR and ¹³C NMR spectrum of compound 4.




Figure S4: ¹H-NMR and ¹³C NMR spectrum of compound 5.




Figure S5: ¹H-NMR and ¹³C NMR spectrum of compound 6.




Figure S6: ¹H-NMR and ¹³C NMR spectrum of compound 7.

7 of 8




Figure S7: ¹H-NMR and ¹³C NMR spectrum of compound 8.