

Supplementary information

Multi-steps fragmentation-ion trap mass spectrometry coupled to liquid chromatography diode array system for investigation of olaparib related substances

Alaa Khedr¹. Soad S. Abd El-Hay². Ahmed K. Kammouna¹

¹ Department of Pharmaceutical Chemistry, Faculty of Pharmacy, King Abdulaziz University,
Jeddah 21589, P.O. Box 80260, Saudi Arabia

² Department of Analytical Chemistry, Faculty of Pharmacy, Zagazig University, Egypt

IT-MS spectra of OLA and OLA-related substances, +MS^{2,3} and -MS^{2,3}




Figure S1: Average negative and positive IT-MS^{2,3} spectra of m/z +435, +435→+367 and -433, -433→253




Figure S2: Average negative and positive IT-MS^{2,3} spectra of m/z +299, +299→+281 and -297, -297→253




Figure S3: Average negative and positive IT-MS^{2,3} spectra of m/z +367, +367→+281 and -365, -365→322.




Figure S4: Average negative and positive IT-MS^{2,3} spectra of m/z +409, +409→+367 and -407, -407→253.




Figure S5: Average positive IT-MS^{2,3} spectra of m/z +417, +417 → +263.




Figure S6: Average positive IT-MS^{2,3} spectra of m/z +326, +326 → +281.