

Article

Alkaline extraction, structural characterization, and bioactivities of novel (1→6)-β-D-glucan from *Lentinus edodes*

Jia Li ¹, Chao Cai ^{1,2,*}, Mengmeng Zheng ³, Jiejie Hao ^{1,2}, Ya Wang ¹, Minghua Hu ⁴, Luodi Fan ⁴, Guangli Yu ^{1,2,*}

¹ Key Laboratory of Marine Drugs of Ministry of Education & Shandong Provincial Key Laboratory of Glycoscience and Glycotechnology, School of Medicine and Pharmacy, Ocean University of China, Qingdao 266003, China.; lijia199208@163.com (J. Li); caic@ouc.edu.cn (C. Cai); 18364166659@163.com (M. Zheng); 2009haojie@ouc.edu.cn (J. Hao); yawangouc@163.com (Y. Wang); Mandy.Hu@infinitus-int.com (M. Hu); Franz.Fan@infinitus-int.com (L. Fan); glyu@ouc.edu.cn (G. Yu)

² Laboratory for Marine Drugs and Bioproducts, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, China., School of Medicine and Pharmacy, Ocean University of China, Qingdao 266003, China.

³ Laboratory of Chinese medicine pharmacy, School of Pharmacy, Shandong University of Traditional Chinese Medicine, Jinan, 250355, China.

⁴ Infinite Pole (China) Co., LTD., Guangdong, Guangzhou 510600, China

* Correspondence: caic@ouc.edu.cn (C. Cai); glyu@ouc.edu.cn (G. Yu) Tel.: +86-532-8203-1609

Contents

Table S1. The results of L₉ (3³) orthogonal test for *lentinus edodes* polysaccharide

Table S2. Orthogonal extraction design of *lentinus edodes* polysaccharides

Figure S1. The ¹H NMR spectrum of OEs in DMSO/D₂O (6:1 v/v) at 60 °C.

Figure S2. FT-IR spectrum of LeP-N2 after full methylation.

Figure S3. The ¹H-¹H COSY spectrum of LeP-N2

Table S1. The results of L₉ (3³) orthogonal test

Orthogonal Experiment number	Factor			Extraction yield(%)
	Temperature A (°C)	Time B (h)	Concentration C (mol/L)	
1	0	0.5	0.1	3.59
2	0	1	0.25	6.11
3	0	2	0.5	9.84
4	20	0.5	0.25	5.98
5	20	1	0.5	9.50
6	20	2	0.1	5.28
7	60	0.5	0.5	8.95
8	60	1	0.1	7.11
9	60	2	0.25	7.73
K ₁	6.513	6.173	5.327	6.940
K ₂	6.920	7.573	6.607	6.780
K ₃	7.930	7.617	9.430	7.643
R	1.417	1.444	4.103	

Table S2. Orthogonal extraction design of *lentinus edodes* polysaccharides

Level	Factor		
	Temperature A (°C)	Time B (h)	Concentration C (mol/L)
1	0	0.5	0.1
2	20	1	0.25
3	60	2	0.5




Figure S1. The ^1H -NMR spectrum of OEs in DMSO/D $_2$ O (6:1 v/v) at 60°C.




Figure S2. FT-IR spectrum of LeP-N2 after full methylation.




Figure S3. The ^1H - ^1H COSY spectrum of LeP-N2