One-Pot Multicomponent Synthesis and Bioevaluation of Tetrahydroquinoline Derivatives as Potential Antioxidants, α -Amylase Enzyme Inhibitors, Anti-Cancerous and Anti-Inflammatory Agents.

Samra Farooq^{1,}, Aqsa Mazhar¹, Areej Ghouri¹, Ihsan-Ul-Haq¹ and Naseem Ullah^{1,*}

¹Department of Pharmacy, Faculty of Biological Science, Quaid-I-Azam University Islamabad, 45320, Pakistan

samrafarooq@bs.qau.edu.pk (S.F.,); aqsamazhar1947@gmail.com (A.M.,); areejkhan27@gmail.com (A.G.,); ihaq@qau.edu.pk (I.H.,)

* Correspondence: Naseem Ullah (N.U.,) Assistant Professor <u>nullah@qau.edu.pk</u>; Tel: +92-334-5023566



Figure S1: Proton NMR of compound SF1









Figure S4: Carbon NMR of compound SF2



Figure S5: Proton NMR of compound SF3



Figure S6: Carbon NMR of compound SF3







Figure S9: Proton NMR of compound SF5







Figure S12: Carbon NMR of compound SF6



Figure S13: Proton NMR of compound SF7







Figure S15: Proton NMR of compound SF8



Figure S16: Carbon NMR of compound SF8



Figure S17: Proton NMR of compound SF9



Figure S18: Carbon NMR of compound SF9



Figure S19: Proton NMR of compound SF10







Figure S21: Proton NMR of compound SF11



Figure S22: Carbon NMR of compound SF11



Figure S23: Proton NMR of compound SF12



Figure S24: Carbon NMR of compound SF12



Figure S25: Proton NMR of compound SF13



