

Supplemental Information

Preparation of Chitosan Molecularly Imprinted Polymers and the Recognition Mechanism for Adsorption of Alpha-Lipoic Acid

Long Xu ^{1,2}, Ze-Xin Zhao ³, Yun-An Huang ⁴, Qiu-Jin Zhu ^{1,5*}

¹ School of Liquor and Food Engineering, Guizhou University, Guiyang 550025, China; xulong19891026@163.com

² School of Food Science and Engineering, South China University of Technology, Guangzhou 510640, China;

³ School of Bioscience and Bioengineering, South China University of Technology, Guangzhou 510006, China; bizexin-zhao@mail.scut.edu.cn

⁴ Department of Food Engineering, Guizhou Vocational College of Foodstuff Engineering, Guiyang 551400, China; hya127@163.com

⁵ Key Laboratory of Agricultural and Animal Products Store and Processing of Guizhou Province, Guiyang 550025, China.

*Correspondence: ls.qjzhu@gzu.edu.cn; Tel.: +86-851-8385-1195




Figure S1. The initial model of ALAs (ball-and-stick model) and chitosan hexamers (vdW surface model) in water box.




Figure S2. RMSD analysis. (A) The chitosan micelle model formed after 16 ns simulation time, (B) The RMSD curve of chitosan.