

Supplementary data

Bersavine: a novel bisbenzylisoquinoline alkaloid with cytotoxic, antiproliferative and apoptosis-inducing effect on human leukemic cells

Darja Koutova^a, Monika Kulhava^a, Radim Havelek^{a*}, Martina Majorosova^a, Karel Kralovec^b, Klara Habartova^a, Anna Hostalkova^c, Lubomir Opletal^c, Lucie Cahlikova^c, Martina Rezacova^a

^aDepartment of Medical Biochemistry, Faculty of Medicine in Hradec Kralove, Charles University, Simkova 870, Hradec Kralove 500 03, Czech Republic

^bDepartment of Biological and Biochemical Sciences, Faculty of Chemical Technology, University of Pardubice, Studentska 573, Pardubice 532 10, Czech Republic

^cADINACO Research Group, Department of Pharmaceutical Botany, Faculty of Pharmacy, Charles University, Heyrovskeho 1203, Hradec Kralove 500 05, Czech Republic

**Corresponding author*

Radim Havelek, Ph.D.
Department of Medical Biochemistry,
Faculty of Medicine in Hradec Králové,
Charles University,
Šimkova 870,
500 03 Hradec Králové,
Czech Republic,
Tel.: +420495816293,
E-mail address: havelekr@lfhk.cuni.cz

IC₅₀ curves of bersavine for the most sensitive cancer cell lines (cell lines demonstrating at least 50% inhibition of proliferation at 10 μ M single-dose treatment)




Fig. S1. IC₅₀ curves of bersavine for the most sensitive cancer cell lines HeLa, MCF-7, HT-29, Jurkat and MOLT-4. IC₅₀ values are given in the table 3. The results are expressed as mean values of three independent experiments $\pm$ SD.