

Review

Recent Advances in the Chemical Synthesis and Evaluation of Anticancer Nucleoside Analogues

Mieke Guinan ¹, Caecilie Benckendorff ¹, Mark Smith ² and Gavin J. Miller ^{1,*}

¹ Lennard-Jones Laboratory, School of Chemical and Physical Sciences, Keele University, Keele, Staffordshire, ST5 5BG, UK; m.guinan@keele.ac.uk (M.G.); c.m.m.benckendorff@keele.ac.uk (C.B.)

² Medicinal Chemistry Knowledge Center, Stanford ChEM-H, 290 Jane Stanford Way, Stanford, CA 94305, USA; mxsmith@stanford.edu

* Correspondence: g.j.miller@keele.ac.uk

Academic Editor: Derek J. McPhee

Received: 2 April 2020; Accepted: 25 April 2020; Published: 28 April 2020

Abstract: Nucleoside analogues have proven to be highly successful chemotherapeutic agents in the treatment of a wide variety of cancers. Several such compounds, including gemcitabine and cytarabine, are the go-to option in first-line treatments. However, these materials do have limitations and the development of next generation compounds remains a topic of significant interest and necessity. Herein, we discuss recent advances in the chemical synthesis and biological evaluation of nucleoside analogues as potential anticancer agents. Focus is paid to 4'-heteroatom substitution of the furanose oxygen, 2'-, 3'-, 4'- and 5'-position ring modifications and the development of new prodrug strategies for these materials.

Keywords: nucleoside analogue; anti-cancer; chemical synthesis; heteroatom replacement; chemotherapeutic; prodrug

1. Introduction

A significant proportion of current chemotherapeutic treatments for cancer involve the use of anti-metabolites, particularly modified nucleoside analogues that possess a capability to mimic native purine or pyrimidine nucleosides which can disrupt metabolic and regulatory pathways [1]. These molecules can be taken up by nucleoside transporters and then phosphorylated to their mono-, di- and triphosphate forms where they are able to interfere with DNA/RNA synthesis and repair; for example, by acting as chain terminators [2] or ribonucleotide reductase inhibitors [3]. Other notable modes of action include epigenetic regulation, through inhibition of DNA regulatory proteins, such as DNA methyltransferase [4]. Selected current examples of anticancer nucleoside analogues approved for chemotherapeutic treatment regimens include capecitabine, gemcitabine **1**, clofarabine **2** and cytarabine (Ara-C) **3** (Figure 1).




Figure 1. Gemcitabine **1**, clofarabine **2** and Ara-C **3**. Modifications compared to native *D*-ribo-configured purine or pyrimidine nucleosides are shown in blue.

Therapeutic intervention using nucleoside analogues is not without its problems and their use is often limited by poor cellular uptake, low conversion to the active triphosphate metabolite, rapid degradation or clearance and development of resistance profiles in certain cell types [5,6]. Consequently, research continues to develop next generations of nucleoside analogues that overcome some of these limitations and provide new therapeutic options.

This class of antimetabolite also possess a proud history, and current frontline treatment, as antiviral [7–12] and, more recently, antibacterial agents [13]. Indeed, the development of nucleoside analogues has a symbiotic relationship between compound class and final therapeutic treatment. For example, gemcitabine **1** was developed as an antiviral, but was subsequently shown to be very toxic to leukaemia cells.

In this review, we survey developments from 2010 onwards for the chemical synthesis and evaluation of modified nucleoside analogues for anticancer research. Specifically, focusing on alterations to the native furanose ring (Figure 2) and generally retaining native purine or pyrimidine nucleobases. Comprehensive reviews concerning hetero-base modifications and general trends in nucleotide synthesis have been covered elsewhere [14,15]. The review is divided into sections that systematically consider: i) furanose 4'-oxygen atom replacements with N, S, Se and C ii) 2', 3', 4'- or 5'-position furanose ring modifications and iii) new prodrug approaches to deliver nucleoside analogues.




Figure 2. General scope for nucleoside analogues covered in this review. Base = purine or pyrimidine (i.e., C, U, T, A, G or close derivative thereof). X = heteroatom or carbon and Y and Z = ring functional group or modification of native *D-ribo* stereochemistry.

2. Furanose Oxygen Atom Replacements

2.1. Azanucleosides

Azanucleosides were originally defined as nucleoside analogues where the furanosyl oxygen is replaced by nitrogen, however this group of analogues has been extended to include nucleosides where the resultant pyrrolidine core has been replaced by other nitrogen containing rings, including heterocycles, heterotricycles and acyclic nitrogen-containing nucleosides [16,17]. This class of compound have proven successful in the treatment of cancer [18] and have also been established as having antiviral and antibacterial properties [19].

2.1.1. Development of Forodesine

Purine nucleoside phosphorylases (PNPs) are responsible for the phosphorolytic metabolism of purine nucleosides to ribose/deoxyribose phosphate and the corresponding nucleobase. Patients with abnormally low levels of PNP possess little T-cell immunity due to a severely reduced degradation of deoxyguanosine, which results in the accumulation of the corresponding triphosphate (dGTP). This then reduces the activity of ribonucleotide reductase and induces apoptosis. As such, human PNP inhibitors are potential treatments for T-cell lymphomas [17]. Immucillin H (forodesine) **4** (Figure 3) is a highly potent PNP inhibitor ($IC_{50} = 0.48\text{--}1.57\text{ nM}$) which is effective against T-cell malignancies and was found to have excellent oral bioavailability in mice (63%) [20].

Forodesine is a gradual onset drug which binds tightly to PNP with a high affinity ($K_i = 0.023\text{ nM}$) [21]. Whilst clinical development of **4** was discontinued in the US and Europe, it was recently approved for use in the treatment of relapsed/refractory peripheral T-cell lymphoma (PCTL) in Japan (April 2017) [22].





Figure 3. Structure of forodesine 4.

Forodesine is a guanosine analogue and a transition state inhibitor of PNP with 100–1000-fold higher potency than previously identified inhibitors [23]. Due to the combined substitution of the furanosyl oxygen with nitrogen and the C-glycosidic bond, 4 is not incorporated into DNA, acting only as a highly selective PNP inhibitor [22]. Also noteworthy is an adenosine mimetic of 4 which is currently under development as a broad-spectrum antiviral [24].


In 2000, Tyler et al. described a linear synthesis of 4 in a satisfactory 39% yield over 10 steps (Scheme 1) [25]. Starting from 5, synthesised in nine steps by known methods from D-gulonolactone [26], the pyrrolidine was treated with *N*-chlorosuccinimide (NCS), obtaining the 1-chloro anomeric glycoside which subsequently underwent elimination using lithium tetramethylpiperidine (LiTMP) to afford imine 6. The nucleobase was next assembled via addition of lithiated acetonitrile to afford 7, followed by protection of the furanosyl nitrogen giving 8 and treatment with Bredereck's reagent to afford enamine 9. Acid-catalysed hydrolysis of 9 delivered enol 10 which was reacted with ethyl glycinate to obtain enamine 11. Treatment of 11 with excess benzyl chloroformate and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) revealed 12 and subsequent hydrogenolysis of the *N*-Cbz group provided pyrrole 13. Completion of the carbocyclic nucleobase was achieved via treatment of 13 with formamidine acetate and acidic removal of the silicon, nitrogen and isopropylidene protecting groups, to afford 4.



Scheme 1. Reagents and conditions: (i) NCS, pentane; (ii) LiTMP, $-78\text{ }^{\circ}\text{C}$, 36% over two steps; (iii) $^n\text{BuLi}$, MeCN, THF, $-78\text{ }^{\circ}\text{C}$ then tetramethylpiperidine $-78\text{ }^{\circ}\text{C}$, 100%; (iv) $(\text{Boc})_2\text{O}$, CH_2Cl_2 ; (v) $^t\text{BuOCH}(\text{NMe}_2)_2$, DMF, $70\text{ }^{\circ}\text{C}$; (vi) THF, AcOH, H_2O , 72% from 7; (vii) $\text{H}_2\text{NCH}_2\text{CO}_2\text{Et}:\text{HCl}$, NaOAc, MeOH (viii) ClCO_2Bn , DBU, CH_2Cl_2 , reflux, 67% from 10; (ix) H_2 , Pd/C, EtOH; (x) $\text{H}_2\text{NCH}=\text{NH}:\text{AcOH}$, EtOH, reflux, 91% from 12 and (xi) TFA, 81%.

Forodesine was found to have low oral bioavailability (<11%) in primates, contrary to the case in mice (63% [20]) and was thus originally developed as an intravenous formulation [21]. In 2005, Morris, Jr et al. synthesised BCX-3040, the 2'-deoxy analogue of 4, and comparatively evaluated its

oral pharmacokinetic and pharmacodynamic properties in an effort to maintain potency and deliver oral bioavailability [27]. This was hypothesised from 2'-deoxyguanosine exhibiting tight binding to PNP [28] and therefore a possible redundancy for the 2'-OH. Starting from vinyl bromide **14** (Scheme 2), the 9-deazapurine underwent bromine-lithium exchange and addition to imine **6**, obtaining nucleoside **15** in 85% yield. Subsequent removal of 5'-O-TBS and 2',3'-O-isopropylidene groups gave **16**, followed by 3',5'-OTIPDS protection to afford **17**. The free 2'-hydroxyl group was then converted to thiocarbonate **18** and deoxygenated via treatment with 1,1'-azobis(cyclohexane-1-carbonitrile) in excellent yield (91%) to give **19**. Deprotection was completed in two steps, first cleaving the CH₂OCH₂Ph (BOM) group, followed by acidic hydrolysis and hydrochloride salt formation to obtain **BCX-3040** in 83% yield (from **19**)



Scheme 2. Reagents and conditions: (i) **6**, ⁿBuLi, anisole, ether, -70 °C; (ii) (Boc)₂O, CH₂Cl₂, 85% over two steps; (iii) 1M HCl, MeOH, 30 °C, 96%; (iv) 1,3-Dichloro-1,1,3,3-tetraisopropylidisiloxane, pyridine, 81%; (v) O-Phenyl chlorothionoformate, MeCN, 90%; (vi) 1,1'-Azobis(cyclohexane-1-carbonitrile), toluene reflux, 91%; (vii) Pd(OH)₂, H₂, EtOH, conc. NH₄OH, 90% and (viii) conc. HCl, MeOH, reflux, 83%.

In vitro biological evaluation of **BCX-3040** confirmed it to be a potent PNP inhibitor, with near identical IC₅₀ values to **4** (**BCX-3040** IC₅₀ = 3.1 ± 0.50 nM and **4** IC₅₀ = 1.2 ± 0.21 nM). Administration of 5.0 mg of **4** had a 12.6-fold greater 2'-deoxyguanosine response than administration of 10.0 mg of **BCX-3040**, indicating a reduced bioavailability for **BCX-3040**. Furthermore, following IV administration of 5 mg/kg of **BCX-3040**, the plasma concentration of **BCX-3040** dropped rapidly to 3.0 ± 0.31 µg/mL. Overall, due to the observed rapid clearance and reduced bioavailability of **BCX-3040**, it was concluded to be a poorer PNP inhibitor candidate in comparison to **4**.


2.2. Thionucleosides

4'-Thiofuranoses are of known importance in biological systems, for example as chemical biology or biomedical tools [29], and possess chemotherapeutic activity [30,31]. Furthermore, the thioaminal moiety within 4'-thiofuranosyl nucleosides has been proven to be resistant to metabolic hydrolysis in comparison to native 4'-oxa analogues [32]. In the early 1990s, Secrist, Montgomery and co-workers stimulated interest in this class of molecule with their synthesis and biological evaluation

of 2'-deoxy-4'-thiopyrimidine nucleosides [30]. Since then, there has been a resurgence of interest in the synthesis and evaluation of these compounds as potential antiviral and chemotherapeutic agents.

2.2.1. 2'-Modified Thionucleosides


Yoshimura et al. reported the synthesis and biological evaluation of 4'-thia-1-(2-deoxy-2-C-methylene- β -D-erythro-pentofuranosyl)cytosine (4'-thio-DMDC) **28 β** and 2'-deoxy-2'-fluoro-*arabino*-4'-thiacytidine **33 β** as potential antitumour agents [33–35]. Their synthesis of **28** started from 1,2,5,6-diisopropylidene-D-glucose **20** and a series of protecting group manipulations delivered 3-O-benzyl D-xylose methyl glycoside **21** (Scheme 3). Mesylation of the 2- and 5-hydroxyl groups enabled reaction with sodium sulfide to afford 2,5-bicyclic intermediate **22** as a mixture of anomers. Following conversion of **22** to 4-thio-*arabinofuranose* **23**, the 2-position hydroxyl group was oxidised and the ketone homologated using a Wittig reagent to install the 2-methylene component, with this material then oxidised to sulfoxide **24**.



Scheme 3. Reagents and conditions: (i) BnBr, NaH, DMF, THF; (ii) 2M HCl, THF; (iii) NaIO₄, H₂O, MeOH; (iv) NaBH₄, MeOH, 84% from **20**; (v) 5% HCl/MeOH, 91%; (vi) MsCl, pyridine; (vii) Na₂S, DMF, α -anomer 78% from **21**, β -anomer 73% from **21**; (viii) 4M HCl, THF; (ix) NaBH₄, MeOH, 90% from **2**; (x) TBDPSCI, imidazole, DMF, 87%; (xi) Ac₂O, DMSO; (xii) Ph₃PCH₃Br, NaH, *t*-amyl alcohol, THF; (xiii) BCl₃, CH₂Cl₂, -78 °C then MeOH, pyridine, 92%; (xiv) *m*-CPBA, CH₂Cl₂, -78 °C, 74% from **23**; (xv) **25**, TMSOTf, ClCH₂CH₂Cl, 0 °C, 29%; (xvi) TBAF, THF and (xvii) aqueous NH₃, MeOH then HPLC separation.


The cytidine nucleobase was installed using a Pummerer-type thioglycosylation, via sulphenium ion **26**, which afforded **27** as a mixture of anomers. These were fully deprotected to afford **28 α** and **28 β** and the desired **28 β** isolated using HPLC separation.

Additionally, 2'-deoxy-2'-fluoro-*arabino*-4'-thiacytidine **33** was prepared from intermediate **23** (Scheme 4). Stereospecific DAST fluorination of **23** proceeded through epi-sulphonium intermediate **29** which delivered the 2-deoxy-2-fluoro-*arabino* intermediate **30** in 68% yield. *m*-CPBA oxidation to the sulfoxide **31** and subsequent Pummerer rearrangement formed anomeric acetate **32** in 77% yield from **30**. Finally, thioglycosylation was successfully employed to access the corresponding 4'-thionucleoside mixture **33** (54% yield). Comparatively, the group found that using **32** as a donor and employing silyl-Hilbert-Johnson glycosylation conditions yielded **33 α/β** in higher yield (93%), but still as a mixture of anomers (2.9:1 α/β).



Scheme 4. Reagents and conditions: (i) DAST, CH_2Cl_2 , $-78\text{ }^\circ\text{C}$, 77%; (ii) *m*-CPBA, CH_2Cl_2 , $-78\text{ }^\circ\text{C}$; (iii) Ac_2O , $100\text{ }^\circ\text{C}$, 77% from **23**; (iv) **25**, SnCl_4 , MeCN, 93%; (v) BBr_3 , MeOH; (vi) NH_4F , MeOH, $60\text{ }^\circ\text{C}$ and (vii) aqueous NH_3 , MeOH then HPLC separation, 43% (β -anomer) and 17% (α -anomer) from **32**.

Finally, the group synthesised 2'-deoxy-2'-difluoro-1'(4'-thia-D-ribofuranose)cytosine **37**, a thionucleoside analogue of the potent chemotherapeutic agent gemcitabine **1** (Scheme 5). Intermediate **23** was again utilised and oxidised using Albright-Goldman conditions to obtain ketone **34** which was difluorinated at C2 using DAST. The 3-O-benzyl group was next removed and replaced with benzoate to afford **35** in 79% overall yield from **34**. Oxidation to the sulfoxide afforded **36** which subsequently underwent Pummerer rearrangement-glycosylation, before the remaining protecting groups were removed to afford **37α/β** ($\alpha/\beta = 2.4/1$) in a moderate yield of 51% from **36**.



Scheme 5. Reagents and conditions: (i) Ac_2O , DMSO; (ii) DAST, benzene, $0\text{ }^\circ\text{C}$ -r.t., 48%; (iii) BCl_3 , CH_2Cl_2 , $-78\text{ }^\circ\text{C}$, then MeOH, pyridine; (iv) Bz_2O , Et₃N, DMAP, MeCN, 79% from **34**; (v) *m*-CPBA, CH_2Cl_2 , $-78\text{ }^\circ\text{C}$; (vi) **25**, TMSOTf, $\text{ClCH}_2\text{CH}_2\text{Cl}$, $0\text{ }^\circ\text{C}$, 57% from **35**; (vii) TBAF, THF and (viii) aqueous NH_3 , MeOH, then HPLC separation, 36% (α -anomer) and 15% (β -anomer) from **36**.

The antineoplastic activities of **28**, **33** and **37** were evaluated and compared to arabinocytidine **3** (Ara-C) and 1-(2-deoxy-2-C-methylene- β -D-erythro-pentofuranosyl)cytosine (DMDC) (Table 1). As expected, all the α -anomer forms were found to be inactive against T-cell leukemia CCRF-HSB-2 cells. However, the β -anomers showed considerable cytotoxic activity against the same cell line. Notably, **28β** and **33β** were highly potent against both T-cell leukemia CCRF-HSB-2 cells and human solid tumour KB cells, with IC_{50} values of $0.01\text{ }\mu\text{g/mL}$ (CCRF-HSB-2) and $0.05\text{ }\mu\text{g/mL}$ (CCRF-HSB-2) for **28β** and **33β**, respectively. Indeed, the activity of **28β** was significantly higher than that of its native counterpart, DMDC, against both cell lines, with an IC_{50} value 2.4 times lower in CCRF-HSB-2 cells and 3.7 times lower in KB cells. Interestingly, 4'-thiogemcitabine analogue **37β** had poorer antineoplastic activity compared to **1**, which the authors suggest may be due to a reduction in the phosphorylation efficacy of **37β** by deoxycytidine kinase, a key enzyme which converts 2'-deoxycytidine analogues to their corresponding monophosphates.


Table 1. Antineoplastic activities of 2'-modified-4'-thionucleosides.

Compound (Anomer)	2'-substituent	Antineoplastic Activities IC ₅₀ (µg/mL)	
		CCRF-HSB-2 ^a	KB cells ^b
28 α	=CH ₂	>10	ND ^c
28 β	=CH ₂	0.01	0.12
33 α	F (arabino)	>10	ND
33 β	F (arabino)	0.05	0.02
37 α	F ₂	>10	ND
37 β	F ₂	1.5	17
Ara-C 3		0.05	0.26
DMDC		0.02	0.44


^aMTT assay [36]; ^bdye uptake method [36]; ^cnot determined.

2.2.2. 4'-Modified-2'-deoxythionucleosides

Following earlier work by Parker and colleagues, who identified 4'-thia-2'-deoxycytidine (T-dCyd) as being able to inhibit tumour growth [37], Haraguchi and colleagues reported the synthesis and evaluation of a small library of 4'-position modified 4'-thia-2'-deoxycytidine nucleosides **38–41** for their antineoplastic and antiviral activity (Figure 4) [38].


**Figure 4.** Structures of 2'-deoxy-4'-thiacytidine nucleosides **38–41**.

Towards analogue **38** the group started from thioglycal **42** (Scheme 6), obtained using established procedures from 2-deoxy-D-ribose in 12 steps [39]. This material was transformed into a glycosyl donor through treatment with *N*-iodosuccinamide (NIS) and pivalic acid, giving one diastereoisomer of 2-iodo derivative **43**. Silylated uracil was then glycosylated with **43** using Vorbrüggen-type conditions to give β -4'-thiouridine **44** which was 2'-deoxygenated via a Bu₃SnH mediated radical reduction to give **45**. Subsequent TIPDS deprotection, followed by C3' and C5'-*O*-acetylation delivered **46** to enable a four-step procedure to deliver 4',5'-unsaturated-4'-thiouridine derivative **47**. *Exo*-thioglycal **47** was next converted to silyl-protected 4'-thionucleoside **48**, which when treated with Pb(OBz)₄ yielded dibenzoate **49**. S_N2 inversion back to the native 4'-D-ribo configuration and 4'-azido installation was achieved by reaction with MeSiN₃ in the presence of SnCl₄ and afforded the desired 4'- α -azido derivative **50** as the major product. Finally, the nucleoside was converted to the cytidine form via intermediate **51** to give **38** in 7% yield over 18 steps.



Scheme 6. Reagents and conditions: (i) NIS, pivalic acid, MeCN/CH₂Cl₂, 93%; (ii) Uracil, *N*,*O*-bis(trimethylsilyl)acetamide (BSA), TMSOTf, MeCN/CH₂Cl₂, 87%; (iii) Bu₃SnH, Et₃B, toluene, O₂, −60 °C, 98%; (iv) TBAF, Ac₂O, THF, 100%; (v) NH₃/MeOH; (vi) I₂, PPh₃, pyridine, dioxane; (vii) Ac₂O, DMAP, DIPEA, CH₂Cl₂; (viii) BDN, MeCN, 65% over four steps; (ix) NH₃/MeOH; (x) TBDMSCl, imidazole, DMF, 61% over two steps; (xi) Pb(OBz)₄, toluene, 63%; (xii) TMSN₃, SnCl₄, CH₂Cl₂, 61%; (xiii) NaOMe, MeOH; (xiv) Ac₂O, DMAP, DIPEA, MeCN, 83% over two steps; (xv) TPSCl, K₂O₄, MeCN, 60 °C; (xvi) NH₄OH, THF; (xvii) TBAF, Ac₂O, THF and (xviii) NaOMe, MeOH, 66% over four steps.


In order to access 2'-deoxy-4'-*C*-fluoromethyl-4'-thiacytidine **39**, alcohol **53** was prepared from known aldehyde **52** [40,41]. Treatment of **53** with DAST successfully installed the key 4- α -fluoromethyl group which was then elaborated to **39** (Scheme 7). 4- α -alkynyl and nitrile analogues **40** and **41** were synthesised from aldehyde **54** using a late-stage insertion of the key functional group at the 4'-position in the presence of the nucleobase. Of the four 2'-deoxy-4'-modified thionucleosides **38–41**, two analogues, **38** and **39**, showed moderate cytotoxicity against human B-cell leukaemia (CCRF-SB; IC₅₀ = 7.14 μ M and 3.19 μ M for **38** and **39** respectively) and human T-cell leukaemia cell lines (Molt-4; IC₅₀ = 2.72 μ M and 2.24 μ M for **38** and **39** respectively).



Scheme 7. Key intermediates to access 2'-deoxy-4'-thiacytidine nucleosides **39–41**.

2.3. Selenonucleosides

A first synthesis of pyrimidine 4'-selenonucleosides was reported by Jeong and colleagues in 2008 (Scheme 8), [42] starting from lyxose derivative **55**, synthesised from D-gulonic- γ -lactone in four steps using established procedures [43].



Scheme 8. Reagents and conditions: (i) TBDPSCl, Et₃N, DMAP, CH₂Cl₂, 92%; (ii) NaBH₄, MeOH, 98%; (iii) MsCl, Et₃N, DMAP, CH₂Cl₂, 97%; (iv) Se, NaBH₄, EtOH, THF, 60 °C, 96%; (v) *m*-CPBA, CH₂Cl₂, –78 °C, 85%; (vi) Uracil, Et₃N, TMSOTf, toluene, CH₂Cl₂, 53%; (vii) 50% aq. TFA, 81%; (viii) N³-benzoylcytosine, Et₃N, TMSOTf, toluene, CH₂Cl₂, 35%; (ix) 50% Aq. TFA and (x) NH₃, MeOH, 82% over two steps.

Selective protection of the primary alcohol in **55** was achieved by reaction with TBDPSCl, giving hemiacetal **56** which was reduced with NaBH₄, furnishing diol **57**. Following double mesylation of **57** to give **58**, cyclisation to give 4-selenosugar **59** was achieved by treatment with selenium in the presence of NaBH₄. Oxidation of **59** to a diastereomeric selenoxide mixture **60** then enabled either a uracil or cytosine nucleobase to be installed via a Pummerer-type glycosylation, furnishing the β -anomers **61** or **63**. 4'-selenouridine **62** was obtained after global deprotection of **61** with 50% aqueous TFA in an overall yield of 12% over 11 steps. 4'-selenocytidine **64** was similarly obtained in an overall yield of 9% over 11 steps. The crystal structure of **62** revealed the non-native ring adopted an unusual C2'-*endo*/C3'-*exo* twist (Southern confirmation), contrary to uridine, which shows a C2'-*exo*/C3'-*endo* twist (Northern conformation). This difference was explained by the introduction of the bulky selenium, whereby stereoelectronic effects observed in 4'-oxanucleosides are outweighed by the size of the heteroatom.

2.3.1. 2'-Substituted-4'-selenoribofuranosyl Pyrimidines

Building on their work in this area, Jeong and co-workers reported the synthesis and biological evaluation of a small library of 2'-substituted 4'-selenoarabinofuranosyl pyrimidine analogues **65–68** (Figure 5) [44,45].





Figure 5. Structures of 2'-substituted 4'-selenoarabinofuranosyl pyrimidine analogues **65–68**.

A 2'-fluoroarabino analogue was obtained via a 2'-position DAST fluorination of intermediate **69**, synthesised from 4'-selenouridine **61** using a four-step process [44]. At first attempt, DAST fluorination of **69** led to the formation of the desired 2'-fluoro product **70** as the minor product in 23% yield, along with a major product, 2,2'-*O*-anhydro nucleoside **71**, in 60% yield (Scheme 9). This ratio was subsequently improved by instead fluorinating an *N*³-benzoyl derivative **72**, giving **73** in 45% yield, now as the major product, and **71** in reduced 30% yield.



Scheme 9. Reagents and conditions: (i) DAST, CH₂Cl₂, -78 °C, 23%; (ii) TESCl, DMAP, CH₂Cl₂; (iii) BzCl, pyridine, DIPEA, 70 °C; (iv) TBAF, THF, 0 °C, 81 % over three steps and (v) DAST, CH₂Cl₂, -78 °C, 45%.

Following global deprotection and uracil to cytosine conversion, four nucleoside analogues **65**–**68** were evaluated against several human cancer cell lines (HCT116, A549, SU638, T47D, PC-3 and K562) and compared to the established anticancer nucleosides **1** and **3** (Table 2). From this study, it was found that 2'-fluoro-4'-selenoarabinocytidine **65** was the most potent analogue, showing even greater potency than **3**, in the majority of cell lines tested.

Table 2. Anticancer activity of 2'-modified 4'-selenoarabino nucleosides **65**–**69** compared to **1** and **3** across several human cancer cell lines.

Compound	IC ₅₀ (μM)					
	HCT116 ^a	A549 ^b	SNU638 ^c	T47D ^d	PC-3 ^e	K562 ^f
65	1.1	0.47	0.14	0.79	0.58	0.63
66	7.13	8.83	4.72	ND	ND	86.6
67	>100	>100	>100	>100	>100	>100
68	>100	>100	>100	>100	>100	>100
3	5.30	1.90	0.15	2.70	55.9	0.05
1	0.01	0.09	ND	ND	0.04	ND

Human cancer cell tissue type ^acolon; ^blung; ^cstomach; ^dbreast; ^eprostate; ^fmyelogenous leukemia.

The group have also reported further 2'-substituted-4'-selenoribofuranosides [46], synthesising a series of pyrimidine 2'-substituted analogues and utilising a 2,2'-*O*-anhydro intermediate to enable regioselective nucleophilic ring opening to the desired *ribo*-configured products (Figure 6). Only a 2'-fluoro *riboselenocytosine* derivative showed significant activity (uracil, thymine and 5-halo uracil derivatives showed no activity up to 100 μM), but this was less potent than the *arabino*-configured analogue previously described [45].




Figure 6. 2'-Substituted-4'-selenoribofuranosyl nucleosides reported by Jeong.

2.4. Carbocyclic Nucleosides

In carbocyclic nucleosides the furanosyl oxygen is replaced by CH_2 , forming a cyclopentane ring. The lack of a hemiaminal linkage between the nucleobase and the sugar leads to an increased chemical stability. Furthermore, due to the lack of a glycosidic bond, these nucleosides show an enhanced resistance towards phosphorylases [47]. Although considered a second generation of nucleoside analogues, there are two naturally occurring carbocyclic nucleosides, aristeromycin **74** and neplanocin A **75** (Figure 7), both of which exhibit substantial biological activity as antitumor agents [47].




Figure 7. Structures of aristeromycin, **74**, and neplanocin A, **75**.

2.4.1. Fluorinated Derivatives of Neplanocin A

In 1988, Driscoll et al. reported that a cytosine analogue (CPE-C) of **75** showed substantial antitumour and antiviral activity [48]. On the basis of these findings, and the continuing clinical success of fluorine containing nucleoside chemotherapeutics [49,50], Jeong et al. subsequently reported the synthesis and biological evaluation of a small library of fluorocyclopentenyl pyrimidine nucleosides **76–79** (Figure 8) [51].





Figure 8. Structures of fluorocyclopentenyl nucleoside analogues **76–79**.

Fluorocyclopentenyl cytosine **76** was synthesised from cyclopentenone **80** (Scheme 10), which was obtained from D-ribose in nine steps using established procedures [52,53]. Iodination of **80** was accomplished by treatment with I_2 and pyridine in THF, to give **81**. Stereo- and regioselective reduction of **76** was achieved using Luche conditions to give **82**, followed by TBDPS protection of the resulting alcohol to deliver **83**. Following lithium-halogen exchange of **83** using *n*-BuLi, electrophilic fluorination was achieved by treatment with *N*-fluorobenzenesulfonimide (NFSI). Anomeric desilylation with TBAF then gave **84** to which a uracil nucleobase was installed via condensation with *N*³-benzoyluracil under Mitsunobu conditions. Finally, global deprotection was achieved by sequential treatment with methanolic ammonia and BBr_3 , providing uracil derivate **85** which was converted to **76** via a standard four-step process.



Scheme 10. Reagents and conditions: (i) I_2 , pyridine, THF, 55%; (ii) $NaBH_4$, $CeCl_3$, MeOH, 93%; (iii) TBDPSCl, imidazole, DMF, 97%; (iv) NFSI, $nBuLi$, THF, $-78\text{ }^\circ\text{C}$, 80%; (v) TBAF, THF, 80%; (vi) N^3 -benzoyluracil, DEAD, PPh_3 , THF; (vii) $NH_3/MeOH$; (viii) BBr_3 , DCM, $-78\text{ }^\circ\text{C}$, 49% over three steps; (ix) Ac_2O , pyridine; (x) $POCl_3$, Et_3N , 1,2,3-triazole; (xi) NH_4OH , 1,4-dioxane and (xii) $NH_3/MeOH$, 40% over four steps.

Of the four nucleoside analogues evaluated (Figure 8), derivative **76** showed significant potency against several human cancer cell lines (Table 3) [51,54,55]. Furthermore, Jeong reported that **76** also showed significant antitumour activity in a nude mouse xenograft model implanted with A549 human lung cancer cells, wherein after 38 days the inhibition of tumour growth was 32% and 58% at 3 and 10 mg/kg doses, respectively [54].

Table 3. Anticancer activity of **76** in human cancer cell lines.

	Cancer Cell Line						
	HTC-116 ^a	MDA-MB-231 ^b	PANC-1 ^c	MCF-7 ^d	A549 ^e	MKN45 ^f	U251 ^g
IC ₅₀ (μM)	0.39	0.18	0.62	0.34	0.34	0.50	0.83

Human cancer cell tissue type ^acolon; ^bbreast; ^cpancreatic; ^dhormone-dependent breast; ^elung; ^fstomach; ^gbrain.

Carbocyclic nucleoside **76** has now been evaluated in more than 100 different cell lines, as well as several xenograft models, showing high potencies against numerous types of cancer, including gemcitabine resistant cell lines [56,57]. Pharmacokinetics and oral bioavailability for **76** were investigated in phase 0 clinical trials, wherein a small cohort of patients were administered a single oral dose (50 mg or 100 mg) of **76**, or a single intravenous dose (20 mg, Table 4) [58]. This study found that the absolute bioavailability for **76** was 56% and 33% for 50 and 100 mg doses, respectively, suggesting it not to be dose-proportional. However, $t_{1/2}$ was found to be 14 h and 21 h for 50 and 100 mg doses, respectively. This may suggest that **76** does exhibit some dose proportionality in some parameters, but not in others and it was noted that this result may be due to the small patient sample size. Analogue **76** is currently in phase II clinical trials for metastatic pancreatic cancer and advanced bladder cancer.


Table 4. Pharmacokinetic data for **76** from phase 0 clinical studies.

Dose (mg)	T _{max} (h)	C _{max} (ng/mL)	t _{1/2} (h)	Oral Bioavailability (%)
20 *	0.3	1144	-	-
50	2.2	303	14	56
100	2.5	311	21	33

* Intravenous.

2.4.2. Norbornane-Derived (C2',C4'-bridged) Carbocyclic Nucleosides


Nencka and colleagues reported the synthesis of norborane, C2',C4'-bridged carbocyclic nucleosides [59], with a hypothesis of locking the nucleoside analogue in the 2'-exo or North conformation which has garnered attention in the field of carbocyclic antiviral nucleoside analogue development [60]. To access these compounds their synthesis started from known hydroxy ester **86** which was stereochemically inverted at the C1 position via an oxidation-reduction sequence to deliver **87** (Scheme 11). This enabled the desired stereochemistry to be attained at C1 when subsequently inserting an azide by nucleophilic displacement of a mesylate to afford **88** which was reduced to give **89**. With scaffold **89** in hand, the group then elaborated the amine at C1 to several purine and pyrimidine nucleobase forms (including C6 purine analogues) and evaluated their cytotoxic potential. From this series, derivatives **90** and **91** showed IC₅₀ activities below 100 μM in human T-lymphocyte (CEM) cells (IC₅₀ = 88 μM for **90** and IC₅₀ = 78 μM for **91**), but no significant activity was observed in murine leukemia (L1210) or HeLa cell lines.



Scheme 11. Reagents and conditions: (i) PDC, CH₂Cl₂, 82%; (ii) NaBH₄, MeOH, 88%; (iii) MsCl, pyridine, 99%; (iv) NaN₃, DMF, 115 °C, 92% and (v) LiAlH₄, THF, 59%.

2.4.3. C3',C5'-Bridged Carbocyclic L-Nucleosides

Tănase and colleagues reported the synthesis of an alternative carbocyclic system based on a similar bicyclo[2.2.1]heptane fragment, accessing a small series of 3'-5'-linked L-nucleoside analogues containing C6-amino modifications (Scheme 12) [61]. Their synthesis started from known alcohol **92** and proceeded through azide incorporation (to give **93**) and reduction steps in good yields to deliver amine **94**. The 6-chloropurine ring was then introduced using standard methods and elaborated at C6 with a series of amines via nucleophilic aromatic substitution. This small library was then screened in vitro at a single high dose (10⁻⁵ M) in the full NCI 58 human tumor cell screen panel. Phenethylamine derivative **95** exhibited growth inhibition of 74% on SK-MEL-5 melanoma and UO-31 renal cancer cell lines, but was not deemed sufficiently cytotoxic for studies to proceed further. The group followed up this report with further synthesis of 6-position carbocyclic analogues [62], derived from **94**, noting that a 6-(4-methoxy-phenethyl)amino group was active, but again the analogue was not progressed further.




Scheme 12. Reagents and conditions: (i) MsCl, pyridine, CH_2Cl_2 , 98%; (ii) NaN_3 , DMF, 110 °C, 91% and (iii) $Pd(OH)_2/C$, MeOH, 87%.

3. 2′-, 3′- and 5′-Furanose Ring Modifications

3.1. 2′-Furanose Modifications

Clofarabine **2** (Figure 1) is a purine nucleoside analogue which, in its corresponding 5′-O-triphosphate form, inhibits both ribonucleotide reductase and DNA polymerases and prevents effective DNA synthesis. Ultimately, this leads to cell apoptosis, particularly in rapidly proliferating and dormant cancer cells. The nucleoside analogue exhibits excellent cytotoxic activity *in vitro*, with an IC_{50} range of 0.028–0.29 μM across a variety of solid tumour and leukaemia cell lines [63], alongside substantial tissue distribution and a half-life of at least 24 h for the active triphosphate metabolite [64]. The use of **2** for treatment of paediatric patients with relapsed or refractory acute lymphoblastic leukaemia was approved by the food and drug administration (FDA) in 2004, the first nucleoside analogue of its kind to be approved in over a decade [65].

In 2010, Sauve and colleagues reported an improved, stereoselective synthesis of **2** [66], in an overall yield of 38% (Scheme 13). This compared favourably to a prior report by ILEX Products Inc., which detailed an overall yield of 14% in six steps, starting from a fully protected ribose derivative. Sauve’s work began from commercially available lactone **96** with 3,5-*O* protection using TIPS0 affording **97**, which was then diastereoselectively fluorinated at the 2-position, obtaining the *arabino*-configured **98** exclusively in 72% yield. Conversion to the anomeric chloride (via hemi-acetal **99**) delivered **100** which was condensed with 2,6-dichloropurine to afford **101** in 66% yield over two steps and a β/α ratio of 3.5:1. Access to **2** was gained following diastereomeric separation of **101**, 4-position ammonolysis and deprotection.




Scheme 13. Reagents and conditions: (i) TIPS0Cl, imidazole, DMF, 92%; (ii) NFSI, LiHMDS, THF, −78 °C, 72%; (iii) DIBAL-H, toluene, −78 °C, 91%; (iv) MsCl, Et_3N , CH_2Cl_2 , quant.; (v) 2,6-dichloropurine, dichloroethane, reflux; (vi) NH_3 /isopropanol, sealed tube, 105 °C, 66% over two steps and (vii) Me_4NF , AcOH, DMF, 90%.

3.2. 2′-O,4′-Bridged Nucleosides

In 2011, Nicolaou and colleagues reported the synthesis and evaluation of a small library of 2′,4′- and 3′,4′-bridged nucleoside analogues, presenting conformationally restrained 3′-endo (North) and

2'-endo (South) systems [67]. These compounds were evaluated for their antiviral, antitumour and antibacterial properties, with 2',4'-bridged purines **105** and **106** showing μM inhibitory activity against CEM ($\text{IC}_{50} = 0.36 \mu\text{M}$ for **105** and $7.6 \mu\text{M}$ for **106**) and Raji ($\text{IC}_{50} = 0.25 \mu\text{M}$ for **105** and $5.8 \mu\text{M}$ for **106**) cancer cell lines.

The synthesis of **105** and **106** derived from a common 4-disubstituted acetate **102**, available using established chemistry from diacetone-D-glucose (Scheme 14) [68,69]. The group used Vorbrüggen glycosylation of either 2,6-diaminopurine or 2,6-dichloro-9H-purine to afford **103** or **104** respectively, with the substitution proceeding in good yields (100% for **103** and 64% for **104**), but noting a requirement to control the amount of *N,O*-bis(trimethylsilyl)acetamide (BSA) used to silylate the purine to 2.5 equivalents. 2',4'-bridged compound **105** was constructed via base mediated cyclisation between C2' and C4', followed by per-benzoylation of purine nitrogen and silicon protecting group removal.




Scheme 14. Reagents and conditions: (i) 2,6-diaminopurine, BSA, TMSOTf, MeCN, 100%; (ii) 2,6-dichloro-9H-purine, BSA, TMSOTf, 64%; (iii) NaOH, THF, 94%; (iv) BzCl, pyridine, 52%; (v) HF, pyridine, 50%; (vi) Allyl(tri-*n*-butyl)tin, PdCl₂(PPh₃)₂, DMF then NaOH, THF, 42%, 2 steps and (vii) HF, pyridine, 87%.

To access allyl substituted 2',4'-system, Stille couplings were completed at positions 2 and 6 of the purine, followed by a similar base-mediated C2' deacetylation and intramolecular cyclisation to afford, after C5'-protecting group removal, **106**. Isomerisation of the allyl group at C6 of the purine ring was observed during the palladium and base-mediated cyclisation steps. The antitumour activities of **105** and **106** were considerably lower than for the known anticancer nucleoside cladribine (CEM $\text{IC}_{50} = 0.5 \text{ nM}$ and Raji $\text{IC}_{50} = 9.0 \text{ nM}$), as was the activity ($\text{IC}_{50} > 10 \mu\text{M}$) of the 2',4'-bridged analogue of cladribine, suggesting that the inclusion of this conformational restraint or the addition of an extra CH₂O unit was enough to remove antitumour properties.

3.3. 3'-Modified Nucleosides

Cheng and co-workers reported an asymmetric synthesis of 2',3'-dideoxy-3'-boronic acid pyrimidine nucleosides [70]. The group used two highly diastereoselective reactions of chiral boronic esters with (dihalomethyl)lithium reagents to install the ultimate stereochemistry required at C3' and C4' (Scheme 15). Starting from (*S,S*)-1,2-dicyclohexyl-1,2-ethanediol (DICHEd) derived boronate ester **107**, homologation was completed with (dichloromethyl)lithium to afford **108**, the newly formed stereochemistry of which was inverted to give boronate ester **109**. The chiral auxiliary component was next switched from DICHEd to a pinanediol (derived from (+)-pinene), delivering **110**. A second diastereoselective homologation was completed with (dibromomethyl)lithium, affording bromo boronic ester **111**. Nucleophilic displacement and inversion of this bromide with allylmagnesium bromide followed by oxidative cleavage of the alkene afforded aldehyde **112**. Hydrogenolysis of the C4-O-Bn group in **112**, concomitant cyclisation and anomeric acetylation delivered the final 2-deoxy-ribo configured scaffold **113**. This was divergently converted to a series of pyrimidine containing nucleoside analogues containing a unique C3'-boronic acid. Unfortunately, biological evaluation of these compounds demonstrated no significant cytotoxicity towards the

HepG2 cell line and all derivatives were observed to undergo gradual hydrolytic cleavage under biological conditions.




Scheme 15. Reagents and conditions: (i) LiCHCl_2 , THF then ZnCl_2 ; (ii) PhCH_2ONa , 97%; (iii) Pinanediol, Et_2O , H_2O , 96%; (iv) CH_2Br_2 , LDA, THF then ZnCl_2 , 83%; (v) $\text{CH}_2\text{CHCH}_2\text{MgBr}$, 86%; (vi) $\text{NaIO}_4/\text{K}_2\text{OsO}_4$, 2,6-lutidine, dioxane/ H_2O , 63%; (vii) H_2 Pd/C, EtOAc, 81% and (viii) Ac_2O , DMAP, CH_2Cl_2 , 98%. Cy = cyclohexyl, T = thymine, 5F-U = 5-fluorouracil, C = cytosine, 5I-U = 5-iodouracil.

More recently, Borbas and colleagues reported a small library of 3'-deoxy-3'-thio substituted *xylofuranosyl* pyrimidines [71]. Utilising a photoinduced thiol-ene reaction the workers were able to effect addition of several different thiols to an appropriately protected 3'-exomethylene ribopyrimidine system. This afforded D-*xylo* configured products in high diastereomeric excess which were shown to have cytostatic activity in the low micromolar range.

3.4. C5'-N-Cyclopropylcarboxamido-C6-amino-C2-alkynylated analogues

In 2017, Mohan et al. synthesised and evaluated a series of C5'-N-cyclopropylcarboxamido-C6-amino-C2-alkynylated purine nucleoside analogues (Scheme 16) [72]. Starting from guanosine, the group accessed C2-aryl iodide derivative **114** which was oxidised at C5' using KMnO_4 , followed by amide coupling to install the C5'-N-cyclopropylcarboxamido group. Following 2',3'-O-acetonide removal, the C2-iodide underwent a series of divergent Sonogoshira couplings to deliver a library of seven purine analogues. From this library, compounds **115** and **116** showed in vitro cytotoxic effects comparable to doxorubicin against human breast (MDA-MB-2312) and human colon (Caco2) cell lines.



Scheme 16. Reagents and conditions: (i) KMnO_4 , KOH, H_2O , 79%; (ii) EDC, HOBT, cyclopropylamine, Et_3N , DMF, 70%; (iii) 50% Aq. HCO_2H , 73% and (iv) $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$, CuI, Et_3N , DMF, MeCN, R = **115** = 86%, R = **116** = 85%.

3.5. 5'- β -Hydroxyphosphonate Analogues

In 2014, Peyrottes and co-workers reported their synthesis of a series of β -hydroxyphosphonate nucleosides, targeting 2' and 3' hydroxyl group stereochemistry changes and the introduction of 5-position substituents to the nucleobase (Figure 9) [73]. This report built upon the group's earlier work developing this class of compound as 5'-nucleotidase inhibitors (specifically the cytosolic cN-II 5'-nucleotidase II) [74]. These enzymes catabolise nucleoside 5'-monophosphates and their expression level is of crucial interest for patients undergoing nucleoside analogue chemotherapy, with a higher expression level often associated with a worsened clinical outcome [75].




Figure 9. 5'- β -Hydroxyphosphonate analogues developed by Peyrottes and active compound **117** from this series, Y = NH₂ or OH.

The group completed the synthesis of a library of 32 β -hydroxyphosphonate analogues which included D-allo, D-altro, D-manno and D-gluco configurations of the furanose 2'- and 3'-positions along with C2'-C3' ring opened derivatives. They also installed alkynyl, aryl or ethenylaryl groups at C5 of the cytidine or uridine nucleobase using transition-metal cross-couplings of the corresponding C5-iodide. Biological activity data was obtained using recombinant, purified cN-II with inosine monophosphate as substrate. From the resultant SAR study, cytosine-based analogues were generally more active than their uracil counterparts, but the configurational isomers at C2' or C3' were generally less active than the parent compounds (allofuranose). Modification of the nucleobase was generally well tolerated with an observed K_i value of 1.14 mM for phenylethenyl substituted derivative **117** (Figure 9), noting that high substrate concentrations (in the mM range) are required for activity with cN-II.

3.6. Ferronucleosides

Tucker and colleagues recently described their synthesis of ferronucleosides, an important new development in the field of medicinal bioorganometallic chemistry [76]. Here the furanose ring was exchanged for the five-membered cyclopentadienyl ring of a ferrocene unit, but retained the nucleobase and hydroxymethyl group as key components appended in a 2,3 relationship to the ferrocene core (compounds **118** and **119**, Figure 10). Using thymine or adenine as the nucleobase, the compounds were tested with a control series (where the hydroxymethyl or base component were absent) for their cytostatic activity and compared to established chemotherapeutic agents 5-fluorouracil (5F-U) and *cis*platin. In a proliferation activity assay on three tumour cell lines (L1210, HeLa and CEM), both **118** and **119** had activities in the low μ M range, with **118** and **119** 20 to 50-fold more active than 5F-U in CEM cell cultures (0.9 μ M for **118**, 0.35 μ M for **119** versus 18 μ M for 5F-U). The compounds also performed promisingly in cell growth inhibition (oesophageal cancer cell line) and cellular viability (MTT assay) studies, with the data indicating that both functional groups appended to the ferrocene component were required for optimum cytostatic activity.





Figure 10. Ferronucleosides developed by Tucker.

4. Nucleoside Analogue Prodrugs

4.1. Phosphorodiamidate Prodrugs

In 2018, Slusarczyk's team reported the synthesis of phosphorodiamidates for a series of established anticancer nucleoside analogues [77]. This was envisioned following the successful ProTide technology developed by McGuigan [78], to deliver nucleoside monophosphate analogues into cells and recent reports of phosphorothioamidate systems [79]. A lack of chirality at phosphorous in a phosphorodiamidate was envisioned to confer advantages in not having to resolve diastereomeric mixtures, often a requirement in earlier generations of ProTides, as *S_P* and *R_P* diastereoisomers exhibited markedly different biological activities [80]. Accordingly, seven different anticancer nucleosides (**1**, FUdR, 8-chloroadenosine, fludarabine, AraG, thioinosine and thioguanosine) were converted to their phosphorodiamidate form using a one-pot, two-stage strategy (Scheme 17). First the 5'-OH was converted to a phosphorodichloridate intermediate **120**, followed by double phosphoramidation using an appropriate amino acid ester salt. The panel of analogues were then evaluated *in vitro* against a wide range of solid tumour and haematological cell lines with the potential for this approach confirmed for FUdR and 8-chloroadenosine, where similar or improved inhibitory activities compared to the parent nucleosides were observed. FUdR and its phosphorodiamidate prodrug showed activity in the sub-micromolar range with IC₅₀ values of 0.0046–0.073 μM for FUdR and 0.01–0.40 μM for the FUdR phosphorodiamidates against the wild-type cell lines. Also of note was the inactivity of *arabino* configured analogues, suggesting conversion to the monophosphate might be prevented through their being poor substrates for phosphoramidase activity [81]. Enzymatic studies were undertaken to investigate the bioactivation pathway of this class of nucleoside prodrug with a carboxypeptidase Y assay and ³¹P-NMR confirming the hydrolysis of both esters followed by loss of one phosphoramidate group.



Scheme 17. Reagents and conditions: (i) (Me)₃PO₄, POCl₃ and (ii) Amino acid ester salt, DIPEA. X and Y represent different functional groups for analogue classes e.g. X = Y = F, Base = C = gemcitabine phosphorodiamidate. See Reference [77] for full list of analogues synthesised and evaluated.

4.2. Vitamin E Phosphate Prodrugs

Nucleoside analogue therapy can suffer from inducible and constitutive resistance, which limits the efficacy of the treatment. Isoforms of vitamin E, in particular δ -tocopherol and tocotrienols have displayed anticancer activity. As such, vitamin E phosphate nucleoside analogue prodrugs were envisaged to combat two mechanisms of resistance: i) downregulation of metabolic kinases (deoxycytidine kinase, dCK, in the case of **1**) and ii) nucleoside transport [82,83]. Accordingly,

Daifuku and co-workers synthesised and evaluated four isoforms of vitamin E conjugated to **1** (Figure 11, compounds **121–124**) [84].




Figure 11. Structures of vitamin E-gemcitabine conjugates **121–124**, highlighting changes to the tocopherol or tocotrienol component.

The *in vitro* GI₅₀ of **1**, vitamin E phosphate (VEP) isoforms and compounds **121–124** showed these VEP-gemcitabine prodrugs to exhibit anticancer activity, consistent with their catabolism to vitamin E and gemcitabine monophosphate. Conjugate **122** displayed the best activity with GI₅₀ values <5 μ M in breast MDA (MB-231), non-small cell lung (NCI-H460) and colon (HCT-116) cell lines. The authors suggested this was due to steric hindrance, from the methyl groups proximal to the vitamin E-phosphate bond, reducing the rate of enzymatic cleavage to the monophosphate counterpart.

Prodrugs **122** and **123** were then evaluated in the presence of an inhibitor of nucleoside transport, dipyridamole (DP, Table 5) with the data indicating that both were largely unaffected by the presence of DP in comparison to **1**. This suggested that these prodrugs bypass nucleoside membrane transporters and may thus be beneficial in the treatment of patients with gemcitabine resistant cells.

Table 5. GI₅₀ values of **1**, **122** and **123** in the presence or absence of dipyridamole.

Compound	Cancer Cell Line					
	Breast		Non-Small Cell Lung		Colon	
	MDA-MB-231 (μ M)		NCI-H460 (μ M)		HCT-116 (μ M)	
	DP (–)	DP (20 μ M)	DP (–)	DP (20 μ M)	DP (–)	DP (20 μ M)
1	3.08	56.8	0.02	0.82	0.03	2.39
122	30.3	27.8	7.16	16.0	5.55	12.6
123	17.2	23.3	2.14	1.47	3.07	6.74

In particular, as conjugate **123** displayed significant potency against the three DP(–) cell lines and moderate potency against the DP-dosed cell lines (Table 5), it was selected to further compare activity against **1** in *in vitro* wild-type leukemic CEM cells and CEM cells deficient in dCK. In dCK(–) cells **1** is not phosphorylated, with GI₅₀ values increasing from 0.002 μ M in wild-type to 124.5 μ M in dCK(–). For **123** the GI₅₀ increase was significantly lower than that of **1** (from 0.59 μ M to 19.2 μ M).

Furthermore, the half-life of **123** in mice was shown to be 4 h, a 13.9-fold increase to that of **1** (0.3 h [85]). Overall, this study showed an interesting proof of concept for VEP-1 prodrugs against resistant cancer cell lines. However, further optimisation will be required to obtain promising candidate compounds with increased potencies.

5. Conclusion

Nucleoside analogues are an historically accomplished class of drugs with highly diversifiable scaffolds and proven potential to treat a wide range of cancer cell types. Recent synthetic trends have focused on furanose oxygen substitution with heteroatoms or carbon and fluorocyclopentenyl cytosine is proving a promising new clinical candidate in this regard. Alongside this, examples of templating such modifications onto established nucleoside analogue scaffolds, such as Ara-C and forodesine, are emerging. As further pharmacokinetic and pharmacodynamic parameters are evaluated for these architectures, their pharmaceutical utility will be established. Finally, the continued advancement of prodrug strategies to deliver these compounds more effectively and provide options to treat drug-resistant cancer cell types sets an exciting future for nucleoside analogue chemotherapeutics and the underpinning requirement of chemical synthesis in realising this.

Author Contributions: G.J.M., M.G. and C.B. conceptualised and wrote the manuscript. M.G., C.B., M.S., G.J.M. reviewed the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: Riboscience LLC and Keele University are thanked for PhD scholarship funding to M.G and C.B.

Conflicts of Interest: M.S. is a Founder and owns equity in Riboscience LLC.

References

1. Jordheim, L.P.; Durantel, D.; Zoulim, F.; Dumontet, C. Advances in the development of nucleoside and nucleotide analogues for cancer and viral diseases. *Nat. Rev. Drug. Discov.* **2013**, *12*, 447–464.
2. Clercq, E.D.; Neyts, J. Handbook of Experimental Pharmacology. *Handb. Exp. Pharmacol.* **2009**, 53–84.
3. Chapman, T.R.; Kinsella, T.J. Ribonucleotide reductase inhibitors: A new look at an old target for radiosensitization. *Front. Oncol.* **2012**, *1*, 1–6.
4. Flotho, C.; Claus, R.; Batz, C.; Schneider, M.; Sandrock, I.; Ihde, S.; Plass, C.; Niemeyer, C.M.; Lübbert, M. The DNA methyltransferase inhibitors azacitidine, decitabine and zebularine exert differential effects on cancer gene expression in acute myeloid leukemia cells. *Leukemia* **2009**, *23*, 1019–1028.
5. Galmarini, C.; Mackey, J.; Dumontet, C. Nucleoside analogues: Mechanisms of drug resistance and reversal strategies. *Leukemia* **2001**, *15*, 875–890.
6. Jordheim, L.; Dumontet, C. Review of recent studies on resistance to cytotoxic deoxynucleoside analogues. *Biochim. Biophys. Acta* **2007**, *1776*, 138–159.
7. Yates, M.K.; Seley-Radtke, K.L. The evolution of antiviral nucleoside analogues: A review for chemists and non-chemists. Part II: Complex modifications to the nucleoside scaffold. *Antivir. Res.* **2018**, *162*, 5–21.
8. Seley-Radtke, K.L.; Yates, M.K. The evolution of nucleoside analogue antivirals: A review for chemists and non-chemists. Part 1: Early structural modifications to the nucleoside scaffold. *Antivir. Res.* **2018**, *154*, 66–86.
9. Siegel, D.; Hui, H.C.; Doerffler, E.; Clarke, M.O.; Chun, K.; Zhang, L.; Neville, S.; Carra, E.; Lew, W.; Ross, B.; et al. Discovery and Synthesis of a Phosphoramidate Prodrug of a Pyrrolo[2,1-f][triazin-4-amino] Adenine C-Nucleoside (GS-5734) for the Treatment of Ebola and Emerging Viruses. *J. Med. Chem.* **2017**, *60*, 1648–1661.
10. Warren, T.K.; Jordan, R.; Lo, M.K.; Ray, A.S.; Mackman, R.L.; Soloveva, V.; Siegel, D.; Perron, M.; Bannister, R.; Hui, H.C.; et al. Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature* **2016**, *531*, 381–385.
11. Sheahan, T.P.; Sims, A.C.; Leist, S.R.; Schäfer, A.; Won, J.; Brown, A.J.; Montgomery, S.A.; Hogg, A.; Babush, D.; Clarke, M.O.; et al. Comparative therapeutic efficacy of remdesivir and combination lopinavir, ritonavir, and interferon beta against MERS-CoV. *Nat. Commun.* **2020**, *11*, 222.

12. Cho, A.; Saunders, O.L.; Butler, T.; Zhang, L.; Xu, J.; Vela, J.E.; Feng, J.Y.; Ray, A.S.; Kim, C.U. Synthesis and antiviral activity of a series of 1'-substituted 4-aza-7,9-dideazaadenosine C-nucleosides. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 2705–2707.
13. Serpi, M.; Ferrari, V.; Pertusati, F. Nucleoside Derived Antibiotics to Fight Microbial Drug Resistance: New Utilities for an Established Class of Drugs? *J. Med. Chem.* **2016**, *59*, 10343–10382.
14. Romeo, G.; Chiacchio, U.; Corsaro, A.; Merino, P. Chemical Synthesis of Heterocyclic-Sugar Nucleoside Analogues. *Chem. Rev.* **2010**, *110*, 3337–3370.
15. Roy, B.; Depaix, A.; Périgaud, C.; Peyrottes, S. Recent Trends in Nucleotide Synthesis. *Chem. Rev.* **2016**, *116*, 7854–7897.
16. Kicsák, M.; Mándi, A.; Varga, S.; Herczeg, M.; Batta, G.; Bényei, A.; Borbás, A.; Herczegh, P. *Org. Biomol. Chem.* **2018**, *16*, 393–401.
17. Hernández, D.; Boto, A. Nucleoside Analogues: Synthesis and Biological Properties of Azanucleoside Derivatives. *Eur. J. Org. Chem.* **2014**, *2014*, 2201–2220.
18. Quintás-Cardama, A.; Santos, F.P.S.; Garcia-Manero, G. Therapy with azanucleosides for myelodysplastic syndromes. *Nat. Rev. Clin. Oncol.* **2010**, *7*, 433–444.
19. Bouton, J.; Hecke, K.V.; Calenbergh, S.V. Efficient diastereoselective synthesis of a new class of azanucleosides: 2'-homoazanucleosides. *Tetrahedron* **2017**, *73*, 4307–4316.
20. Bantia, S.; Miller, P.J.; Parker, C.D.; Ananth, S.L.; Horn, L.L.; Kilpatrick, J.M.; Morris, P.E.; Hutchison, T.L.; Montgomery, J.A.; Sandhu, J.S. Purine nucleoside phosphorylase inhibitor BCX-1777 (Immucillin-H)—A novel potent and orally active immunosuppressive agent. *Int. Immunopharmacol.* **2001**, *1*, 1199–1210.
21. Miles, R.W.; Tyler, P.C.; Furneaux, R.H.; Bagdassarian, C.K.; Schramm, V.L. One-third-the-sites transition-state inhibitors for purine nucleoside phosphorylase. *Biochemistry* **1998**, *37*, 8615–8621.
22. Makita, S.; Maeshima, A.M.; Maruyama, D.; Izutsu, K.; Tobinai, K. Forodesine in the treatment of relapsed/refractory peripheral T-cell lymphoma: An evidence-based review. *Oncotargets Ther.* **2018**, *11*, 2287–2293.
23. Al-Kali, A.; Gandhi, V.; Ayoubi, M.; Keating, M.; Ravindi, F. Forodesine: Review of preclinical and clinical data. *Future Oncol.* **2010**, *6*, 1211–1217.
24. Warren, T.K.; Wells, J.; Panchal, R.G.; Stuthman, K.S.; Garza, N.L.; Tongeren, S.A.V.; Dong, L.; Retterer, C.J.; Eaton, B.P.; Pegoraro, G.; et al. Protection against filovirus diseases by a novel broad-spectrum nucleoside analogue BCX4430. *Nature* **2014**, *508*, 402–405.
25. Evans, G.B.; Furneaux, R.H.; Gainsford, G.J.; Schramm, V.L.; Tyler, P.C. Synthesis of Transition State Analogue Inhibitors for Purine Nucleoside Phosphorylase and N-Riboside Hydrolases. *Tetrahedron* **2000**, *56*, 3053–3062.
26. Fleet, G.W.J.; Son, J.C. Polyhydroxylated pyrrolidines from sugar lactones: Synthesis of 1,4-dideoxy-1,4-imino-d-glucitol from d-galactonolactone and syntheses of 1,4-dideoxy-1,4-imino-d-allitol, 1,4-dideoxy-1,4-imino-d-ribitol, and (2s,3r,4s)-3,4-dihydroxyproline from d-gulonolactone. *Tetrahedron* **1988**, *44*, 2637–2647.
27. Kezar, H.S.; Kilpatrick, J.M.; Phillips, D.; Kellogg, D.; Zhang, J.; Morris, P.E. Synthesis and Pharmacokinetic and Pharmacodynamic Evaluation of the Forodesine HCl Analog BCX-3040. *Nucleosides Nucleotides Nucleic Acids* **2005**, *24*, 1817–1830.
28. Wortmann, R.L.; Andres, C.; Kaminska, J.; Mejias, E.; Gelf, E.; Arnold, W.; Rich, K.; Fox, I.H. Purine Nucleoside Phosphorylase Deficiency. *Arthritis Rheum.* **1979**, *22*, 524–531.
29. Witczak, Z.J.; Culhane, J.M. Thiosugars: New perspectives regarding availability and potential biochemical and medicinal applications. *Appl. Microbiol. Biotechnol.* **2005**, *69*, 237–244.
30. Secrist, J.A.; Tiwari, K.N.; Riordan, J.M.; Montgomery, J.A. Synthesis and biological activity of 2'-deoxy-4'-thio pyrimidine nucleosides. *J. Med. Chem.* **1991**, *34*, 2361–2366.
31. Pejanović, V.; Stokić, Z.; Stojanović, B.; Piperski, V.; Popsavin, M.; Popsavin, V. Synthesis and biological evaluation of some novel 4'-Thio-L-ribonucleosides with modified nucleobase moieties. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 1849–1852.
32. Zheng, F.; Zhang, X.-H.; Qiu, X.-L.; Zhang, X.; Qing, F.-L. Synthesis of 1-β-3'-Deoxy-3',3'-difluoro-4'-thionucleosides. *Org. Lett.* **2006**, *8*, 6083–6086.
33. Yoshimura, Y.; Kitano, K.; Yamada, K.; Satoh, H.; Watanabe, M.; Miura, S.; Sakata, S.; Sasaki, T.; Matsuda, A. A Novel Synthesis of 2'-Modified 2'-Deoxy-4'-thiocytidines from D-Glucose. *J. Org. Chem.* **1997**, *62*, 3140–3152.

34. Yoshimura, Y.; Kitano, K.; Satoh, H.; Watanabe, M.; Miura, S.; Sakata, S.; Sasaki, T.; Matsuda, A. A Novel Synthesis of New Antineoplastic 2'-Deoxy-2'-substituted-4'-thiocytidines. *J. Org. Chem.* **1996**, *61*, 822–823.
35. Yoshimura, Y.; Saito, Y.; Natori, Y.; Wakamatsu, H. Synthesis of 4'-Thionucleosides as Antitumor and Antiviral Agents. *Chem. Pharm. Bull.* **2018**, *66*, 139–146.
36. Yoshimura, Y.; Kano, F.; Miyazaki, S.; Ashida, N.; Sakata, S.; Haraguchi, K.; Itoh, Y.; Tanaka, H.; Miyasaka, T. Synthesis and biological evaluation of 1'-C-cyano-pyrimidine nucleosides. *Nucleosides Nucleotides* **1996**, *15*, 305–324.
37. Thottassery, J.V.; Sambandam, V.; Allan, P.W.; Maddry, J.A.; Maxuitenko, Y.Y.; Tiwari, K.; Hollingshead, M.; Parker, W.B. Novel DNA methyltransferase-1 (DNMT1) depleting anticancer nucleosides, 4'-thio-2'-deoxycytidine and 5-aza-4'-thio-2'-deoxycytidine. *Cancer Chemother. Pharmacol.* **2014**, *74*, 291–302.
38. Haraguchi, K.; Kumamoto, H.; Konno, K.; Yagi, H.; Tatano, Y.; Odanaka, Y.; Matsubayashi, S.S.; Snoeck, R.; Andrei, G. Synthesis of 4'-substituted 2'-deoxy-4'-thiocytidines and its evaluation for antineoplastic and antiviral activities. *Tetrahedron* **2019**, *75*, 4542–4555.
39. Haraguchi, K.; Takahashi, H.; Shiina, N.; Horii, C.; Yoshimura, Y.; Nishikawa, A.; Sasakura, E.; Nakamura, K.T.; Tanaka, H. Stereoselective Synthesis of the β -Anomer of 4'-Thionucleosides Based on Electrophilic Glycosidation to 4-Thiofuranoid Glycals. *J. Org. Chem.* **2002**, *67*, 5919–5927.
40. Jayakanthan, K.; Johnston, B.D.; Pinto, B.M. Stereoselective synthesis of 4'-selenonucleosides using the Pummerer glycosylation reaction. *Carbohydr. Res.* **2008**, *343*, 1790–1800.
41. Haraguchi, K.; Shimada, H.; Kimura, K.; Akutsu, G.; Tanaka, H.; Abe, H.; Hamasaki, T.; Baba, M.; Gullen, E.A.; Dutschman, G.E.; et al. Synthesis of 4'-Ethylnyl-2'-deoxy-4'-thioribonucleosides and Discovery of a Highly Potent and Less Toxic NRTI. *ACS. Med. Chem. Lett.* **2011**, *2*, 692–697.
42. Jeong, L.S.; Tosh, D.K.; Kim, H.O.; Wang, T.; Hou, X.; Yun, H.S.; Kwon, Y.; Lee, S.K.; Choi, J.; Zhao, L.X. First Synthesis of 4'-Selenonucleosides Showing Unusual Southern Conformation. *Org. Lett.* **2008**, *10*, 209–212.
43. Varela, O.; Zunszain, P.A. First synthesis of aldopentono-1,4-thiolactones. *J. Org. Chem.* **1993**, *58*, 7860–7864.
44. Jeong, L.S.; Tosh, D.K.; Choi, W.J.; Lee, S.K.; Kang, Y.-J.; Choi, S.; Lee, J.H.; Lee, H.; Lee, H.W.; Kim, H.O. Discovery of a New Template for Anticancer Agents: 2'-deoxy-2'-fluoro-4'-selenoarabinofuranosyl-cytosine (2'-F-4'-Seleno-ara-C). *J. Med. Chem.* **2009**, *52*, 5303–5306.
45. Kim, J.-H.; Yu, J.; Alexander, V.; Choi, J.H.; Song, J.; Lee, H.W.; Kim, H.O.; Choi, J.; Lee, S.K.; Jeong, L.S. Structure–activity relationships of 2'-modified-4'-selenoarabinofuranosyl-pyrimidines as anticancer agents. *Eur. J. Med. Chem.* **2014**, *83*, 208–225.
46. Alexander, V.; Song, J.; Yu, J.; Choi, J.H.; Kim, J.-H.; Lee, S.K.; Choi, W.J.; Jeong, L.S. Synthesis and biological evaluation of 2'-substituted-4'-selenoribofuranosyl pyrimidines as antitumor agents. *Arch. Pharm. Sci. Res.* **2014**, *38*, 966–972.
47. Marquez, V.E. Advances in Antiviral Drug Design. In *Advances in Antiviral Drug Design*; Clercq, E.D., Ed.; 1996; pp. 89–146.
48. Marquez, V.E.; Lim, M.I.; Treanor, S.P.; Plowman, J.; Priest, M.A.; Markovac, A.; Khan, M.S.; Kaskar, B.; Driscoll, J.S. Cyclopentenylcytosine. A carbocyclic nucleoside with antitumor and antiviral properties. *J. Med. Chem.* **1988**, *31*, 1687–1694.
49. Cavaliere, A.; Probst, K.C.; Westwell, A.D.; Slusarczyk, M. Fluorinated nucleosides as an important class of anticancer and antiviral agents. *Future Med. Chem.* **2017**, *9*, 1809–1833.
50. Wang, Q.; Liu, X.; Wang, Q.; Zhang, Y.; Jiang, J.; Guo, X.; Fan, Q.; Zheng, L.; Yu, X.; Wang, N.; et al. FNC, a novel nucleoside analogue inhibits cell proliferation and tumor growth in a variety of human cancer cells. *Biochem. Pharmacol.* **2011**, *81*, 848–855.
51. Jeong, L.S.; Zhao, L.X.; Choi, W.J.; Pal, S.; Park, Y.H.; Lee, S.K.; Chun, M.W.; Lee, Y.B.; Ahn, C.H.; Moon, H.R. Synthesis and Antitumor Activity of Fluorocyclopentenyl-Pyrimidines. *Nucleosides Nucleotides Nucleic Acids* **2007**, *26*, 713–716.
52. Moon, H.R.; Choi, W.J.; Kim, H.O.; Jeong, L.S. Improved and alternative synthesis of d- and l-cyclopentenone derivatives, the versatile intermediates for the synthesis of carbocyclic nucleosides. *Tet. Asymm.* **2002**, *13*, 1189–1193.
53. Choi, W.J.; Moon, H.R.; Kim, H.O.; Yoo, B.N.; Lee, J.A.; Shin, D.H.; Jeong, L.S. Preparative and Stereoselective Synthesis of the Versatile Intermediate for Carbocyclic Nucleosides: Effects of the Bulky Protecting Groups to Enforce Facial Selectivity. *J. Org. Chem.* **2004**, *69*, 2634–2636.

54. Choi, W.J.; Chung, H.-J.; Chandra, G.; Alexander, V.; Zhao, L.X.; Lee, H.W.; Nayak, A.; Majik, M.S.; Kim, H.O.; Kim, J.-H.; et al. Fluorocyclopentenyl-cytosine with Broad Spectrum and Potent Antitumor Activity. *J. Med. Chem.* **2012**, *55*, 4521–4525.
55. Balboni, B.; Hassouni, B.E.; Honeywell, R.J.; Sarkisjan, D.; Giovannetti, E.; Poore, J.; Heaton, C.; Peterson, C.; Benaim, E.; Lee, Y.B.; et al. RX-3117 (fluorocyclopentenyl cytosine): A novel specific antimetabolite for selective cancer treatment. *Expert. Opin. Inv. Drug.* **2019**, *28*, 1–12.
56. Peters, G.J.; Smid, K.; Vecchi, L.; Kathmann, I.; Sarkisjan, D.; Honeywell, R.J.; Losekoot, N.; Ohne, O.; Orbach, A.; Blaugrund, E.; et al. Metabolism, mechanism of action and sensitivity profile of fluorocyclopentenylcytosine (RX-3117; TV-1360). *Investig. New Drugs* **2013**, *31*, 1444–1457.
57. Yang, M.Y.; Lee, Y.B.; Ahn, C.-H.; Kaye, J.; Fine, T.; Kashi, R.; Ohne, O.; Smid, K.; Peters, G.J.; Kim, D.J. A novel cytidine analog, RX-3117, shows potent efficacy in xenograft models, even in tumors that are resistant to gemcitabine. *Anticancer Res.* **2014**, *34*, 6951–6959.
58. Udvaros, I.; Rethy, A.; Lang, I.; Hitre, E.; Peterson, C. A phase 1 exploratory study of RX-3117 to determine oral bioavailability in cancer subjects with solid tumors. *J. Clin. Oncol.* **2015**, *33*, e13545–e13545.
59. Dejmek, M.; Šála, M.; Hřebabecký, H.; Dračinský, M.; Procházková, E.; Chalupská, D.; Klíma, M.; Plačková, P.; Hájek, M.; Andrei, G.; et al. Norbornane-based nucleoside and nucleotide analogues locked in North conformation. *Bioorg. Med. Chem.* **2015**, *23*, 184–191.
60. Marquez, V.E.; Siddiqui, M.A.; Ezzitouni, A.; Russ, P.; Wang, J.; Wagner, R.W.; Matteucci, M.D. Nucleosides with a Twist. Can Fixed Forms of Sugar Ring Pucker Influence Biological Activity in Nucleosides and Oligonucleotides? *J. Med. Chem.* **1996**, *39*, 3739–3747.
61. Tănase, C.I.; Drăghici, C.; Căproiu, M.T.; Shova, S.; Mathe, C.; Cocu, F.G.; Enache, C.; Maganu, M. New carbocyclic nucleoside analogues with a bicyclo[2.2.1]heptane fragment as sugar moiety; Synthesis, X-ray crystallography and anticancer activity. *Bioorg. Med. Chem.* **2014**, *22*, 513–522.
62. Tănase, C.I.; Drăghici, C.; Cojocaru, A.; Galochkina, A.V.; Orshanskaya, J.R.; Zarubaev, V.V.; Shova, S.; Enache, C.; Maganu, M. New carbocyclic N6-substituted adenine and pyrimidine nucleoside analogues with a bicyclo[2.2.1]heptane fragment as sugar moiety; synthesis, antiviral, anticancer activity and X-ray crystallography. *Bioorg. Med. Chem.* **2015**, *23*, 6346–6354.
63. Waud, W.R.; Schmid, S.M.; Montgomery, J.A.; Secrist, J.A. Preclinical Antitumor Activity of 2-Chloro-9-(2-deoxy-2-fluoro-β-D-arabinofuranosyl)adenine (C1-F-Ara-A). *Nucleosides Nucleotides Nucleic Acids* **2000**, *19*, 447–460.
64. Albertioni, F.; Lindemalm, S.; Reichelova, V.; Pettersson, B.; Eriksson, S.; Juliusson, G.; Liliemark, J. Pharmacokinetics of cladribine in plasma and its 5′-monophosphate and 5′-triphosphate in leukemic cells of patients with chronic lymphocytic leukemia. *Clin. Cancer Res.* **1998**, *4*, 653–658.
65. Pui, C.H.; Jeda, S.; Kirkpatrick, P. Clofarabine. *Nat. Rev. Drug. Discov.* **2005**, *4*, 369–370.
66. Cen, Y.; Sauve, A.A. Efficient syntheses of clofarabine and gemcitabine from 2-deoxyribose. *Nucleosides Nucleotides Nucleic Acids* **2010**, *29*, 113–122.
67. Nicolaou, K.C.; Ellery, S.P.; Rivas, F.; Saye, K.; Rogers, E.; Workinger, T.J.; Schallenberger, M.; Tawatao, R.; Montero, A.; Hessell, A.; et al. Synthesis and biological evaluation of 2′,4′- and 3′,4′-bridged nucleoside analogues. *Bioorg. Med. Chem.* **2011**, *19*, 5648–5669.
68. Obika, S.; Morio, K.; Nanbu, D.; Hari, Y.; Itoh, H.; Imanishi, T. Synthesis and conformation of 3′,4′-BNA monomers, 3′-O,4′-C-methylenedibonucleosides. *Tetrahedron* **2002**, *58*, 3039–3049.
69. Youssefyeh, R.D.; Verheyden, J.P.H.; Moffatt, J.G. 4′-Substituted nucleosides. 4. Synthesis of some 4′-hydroxymethyl nucleosides. *J. Org. Chem.* **1979**, *44*, 1301–1309.
70. Kim, B.J.; Zhang, J.; Tan, S.; Matteson, D.S.; Prusoff, W.H.; Cheng, Y.-C. Synthesis and properties of 1-(3′-dihydroxyboryl-2′,3′-dideoxyribosyl)pyrimidines. *Org. Biomol. Chem.* **2012**, *10*, 9349.
71. Bege, M.; Kiss, A.; Kicsák, M.; Bereczki, I.; Baksa, V.; Király, G.; Szemán-Nagy, G.; Szigeti, M.Z.; Herczegh, P.; Borbás, A. Synthesis and Cytostatic Effect of 3′-deoxy-3′-C-Sulfanylmethyl Nucleoside Derivatives with d-xylo Configuration. *Molecules* **2019**, *24*, 2173.
72. Mohan, A.A.; Sharma, G.V.R.; Vidavalur, S. Synthesis, characterization and biological evaluation of C5′-N-cyclopropylcarboxamido-C6-amino-C2-alkynylated purine nucleoside analogues. *Nucleosides Nucleotides Nucleic Acids* **2017**, *36*, 1–15.
73. Meurillon, M.; Marton, Z.; Hospital, A.; Jordheim, L.P.; Béjaud, J.; Lionne, C.; Dumontet, C.; Périgaud, C.; Chaloin, L.; Peyrottes, S. Structure–activity relationships of β-hydroxyphosphonate nucleoside analogues

- as cytosolic 5'-nucleotidase II potential inhibitors: Synthesis, in vitro evaluation and molecular modeling studies. *Eur. J. Med. Chem.* **2014**, *77*, 18–37.
74. Gallier, F.; Lallemand, P.; Meurillon, M.; Jordheim, L.P.; Dumontet, C.; Périgaud, C.; Lionne, C.; Peyrottes, S.; Chaloin, L. Structural Insights into the Inhibition of Cytosolic 5'-Nucleotidase II (cN-II) by Ribonucleoside 5'-Monophosphate Analogues. *PLoS Comput. Biol.* **2011**, *7*, e1002295.
 75. Galmarini, C.M.; Thomas, X.; Graham, K.; Jafaari, A.E.; Cros, E.; Jordheim, L.; Mackey, J.R.; Dumontet, C. Deoxycytidine kinase and cN-II nucleotidase expression in blast cells predict survival in acute myeloid leukaemia patients treated with cytarabine. *Br. J. Haematol.* **2003**, *122*, 53–60.
 76. Nguyen, H.V.; Sallustrau, A.; Balzarini, J.; Bedford, M.R.; Eden, J.C.; Georgousi, N.; Hodges, N.J.; Kedge, J.; Mehellou, Y.; Tselepis, C.; et al. Organometallic nucleoside analogues with ferrocenyl linker groups: Synthesis and cancer cell line studies. *J. Med. Chem.* **2014**, *57*, 5817–5822.
 77. Slusarczyk, M.; Ferrari, V.; Serpi, M.; Gönczy, B.; Balzarini, J.; McGuigan, C. Symmetrical Diamidates as a Class of Phosphate Prodrugs to Deliver the 5'-Monophosphate Forms of Anticancer Nucleoside Analogues. *ChemMedChem* **2018**, *13*, 2305–2316.
 78. Cahard, D.; McGuigan, C.; Balzarini, J. Aryloxy phosphoramidate triesters as pro-tides. *Mini-Rev. Med. Chem.* **2004**, *4*, 371–381.
 79. Liu, W.; Zhang, L.; Zhou, H.; Yang, C.; Miao, Z.; Zhao, Y. Synthesis of Novel Nucleoside Analogue Phosphorothioamidate Prodrugs and in vitro Anticancer Evaluation Against RKO Human Colon Carcinoma Cells. *Nucleosides Nucleotides Nucleic Acids* **2013**, *32*, 161–173.
 80. Sofia, M.J.; Bao, D.; Chang, W.; Du, J.; Nagarathnam, D.; Rachakonda, S.; Reddy, P.G.; Ross, B.S.; Wang, P.; Zhang, H.-R.; et al. Discovery of a β -d -2'-Deoxy-2'- α -fluoro-2'- β - C -methyluridine Nucleotide Prodrug (PSI-7977) for the Treatment of Hepatitis C Virus. *J. Med. Chem.* **2010**, *53*, 7202–7218.
 81. Mehellou, Y.; Valente, R.; Mottram, H.; Walsby, E.; Mills, K.I.; Balzarini, J.; McGuigan, C. Phosphoramidates of 2'-beta-D-arabinouridine (AraU) as phosphate prodrugs; design, synthesis, in vitro activity and metabolism. *Bioorg. Med. Chem.* **2010**, *18*, 2439–2446.
 82. Jordheim, L.P.; Galmarini, C.M.; Dumontet, C. Gemcitabine resistance due to deoxycytidine kinase deficiency can be reverted by fruitfly deoxynucleoside kinase, DmdNK, in human uterine sarcoma cells. *Cancer Chemother. Pharmacol.* **2006**, *58*, 547–554.
 83. Damaraju, V.L.; Damaraju, S.; Young, J.D.; Baldwin, S.A.; Mackey, J.; Sawyer, M.B.; Cass, C.E. Nucleoside anticancer drugs: The role of nucleoside transporters in resistance to cancer chemotherapy. *Oncogene* **2003**, *22*, 7524–7536.
 84. Daifuku, R.; Koratich, M.; Stackhouse, M. Vitamin E Phosphate Nucleoside Prodrugs: A Platform for Intracellular Delivery of Monophosphorylated Nucleosides. *Pharmaceuticals* **2018**, *11*, 16.
 85. Shipley, L.A.; Brown, T.J.; Cornpropst, J.D.; Hamilton, M.; Daniels, W.D.; Culp, H.W. Metabolism and disposition of gemcitabine, an oncolytic deoxycytidine analog, in mice, rats, and dogs. *Drug Metab. Dispos.* **1992**, *20*, 849–855.

