

Supplementary Materials

Scale-up preparation of crocins I and II from *Gardenia Jasminoides* by a two-step chromatographic approach and their inhibitory activity on ATP citrate lyase

Shuguang Guan ^{1,2}, Qiaoli Pu ¹, Yanan Liu ³, Honghong Wu ⁴, Wenbo Yu ¹, Zifeng Pi ^{2,5,*}, Shu Liu ², Fengrui Song ², Jingya Li ³ and De-an Guo ^{1,3,*}

¹ College of Pharmacy, Changchun University of Chinese Medicine, Changchun, 130117, China; gsgwww@126.com (S.G.); Puqiaolily@163.com (Q.P.); a511cs@126.com (W.Y.)

² Changchun Institute of Applied Chemistry, Chinese Academy of Sciences, Changchun, 130022, China; mslab20@ciac.ac.cn (S.L.); songfr@ciac.ac.cn (F.S.)

³ Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai, 201203, China; ynlou@sim.ac.cn (Y.L.); jyli@sim.ac.cn (J.L.)

⁴ University of Chinese Academy of Sciences, Beijing, 100049, China; s18-wuhonghong@sim.ac.cn

⁵ Changchun Sunnitech Co., Ltd., Changchun, 130061, China

* Correspondence: mslab21@ciac.ac.cn (Z.P.); daguo@sim.ac.cn (D.G.); Tel.: +86-21-50271516; Fax: +86-21-50271516 (D.G.)




Figure S1. 400 MHz ¹H-NMR spectrum of compound 1 (in DMSO-*d*₆) .




Figure S2. 100 MHz ^{13}C -NMR spectrum of compound 1 (in $\text{DMSO}-d_6$)




Figure S3. 400 MHz ¹H-NMR spectrum of compound 2 (in DMSO-*d*₆)




Figure S4. 100 MHz ^{13}C -NMR spectrum of compound 2 (in $\text{DMSO-}d_6$) .




Figure 5. ESI-MS spectrum of compound 1.




Figure S6. ESI-MS spectrum of compound 2.