

Supplementary Material

Phenolic Extract from *Aralia nudicaulis* L. Rhizomes Inhibits Cellular Oxidative Stresses

Quentin Lion ¹, Andre Pichette ^{1,2}, Mouadh Mihoub ¹, Vakhtang Mshvildadze ^{1,2} and Jean Legault ^{1,2,*}

- ¹ Laboratoire d'analyse et de séparation des essences végétales (LASEVE), Département des Sciences Fondamentales, Université du Québec à Chicoutimi, 555 Boulevard de l'Université, Chicoutimi, QC G7H 2B1, Canada; Quentin.lion@uliege.be (Q.L.); Andre_Pichette@uqac.ca (A.P.); Mouadh.mihoub1@uqac.ca (M.M.); vakhtang_mshvildadze@uqac.ca (V.M.)
- ² Centre de Recherche sur la Boréale (CREB), Département des Sciences Fondamentales, Université du Québec à Chicoutimi, 555 Boulevard de l'Université, Chicoutimi, QC G7H 2B1, Canada
- * Correspondence: Jean.Legault@uqac.ca; Tel.: +14-18-545-5011 (ext. 2334)




Figure S1. HPLC profiles of FEPC with composition identification.

Table S1. HPLC quantification of FEPC compounds.

	FEPC ($\mu\text{g}/\text{mg}$; $w/w\%$)
<u>Protocatechuic acid</u>	48 ± 1 (4.8%)
<u>Chlorogenic acid</u>	156 ± 1 (15.6%)
Vanillic acid	10 ± 1 (1.0%)
<u>Caffeic acid</u>	153 ± 2 (15.3%)
3-Feruloylquinic acid	28 ± 2 (2.8%)
p-Coumaric acid	4.7 ± 1 (0.47%)
5-Feruloylquinic acid	4.8 ± 1 (0.48%)
Dicafeoylquinic acid	21 ± 2 (2.1%)