

Article



Soy Isoflavones Inhibit Both GPIb-IX Signaling and α IIb β 3 Outside-In Signaling via 14-3-3 ζ in Platelet

Ming Liu¹, Gan Wang², Runjia Xu², Chuanbin Shen^{3,4,5}, Heyu Ni^{3,4,5,6,7,8} and Ren Lai^{2,*}

- ¹ Department of Molecular and Cell Biology, School of Life Sciences, University of Science and Technology of China, Hefei 230027, China; qhdming@mail.ustc.edu.cn
- ² Key Laboratory of Bioactive Peptides, Yunnan Province/Key Laboratory of Animal Models and Human Disease Mechanisms, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650032, China; wang.gan@outlook.com (G.W.); xurunjia1991@163.com (R.X.)
- ³ Department of Laboratory Medicine and Pathobiology, University of Toronto,
- Toronto, ON M5S 1A1, Canada; shenchuanbin@126.com (C.S.); Heyu.Ni@unityhealth.to (H.N.)
- ⁴ Department of Laboratory Medicine, LKSKI-Keenan Research Centre for Biomedical Science, St. Michael's Hospital, Toronto, ON M5B 1W8, Canada
- ⁵ Toronto Platelet Immunobiology Group, Toronto, ON M5B 1W8, Canada
- ⁶ Department of Physiology, University of Toronto, Toronto, ON M5S 1A1, Canada
- 7 Canadian Blood Services Centre for Innovation, Toronto, ON M5G 2M1, Canada
- 8 Department of Medicine, University of Toronto, Toronto, ON M5S 1A1, Canada
- * Correspondence: rlai@mail.kiz.ac.cn

Abstract: Soy diet is thought to help prevent cardiovascular diseases in humans. Isoflavone, which is abundant in soybean and other legumes, has been reported to possess antiplatelet activity and potential antithrombotic effect. Our study aims to elucidate the potential target of soy isoflavone in platelet. The anti-thrombosis formation effect of genistein and daidzein was evaluated in ex vivo perfusion chamber model under low (300 s^{-1}) and high (1800 s^{-1}) shear forces. The effect of genistein and daidzein on platelet aggregation and spreading was evaluated with platelets from both wildtype and *GPlba* deficient mice. The interaction of these soy isoflavone with 14-3-3 ζ was detected by surface plasmon resonance (SPR) and co-immunoprecipitation, and the effect of α Ilb β 3-mediated outside-in signaling transduction was evaluated by western blot. We found both genistein and daidzein showed inhibitory effect on thrombosis formation in perfusion chamber, especially under high shear force (1800 s^{-1}). These soy isoflavone interact with 14-3-3 ζ and inhibited both GPIb-IX and α Ilb β 3-mediated platelet aggregation, integrin-mediated platelet spreading and outside-in signaling transduction. Our findings indicate that 14-3-3 ζ is a novel target of genistein and daidzein. 14-3-3 ζ , an adaptor protein that regulates both GPIb-IX and α Ilb β 3-mediated platelet activation is involved in soy isoflavone mediated platelet inhibition.

Keywords: soy; isoflavone; daidzein; 14-3-3ζ; platelet; GPIb-IX; αIIbβ3

1. Introduction

A systematic analysis for the Global Burden of Disease Study revealed that cardiovascular disease (CVD) has the highest mortality among all the non-communicable diseases [1]. Thrombus is responsible for high-mortality CVD such as myocardial infarction and stroke, and platelets play important roles in the pathogenesis of these diseases [2–4]. The inhibition of platelet function has been used for a long time to prevent and treat CVD [5]. However, anti-platelet drugs such as aspirin and clopidogrel increase bleeding risk and may precipitate a hemorrhagic stroke and therapeutic resistance [6], which greatly limits the use of these drugs.

The well-known correlation between diet and health provides strong evidence that functional foods may maintain or improve health and prevent CVD [6–8]. Soybeans and

Citation: Liu, M.; Wang, G.; Xu, R.; Shen, C.; Ni, H.; Lai, R. Soy Isoflavones Inhibit Both GPIb-IX Signaling and α IIb β 3 Outside-In Signaling via 14-3-3 ζ in Platelet. 2021, 26, 4911. https://doi.org/10.3390/ molecules26164911

Academic Editor: Zia Shariat-Madar

Received: 11 July 2021 Accepted: 10 August 2021 Published: 13 August 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/). related food products have been an integral part of regular diets all over the world, especially in Asian countries [7,9]. Epidemiological studies suggested that soy consumption is associated with a lower incidence of CVD [10–12]. Therefore, based on clinical trials and epidemiological data, the US Food and Drugs Administration also approved a health claim for soy [13]. Natural isoflavones mostly exist in bean family plants, and they are particularly abundant in soybeans. Dietary soy isoflavones rather than soy proteins exert antiplatelet functions in agonist-induced platelet activation [14,15].

Early studies have shown that isoflavones might exert anti-platelet function through cAMP regulation [16], tyrosine kinase [17], calcium messenger [18], and TxA₂ pathway inhibition [19]. Moreover, genistein and daidzein can prolong the time that thrombotic vessel occlusion was produced in the femoral arteries of mice using the photochemical thrombosis model [20], and they have a good protective effect on the death or paralysis of mice caused by the pulmonary embolism model induced by the injection of collagen and epinephrine in the tail vein [21]. Although multiple factors may be involved in isoflavone-mediated platelet inhibition, the direct target of soy isoflavone in platelet needs further investigation. Our previous study revealed 14-3-3 ζ is important in platelet integrin outside-in signaling and is a potential target for antithrombotic therapy without causing significant bleeding. An isoflavone which identified from a library (LR-NP1792; Mendeley Data doi:10. 17632/2kbjft8jd3.1) targets 14-3-3 ζ [22]. In the present study, we reported that soy isoflavones (genistein and daidzein) directly bind to 14-3-3 ζ protein and inhibit both GPIb-IX and α IIb β 3 mediated platelet activation.

2. Results and Discussion

2.1. Genistein and Daidzein Inhibit Platelet Aggregation and Thrombi Formation in an Ex Vivo Thrombosis Model

Many isoflavones and their analogues have been demonstrated to inhibit platelet activation [17,19]. However, the effect of soy isoflavones on real-time thrombus formation under different shear stresses has seldom been studied. Here we found that in heparinized mouse blood, platelet-mediated thrombi formed more rapidly under a higher shear rate (1800 s⁻¹) in collagen-coated perfusion chamber. Both genistein (Figure 1A, CAS registry numbers are 446-72-0) and daidzein (Figure 1B, CAS registry numbers are 486-66-8) inhibited thrombi formation under the shear rate of 300 s⁻¹ (Figure 1C,D) and 1800 s⁻¹ (Figure 1E,F), while genistein showed higher potency. However, these isoflavones were preferred to inhibit the platelet aggregation and thrombosis at higher shear rate (1800 s⁻¹), suggesting these compounds may have an inhibitory effect on platelet mechanosensor GPIb-IX mediated signaling transduction [23,24].



Figure 1. Genistein and daidzein inhibited thrombus formation in perfusion chamber. The structural formula of genistein (**A**) and daidzein (**B**). The representative images (**C**) and quantification (**D**) of platelet adhesion and aggregation at 300 s⁻¹ shear rate for 5 min, the concentration of genistein and daidzein both are 100 μ M. The platelets were incubated with isoflavone compounds and their solvent (DMSO) for 5 min at 37 °C. The representative images (**E**) and quantification (**F**) of platelet adhesion and aggregation at 1800 s⁻¹ for 3 min. The circle represent platelet mean fluorescence intensity (MFI) and the shaded regions represent SEM of three independent experiments. Statistical analysis was performed using non-parametric test with a Dunn's multiple comparison test. Significance was defined as * *p* < 0.05, ** *p* < 0.01. Green dots: fluorescent platelets. Bar = 50 µm.

2.2. Genistein and Daidzein Inhibit GPIb-IX-Mediated Platelet Aggregation

It has been well characterized that von Willebrand factor binds to collagen and promotes thrombus formation under high shear force via GPIb-IX signaling [25–27]. We found that genistein and daidzein significantly reduced platelet-mediated thrombi formation under high shear force (Figure 1D). Therefore, we investigated whether these compounds affect GPIb-IX-mediated platelet aggregation. Genistein (Figure 2A) and daidzein (Figure 2B) concentration-dependently inhibited ristocetin-induced platelet aggregation. However, Ristocetin-induced platelet aggregation involves GPIb-IX-mediated platelet agglutination and subsequent integrin α IIb β 3-mediated platelet aggregation [28]. To exclude integrin-mediated platelet aggregation, we examined the effect of the soy isoflavones on ristocetin-induced platelet agglutination in the presence of eptifibatide (Figure 2C) or EDTA (Figure 2D), which abrogates integrin-dependent ligand binding [29]. Both genistein and daidzein inhibited ristocetin-induced platelet agglutination in the presence of 1 mM eptifibatide or 5 mM EDTA, suggesting these isoflavones inhibit GPIb-IX signaling independent of integrin.



Figure 2. The platelets were incubated with isoflavone compounds and their solvent (DMSO) for 5 min at 37 °C. Genistein and daidzein inhibited GPIb-mediated platelet agglutination. Genistein (**A**) and daidzein (**B**) concentration-dependently inhibited 1.2 mg/mL ristocetin induced platelet aggregation in mice PRP. Genistein and daidzein inhibited ristocetin-induced platelet agglutination (**C**) and 5 mM EDTA (**D**).

2.3. Genistein and Daidzein Inhibit α IIb β 3-Mediated Platelet Aggregation

Platelets from wild type and *GPlba* deficient mice were used to investigate the effect of genistein and daidzein on α IIb β 3-mediated platelet aggregation. We found that not only genistein and daidzein inhibited ADP (Figure 3A), collagen (Figure 3B), thrombin (Figure 3C), and arachidonic acid (Figure 3D) induced platelet aggregation but also high concentration of collagen (Figure 3E) and integrin α IIb β 3 activator anti- α IIb TM (Figure 3F) induced platelet aggregation, suggesting these soy isoflavones might inhibit α IIb β 3 outside-in signaling [22,30,31]. The inhibitory effect of genistein and daidzein on platelet aggregation was not through GPIb-IX signaling, since both of these compounds inhibited ADP (Figure 3G) and collagen (Figure 3H) induced platelet aggregation in *GPIb* α deficit platelets.



Figure 3. Genistein and daidzein inhibited integrin-mediated platelet aggregation. The platelets were incubated with isoflavone compounds and their solvent (DMSO) for 5 min at 37 °C. Genistein and daidzein at 200 μ M inhibited 5 μ M ADPinduced platelet aggregation in PRP (**A**). Genistein and daidzein at 200 μ M inhibited 2 μ g/mL collagen (**B**), 0.04 U/mL thrombin (**C**), and 3 μ M arachidonic acid (**D**)-induced platelet aggregation in washed platelets. Effect of genistein, daidzein, SQ29548 (3 μ M), and apyrase (0.5 U/mL) on 50 μ g/mL collagen (**E**) or 1 μ g/mL anti- α IIbTM (**F**)-induced platelet aggregation in the presence of 100 μ g/mL fibrinogen. Genistein and daidzein inhibited 5 μ M ADP (**G**) and 2 μ g/mL collagen (**H**)-induced platelet aggregation in *GPIb* α -deficent mice. Black curve: control. Red curve: genistein treated. Green curve: daidzein treated. Blue curve: SQ29548 treated. Grey curve: apyrase treated.

2.4. Genistein and Daidzein Inhibit Platelet Spreading on Immobilized Fibrinogen

Integrin α IIb β 3 is critical for platelet adhesion and spreading [30–32]. Therefore, we investigated whether the soy isoflavones affects platelet adhesion and spreading on immobilized fibrinogen. Compared with the control group, the number of platelets adhered to immobilized fibrinogen was not significantly affected by the addition of isoflavones at 100 μ M (Figure 4A,B), while 30 μ M integrilin reduced the adhering number by 66 % (Figure 4A,B). The average spreading area in genistein and daidzein treated platelets was 63.2 and 63.7 % that of the control group, respectively. These suggest that isoflavones inhibit platelet spreading and lamellipodia formation. The inhibitory effect of genistein and daidzein on platelet spreading was not through GPIb-IX signaling, since both of these compounds inhibited platelet spreading in *GPIba* deficit platelets (Figure 4C,D).



Figure 4. Genistein and daidzein inhibited integrin-mediated platelet spreading. (**A**,**B**) Confocal microscopy of platelet adhesion and spreading on immobilized fibrinogen. Washed wild type platelets pre-added with 0.1 % DMSO, 100 μ M genistein (GEN) and daidzein (DAI), 3 μ M integrilin were allowed to spread on immobilized fibrinogen for 40 min at 37°C, then stained with FITC-conjugated phalloidin. Scale bar = 10 μ m. (**C**,**D**) Confocal microscopy of *GPlba*-deficient platelet adhesion and spreading on immobilized fibrinogen. Scale bar = 20 μ m. Platelet adhesion numbers were calculated with 3 random vision fields, and spreading areas were calculated with more than 100 platelets. All data are expressed as mean ± SD from 3 experiments. One-way ANOVA with Dunnett's multiple comparisons test was used to indicate statistically significant differences between groups. *** *p* < 0.001.

2.5. Genistein and Daidzein Bind to 14-3-3ζ

The activation of integrin outside-in signaling relies on adaptor proteins that bind to integrin cytoplasmic tail, including talin, kindlins, c-Src, G α 13, and 14-3-3 ζ [30,31]. Surface plasmon resonance (SPR) measurements indicated that both genistein and daidzein bound to 14-3-3 ζ proteins with the estimated equilibrium dissociation constants (KD) of 5.2 × 10⁻⁵ and 2.1 × 10⁻⁴ M, respectively (Figure 5A). Binding of recombinant 14-3-3 ζ to cytoplasmic domain of integrin β 3 (β 3CT) was significantly inhibited by these isoflavones at 30 μ M (Figure 5B), suggesting the binding of isoflavones functionally blocked the complex formation of 14-3-3 ζ with the cytoplasmic domain of integrin β 3 in vitro (Figure 5B).



Figure 5. Genistein and daidzein bind to 14-3-3 ζ . (**A**) 14-3-3 ζ was immobilized on sensor chip CM-5 by amine coupling. Surface plasmon resonance (SPR) analysis of the interaction of genistein (GEN) and daidzein (DAI) with 14-3-3 ζ respectively. (**B**) GEN and DAI inhibited 14-3-3 ζ binding to integrin β 3 cytoplasmic tail. All data are expressed as mean ± SD from 3 experiments. One-way ANOVA with Dunnett's multiple comparisons test was used to indicate statistically significant differences between groups. * *p* < 0.05.

2.6. Genistein and Daidzein Inhibit 14-3-3ζ-Integrin β3 Complex Formation and Outside-In Signaling Transduction in Platelet

Our previous studies showed that 14-3-3 ζ binds to the cytoplasmic domain of integrin β 3 and facilitate outside-in signaling transduction [22]. Therefore, we investigated whether genistein and daidzein affect 14-3-3 ζ -integrin β 3 complex formation in platelet. With the activation of platelets, binding of 14-3-3 ζ to integrin β 3 was increased by 2.7 times (Figure 6B, line 3). Isoflavones significantly inhibited 14-3-3 ζ -integrin β 3 complex formation during platelet activation, while the total content of 14-3-3 and integrin β 3 in platelet lysate was not changed during the process (Figure 6B). Activation of c-Src is essential for early integrin outside-in signaling and platelet aggregation [30,33]. Isoflavones significantly inhibited the phosphorylation of c-Src Y416 as c-Src inhibitor PP2 did at the early stage of platelet aggregation, while the total c-Src content remained unchanged (Figure 6C,D). Isoflavones also significantly inhibited the activation of AKT pathway (Figure 6C,D), which is considered to be the downstream of platelet integrin outside-in signaling pathway [34]. These results suggested that 14-3-3-integrin β 3 complex disrupted by the soy isoflavones affects platelet integrin outside-in signaling pathway.



Figure 6. Genistein and daidzein suppress 14-3-3ζ-integrin β 3 complex formation and outside-in signaling. (**A**,**B**) Binding of 14-3-3ζ to platelet integrin β 3 was inhibited by isoflavones. Platelet pretreated with 0.1% DMSO, 100 µM isoflavones and 3 µM SQ29548 (SQ) for 5 min at 37 °C, then stimulated with 0.04 U/mL thrombin in an aggregometer. Platelets were lysed for co-immunoprecipitation by anti-integrin β 3 antibody. (**C**,**D**) Isoflavones suppressed c-Src and AKT1 activation in platelet. Platelets were pretreated with DMSO (0.1 %), aspirin (ASP) and isoflavones at 100 µM, PP2 at 10 µM and integrilin at 3 µM for 5 min at 37 °C, then stimulated with 0.04 U/mL thrombin in an aggregometer. Platelets were lysed after 30 s and subjected to western blot analysis. All data are expressed as mean ± SD from 3 experiments. (**E**,**F**) Genistein and daidzein inhibited GPIb-IX and α IIb β 3 mediated signaling transduction. One-way ANOVA with Dunnett's multiple comparisons test was used to indicate statistically significant differences between groups. * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001.

Based on previous studies, the pharmacokinetics of isoflavones in humans have been exhaustively cognized [35–37]. Genistein and daidzein are the most abundant isoflavones in soybeans [38–40], which are absorbed through the intestine and liver relative rapidly, and reach the maximum plasma concentration in 2 and 8 h post-intake, and are excreted in urine as glucuronides [41].

Soy isoflavones have been recognized as platelet antagonist and potential antithrombotic, the direct target of these isoflavones in platelet need to be further investigated. In this study, we identified 14-3-3 ζ as a novel target of genistein and daidzein. These isoflavones inhibited both GPIb-IX and α IIb β 3 signaling transduction mediated by 14-3-3 ζ in platelet (Figure 6E,F).

Early studies have shown that genistein inhibited platelet activation via tyrosine kinase inhibition. However, daidzein, which is not a tyrosine kinase inhibitor, also suppressed platelet aggregation, suggesting the inhibition of tyrosine kinase is not essential for these soy isoflavones mediated platelet inhibition [17,42]. Some flavonoids and their analogues (including genistein) might compete for binding to TxA₂ receptor, abrogated arachidonic acid and collagen-induced platelet responses [19]. However, in the presence of fibrinogen, isoflavones significantly inhibited high concentration collagen and anti- α IIb TM induced platelet aggregation while TxA₂ receptor antagonist SQ29548 and adenosine diphosphatase apyrase did not (Figure 3E,F) [22]. This indicates that another target or pathway is involved in genistein and daidzein mediated platelet inhibition.

High concentration of collagen (50 µg/mL) activates integrin α IIb β 3 outside-in signaling independent of PLC γ 2, G α q, TxA₂ receptor and protein kinase C [43], we found genistein and daidzein significantly inhibited 50 µg/mL collagen-induced platelet aggregation (Figure 3F), suggesting integrin α IIb β 3 outside-in signaling is involved in these soy isoflavones mediated platelet inhibition. The designed peptide anti- α IIb TM directly targets the transmembrane (TM) region of integrin α IIb β 3 to high affinity state and initiates α IIb β 3-mediated outside-in signaling [29,44]. Anti- α IIb TM induced platelet aggregation was suppressed by genistein and daidzein, but not significantly affected by SQ29548 and adenosine diphosphatase apyrase in the presence of fibrinogen (Figure 3F), indicates α IIb β 3 outside-in signaling was indeed inhibited by these isoflavones. The effect of the soy isoflavones on platelet was not relying on GPIb-IX, as the aggregation and the spreading of the *GPIb* α deficient platelet were also inhibited by these isoflavones (Figures 3G,H and 4C,D).

It has been well documented that 14-3-3 ζ is abundant in platelet and plays a critical role in VWF-mediated GPIb-IX signaling transduction [28,44,45]. Our recent study revealed 14-3-3 ζ also forms complex with β 3 integrin and is very important for integrin outside-in signaling transduction, therefore may be a potential target for antithrombotic treatment without significant bleeding side-effect [22]. Library screening and functionality assay have uncovered some isoflavones or analogues interact with 14-3-3 ζ and inhibit platelet aggregation [22], whether genistein and daidzein interact with 14-3-3 ζ and inhibit α IIb β 3-mediated outside-in signaling transduction needs further elucidation. SPR and coimmunoprecipitation experiments showed that these soy isoflavones directly bind to 14-3-3 ζ (Figure 5A) and functionally inhibited 14-3-3 ζ -integrin β 3 complex formation both in vitro (Figure 5B) and in vivo (Figure 6A). The interaction of soy isoflavones with 14-3-3 ζ blocked both GPIb-IX and α IIb β 3 signaling as the isoflavones inhibited GPIb-IX mediated platelet agglutination (Figure 2) as well as α IIb β 3 mediated platelet aggregation (Figure 3). However, this study could not completely exclude that other proteins or pathways may be also involved in soy isoflavones treated platelets, which may synergistically contribute to the inhibition of platelet.

Soya foods play an important role in typical Asian diets. Early studies have shown that intakes of the isoflavones of soybean and other legumes reach 20–50 mg/d or even higher: 102 mg/d[45]. The concentration of isoflavones is 10-60 μ M in adult. This may be one of the reasons why the prevalence of cardiovascular diseases is relatively lower in Asia [46].

3. Materials and Methods

All the animal experimental protocols in this work were authorized by the Institutional Animal Care and Use Committees at Kunming Institute of Zoology, Chinese Academy of Sciences (Approval ID: SMKX-20171118-157). C57BL/6 male mice at age of 6-8 weeks (19–25 g) were purchased from the Laboratory Animal Research Center of Kunming Medical University, Kunming, China.

3.1. GPIba Deficient (GPIba-/-) Mice

The murine *GPIba* gene is composed of two exons. We replaced the entire *GPIba* coding sequence (9.0 kb *Hind*III DNA fragment) with a phosphoglycerate kinase-neomycinresistance (*neor*) cassette (6.7 kb *Hind*III DNA fragment) in murine embryonic stem (ES) cells. Southern analysis of ES cells suggests the successful replacement of the *GPIba* gene, and Northern analysis indicates the complete absence of *GPIba* mRNA in homozygousdeficient offspring mice [47–49].

3.2. Ex Vivo Perfusion Chamber

To study the effect of soy isoflavones on thrombus formation under shear force, we used an ex vivo perfusion chamber system as previously described [50]. Briefly, μ -Slide (μ -Slide VI 0.1, ibidi, Madison, WI, USA) were coated with 100 μ g/mL collagen (NC9533954, Thermo Fisher, Waltham, MA, USA) for 2 h at room temperature. Heparinized whole blood was collected from healthy C57BL/6 male mice, pretreated with the soy isoflavones (200 μ M) for 10 min at 37 °C and fluorescently labelled by DiOC₆ (1 μ M, Sigma Alderich, Billerica, MA, USA). The platelets were incubated with isoflavone compounds and their solvent (DMSO) for 5 min at 37 °C. Then the blood was perfused over the collagen-coated surface using a syringe pump (Harvard Apparatus, USA) under the shear rate of 300 s⁻¹ or 1800 s⁻¹ for 5 or 3 min, respectively. Platelet aggregation and thrombus formation were recorded in real-time with Zeiss Axiovert 135-inverted fluorescence microscope (60 X-W objective). Quantitative dynamics of platelet fluorescence intensity were acquired by SlideBook software (Intelligent Imaging Innovations Inc., Denver, CO, USA).

3.3. Platelet Preparation and Aggregation

Platelet preparation from wild type (WT) and *GPlba* deficient (*GPlba*^{-/-}) mice were performed as we previously described [47,51]. Platelet aggregation assay was performed as before [22,52]. Animal protocol for this study was reviewed and approved by the Animal Care and Use Committee at Kunming Institute of Zoology, Chinese Academy of Sciences.

3.4. Surface Plasmon Resonance (SPR) Analysis

SPR was performed as previously described [53]. Briefly, recombinant 14-3-3 ζ was immobilized on sensor chip CM-5 by amine coupling, genistein and daidzein in HBS-EP+ running buffer was applied to the chip with the flow rate of 10 µL/min and the real-time binding signal was recorded by BIAcore 3000 (GE, Boston, MA, USA). The KD (equilibrium dissociation constant) for binding was calculated using a Langmuir model with the Biacore evaluation software provided by the manufacturer.

3.5. Streptavidin–Magnetic Beads Pulldown Assay

The interaction of 14-3-3 ζ with the cytoplasmic tail of integrin β 3 was carried out as previously described [22]. Briefly, 1 µg biotin-conjugated peptide β 3CT (biotin-NNPLYKEATSTFTNITYRGT) was incubated with streptavidin-conjugate magnetic beads (CST, 5947S). Then the beads were incubated with 1 µg recombinant 14-3-3 ζ in binding buffer (50 mM Tris, pH 7.4, 150 mM NaCl, 10 mM MgCl₂, 10 % (v/v) glycerol) with or without soy isoflavones for 4 h at 4 °C. The protein/peptide was degeneration in 5 × SDS loading buffer (BL502A, Biosharp, Hefei, Anhui, China) and separated by 12 % SDS-PAGE, and further blotted by streptavidin-HRP and 14-3-3 ζ antibody.

3.6. Immunoprecipitation and Western Blot

Platelets (2 × 10⁹ /mL) pretreated with genistein and daidzein were elicited by 0.04 U/mL thrombin for 30 s, then lysed with ice-cold NP-40 lysis buffer (10 mM Tris, pH 7.4, 150 mM NaCl, 10 mM MgCl₂, 1 % (v/v) NP-40, 10 % (v/v) glycerol) with protease inhibitors (P8340, Sigma) and phosphatase inhibitors (P5726, Sigma Alderich, Billerica, MA, USA). The lysates were degenerated for 5 min at 90 °C in SDS loading buffer, subjected to 10 % SDS-PAGE and blotted with c-Src, c-Src pY416, AKT1 and AKT1 pS473 antibodies. For immunoprecipitation experiment, the lysates were incubated with integrin β 3 antibody (2 µg) or isotype-matched control IgG (2 µg) overnight at 4 °C, followed by the addition of protein A conjugated magnetic beads (10002D, Thermo Fisher, Waltham, MA, USA). After 3 times washes with NP-40 lysis buffer, bound proteins on magnetic beads were eluted by

50 mM Glycine (PH 2.8) and degenerated for 10 min at 70 °C in SDS loading buffer for western blot analysis.

3.7. Platelet Spreading and Confocal Microscopy

Platelet spreading was measured as we previously described [54]. Briefly, washed platelets were allowed to adhere and spread on fibrinogen (16088, Cayman, Ann Arbor, MI, USA) coated coverslips at 37 °C in the absence or presence of genistein and daidzein. Then platelets were fixed in 4 % paraformaldehyde at room temperature for 15 min, permeabilized in 0.3 % Triton X-100 (T0694, Amresco, Radnor, PA, USA) for 10 min, and stained with FITC labeled phalloidin (40735ES75, Yeasen, Shanghai, China). Platelet numbers and areas were viewed with confocal microscope A1 MP+ (Nikon, Tokyo, Japan) and measured by ImageJ 1.35 h (National Institute of Mental Health, Bethesda, MD, USA) software.

3.8. Statistical Analysis

Image J 1.35h software was used to analyze and quantify Western blot bands, platelet spreading area and blood clot retraction area; Flow Jo V10 software (Becton, Dickinson & Company, Ashland, OR, USA) was used to analyze the average fluorescence intensity of platelets bound to FITC-labeled fibrinogen. Statistical significance was assessed by Student's *t*-test, and GraphPad Prism 6.1 software (GraphPad Software, San Diego, CA, USA) was used for analysis and graphing. The statistical results were expressed as mean \pm SD, and it is statistically significant when p < 0.05.

4. Conclusions

Our study presents evidence that soy isoflavone genistein and daidzein target 14-3-3 ζ and inhibit both GPIb-IX and α IIb β 3 signaling transduction mediated by 14-3-3 ζ in platelet. Soy isoflavone or isoflavone-rich dietary may potentially contribute to the prophylaxis or treatment of thrombosis or CVD that related with abnormal platelet activation.

Author Contributions: R.L. and C.S. conceived and designed the study. M.L. and R.X. performed experiments and data analysis. M.L., C.S. and H.N. wrote the first draft of manuscript. G.W. contributed to results explanation and critically reviewed the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by funding from National Science Foundation of China (31930015, 32070444 and 31801975), Chinese Academy of Sciences (XDB31000000, 2018YFA0801403, SAJC202103 and K. C. Wong Education Foundation), the Science and Technology Department of Yunnan Province (202002AA100007, 2019FB040, 202001AW070019, and 2019FB127). Principal Investigator: Ren Lai.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Institutional Animal Care and Use Committees at Kunning Institute of Zoology, Chinese Academy of Sciences (protocol code SMKX-20171118-157).

Informed Consent Statement: Not applicable.

Data Availability Statement: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the compounds are available from the authors.

References

- Wang, H.D.N.M.; Allen, C.; Barber, R.M.; Bhutta, Z.A.; Carter, A.; Casey, D.C.; Charlson, F.J.; Chen, A.Z.; Coates, M.M.; et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: A systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016, *388*, 1459–1544.
- 2. McManus, D.D.; Freedman, J.E., MicroRNAs in platelet function and cardiovascular disease. *Nat. Rev. Cardiol.* 2015, 12, 711–717.
- Yang, Y.; Shi, Z.; Reheman, A.; Jin, J.W.; Li, C.; Wang, Y.; Andrews, M.C.; Chen, P.; Zhu, G.; Ling, W. Plant food delphinidin-3glucoside significantly inhibits platelet activation and thrombosis: Novel protective roles against cardiovascular diseases. *PLoS* ONE 2012, 7, e37323.
- 4. Wang, Y.; Andrews, M.; Yang, Y.; Lang, S.; W.; Jin, J.; Cameron-Vendrig, A.; Zhu, G.; Reheman, A.; Ni, H. Platelets in thrombosis and hemostasis: Old topic with new mechanisms. *Cardiovasc. Haematol. Disord. -Drug Targets (Former. Curr. Drug Targets-Cardiovasc. Hematol. Disord.*) 2012, 12, 126–132.
- 5. Michelson, A.D. Antiplatelet therapies for the treatment of cardiovascular disease. Nat. Reviews. Drug Discov. 2010, 9, 154–169.
- 6. Fuentes, E.; Palomo, I. Antiplatelet effects of natural bioactive compounds by multiple targets: Food and drug interactions. *J. Funct. Foods* **2014**, *6*, 73–81.
- 7. Zaheer, K.; Akhtar, M.H. An updated review of dietary isoflavones: Nutrition, processing, bioavailability and impacts on human health. *Crit. Rev. Food Sci.* 2017, 57, 1280–1293.
- 8. Habauzit, V.; Morand, C. Evidence for a protective effect of polyphenols-containing foods on cardiovascular health: An update for clinicians. *Adv. Chronic Dis.* **2012**, *3*, 87–106.
- 9. Xu, R.J.; Fei, S.H.; Chen, L.Y.; Wang, G.; Liu, M.; Zhang, W.S.; Yan, X.W.; Lai, R.; Shen, C.B. 3'-Methoxydaidzein exerts analgesic activity by inhibiting voltage-gated sodium channels. *Chin. J. Nat. Med.* **2019**, *17*, 413–423.
- 10. Sagara, M.; Kanda, T.; NJelekera, M.; Teramoto, T.; Armitage, L.; Birt, N.; Birt, C.; Yamori, Y. Effects of dietary intake of soy protein and isoflavones on cardiovascular disease risk factors in high risk, middle-aged men in Scotland. *J. Am. Coll. Nutr.* **2004**, 23, 85–91.
- Kokubo, Y.; Iso, H.; Ishihara, J.; Okada, K.; Inoue, M.; Tsugane, S.; Group, J.S. Association of dietary intake of soy, beans, and isoflavones with risk of cerebral and myocardial infarctions in Japanese populations: The Japan Public Health Center-based (JPHC) study cohort I. *Circulation* 2007, 116, 2553–2562.
- 12. Gil-Izquierdo, A.; Penalvo, J.L.; Gil, J.I.; Medina, S.; Horcajada, M.N.; Lafay, S.; Silberberg, M.; Llorach, R.; Zafrilla, P.; Garcia-Mora, P.; et al. Soy isoflavones and cardiovascular disease epidemiological, clinical and -omics perspectives. *Curr. Pharm. Biotechnol.* **2012**, *13*, 624–631.
- 13. Rimbach, G.; Boesch-Saadatmandi, C.; Frank, J.; Fuchs, D.; Wenzel, U.; Daniel, H.; Hall, W.L.; Weinberg, P.D. Dietary isoflavones in the prevention of cardiovascular disease—A molecular perspective. *Food Chem. Toxicol.* **2008**, *46*, 1308–1319.
- 14. Williams, J.K.; Clarkson, T.B. Dietary soy isoflavones inhibit in-vivo constrictor responses of coronary arteries to collageninduced platelet activation. *Coron. Artery Dis.* **1998**, *9*, 759–764.
- 15. Schoene, N.W.; Guidry, C.A. Dietary soy isoflavones inhibit activation of rat platelets. J. Nutr. Biochem. 1999, 10, 421-426.
- 16. Beretz, A.; Cazenave, J.P.; Anton, R. Inhibition of aggregation and secretion of human platelets by quercetin and other flavonoids: Structure-activity relationships. *Agents Actions* **1982**, *12*, 382–387.
- 17. Nakashima, S.; Koike, T.; Nozawa, Y. Genistein, a protein tyrosine kinase inhibitor, inhibits thromboxane A2-mediated human platelet responses. *Mol. Pharm.* **1991**, *39*, 475–480.
- 18. Dobrydneva, Y.; Williams, R.L.; Morris, G.Z.; Blackmore, P.F. Dietary phytoestrogens and their synthetic structural analogues as calcium channel blockers in human platelets. *J. Cardiovasc. Pharm.* **2002**, *40*, 399–410.
- 19. Guerrero, J.A.; Lozano, M.L.; Castillo, J.; Benavente-Garcia, O.; Vicente, V.; Rivera, A. Flavonoids inhibit platelet function through binding to the thromboxane A(2) receptor. *J. Thromb. Haemost.* **2005**, *3*, 369–376.
- 20. Kondo, K.; Suzuki, Y.; Ikeda, Y.; Umemura, K. Genistein, an isoflavone included in soy, inhibits thrombotic vessel occlusion in the mouse femoral artery and in vitro platelet aggregation. *Eur. J. Pharm.* **2002**, *455*, 53–57.
- 21. Choo, M.K.; Park, E.K.; Yoon, H.K.; Kim, D.H. Antithrombotic and antiallergic activities of daidzein, a metabolite of puerarin and daidzin produced by human intestinal microflora. *Biol. Pharm. Bull.* **2002**, *25*, 1328–1332.
- 22. Shen, C.; Liu, M.; Xu, R.; Wang, G.; Li, J.; Chen, P.; Ma, W.; Mwangi, J.; Lu, Q.; Duan, Z.; et al. The 14-3-3ζ-c-Src-integrin-β3 complex is vital for platelet activation. *Blood* **2020**, *136*, 974–988.
- 23. Ruggeri, Z.M. Platelet GPIb: Sensing force and responding. Bloodthe J. Am. Soc. Hematol. 2015, 125, 423–424.
- 24. Zhang, W.; Deng, W.; Zhou, L.; Xu, Y.; Yang, W.; Liang, X.; Wang, Y.; Kulman, J.D.; Zhang, X.F.; Li, R. Identification of a juxtamembrane mechanosensitive domain in the platelet mechanosensor glycoprotein Ib-IX complex. *Bloodthe J. Am. Soc. Hematol.* **2015**, *125*, 562–569.
- Wu, D.; Vanhoorelbeke, K.; Cauwenberghs, N.; Meiring, M.; Depraetere, H.; Kotze, H.F.; Deckmyn, H. Inhibition of the von Willebrand (VWF)–collagen interaction by an antihuman VWF monoclonal antibody results in abolition of in vivo arterial platelet thrombus formation in baboons. *Bloodthe J. Am. Soc. Hematol.* 2002, *99*, 3623–3628.
- Flood, V.H.; Schlauderaff, A.C.; Haberichter, S.L.; Slobodianuk, T.L.; Jacobi, P.M.; Bellissimo, D.B.; Christopherson, P.A.; Friedman, K.D.; Gill, J.C.; Hoffmann, R.G. Crucial role for the VWF A1 domain in binding to type IV collagen. *Blood* 2015, 125, 2297–2304.

- 27. Wang, Y.; Gallant, R.C.; Ni, H. Extracellular matrix proteins in the regulation of thrombus formation. *Curr. Opin. Hematol.* **2016**, 23, 280–287.
- Dai, K.S.; Bodnar, R.; Berndt, M.C.; Du, X.P. A critical role for 14-3-3 zeta protein in regulating the VWF binding function of platelet glycoprotein Ib-IX and its therapeutic implications. *Blood* 2005, *106*, 1975–1981.
- Fong, K.P.; Zhu, H.; Span, L.M.; Moore, D.T.; Yoon, K.; Tamura, R.; Yin, H.; DeGrado, W.F.; Bennett, J.S. Directly Activating the Integrin alphaIIbbeta3 Initiates Outside-In Signaling by Causing alphaIIbbeta3 Clustering. J. Biol. Chem. 2016, 291, 11706–11716.
- 30. Durrant, T.N.; van den Bosch, M.T.; Hers, I. Integrin alphaIIbbeta3 outside-in signaling. Blood 2017, 130, 1607–1619.
- 31. Shen, B.; Delaney, M.K.; Du, X. Inside-out, outside-in, and inside-outside-in: G protein signaling in integrin-mediated cell adhesion, spreading, and retraction. *Curr. Opin. Cell Biol.* **2012**, *24*, 600–606.
- 32. Ni, H.; Freedman, J. Platelets in hemostasis and thrombosis: Role of integrins and their ligands. *Transfus. Apher. Sci.* 2003, 28, 257–264.
- 33. Estevez, B.; Du, X.P. New Concepts and Mechanisms of Platelet Activation Signaling. Physiology 2017, 32, 162–177.
- 34. Su, X.; Mi, J.; Yan, J.; Flevaris, P.; Lu, Y.; Liu, H.; Ruan, Z.; Wang, X.; Kieffer, N.; Chen, S.; et al. RGT, a synthetic peptide corresponding to the integrin beta 3 cytoplasmic C-terminal sequence, selectively inhibits outside-in signaling in human platelets by disrupting the interaction of integrin alpha IIb beta 3 with Src kinase. *Blood* **2008**, *112*, 592–602.
- 35. Setchell, K.D.R. The history and basic science development of soy isoflavones. *Menopause* 2017, 24, 1338–1350.
- 36. Setchell, K.D.; Zimmer-Nechemias, L.; Cai, J.; Heubi, J.E. Exposure of infants to phyto-oestrogens from soy-based infant formula. *Lancet* **1997**, *350*, 23–27.
- 37. Setchell, K.D. Soy isoflavones--benefits and risks from nature's selective estrogen receptor modulators (SERMs). J. Am. Coll. Nutr. 2001, 20, 354S–362S.
- Setchell, K.D.; Brown, N.M.; Zimmer-Nechemias, L.; Brashear, W.T.; Wolfe, B.E.; Kirschner, A.S.; Heubi, J.E. Evidence for lack of absorption of soy isoflavone glycosides in humans, supporting the crucial role of intestinal metabolism for bioavailability. *Am. J. Clin. Nutr.* 2002, *76*, 447–453.
- Goodman-Gruen, D.; Kritz-Silverstein, D. Usual dietary isoflavone intake is associated with cardiovascular disease risk factors in postmenopausal women. J. Nutr. 2001, 131, 1202–1206.
- 40. Jaiswal, N.; Akhtar, J.; Singh, S.P.; Ahsan, F. An Overview on Genistein and its Various Formulations. *Drug Res.* **2019**, *69*, 305–313.
- Axelson, M.; Sjövall, J.; Gustafsson, B.E.; Setchell, K.D. Soya—A dietary source of the non-steroidal oestrogen equol in man and animals. J. Endocrinol. 1984, 102, 49–56.
- 42. Akiyama, T.; Ogawara, H. Use and specificity of genistein as inhibitor of protein-tyrosine kinases. *Methods Enzymol.* **1991**, 201, 362–370.
- Cho, M.J.; Liu, J.L.; Pestina, T.I.; Steward, S.A.; Thomas, D.W.; Coffman, T.M.; Wang, D.M.; Jackson, C.W.; Gartner, T.K. The roles Of alpha(IIb)beta(3)-mediated outside-in signal transduction, thromboxane A2, and adenosine diphosphate in collageninduced platelet aggregation. *Blood* 2003, *101*, 2646–2651.
- 44. Yin, H.; Slusky, J.S.; Berger, B.W.; Walters, R.S.; Vilaire, G.; Litvinov, R.I.; Lear, J.D.; Caputo, G.A.; Bennett, J.S.; DeGrado, W.F. Computational design of peptides that target transmembrane helices. *Science* **2007**, *315*, 1817–1822.
- 45. van Erp-Baart, M.A.; Brants, H.A.; Kiely, M.; Mulligan, A.; Turrini, A.; Sermoneta, C.; Kilkkinen, A.; Valsta, L.M. Isoflavone intake in four different European countries: The VENUS approach. *Br. J. Nutr.* **2003**, *89*, S25–S30.
- 46. Nagata, C.; Wada, K.; Tamura, T.; Konishi, K.; Goto, Y.; Koda, S.; Kawachi, T.; Tsuji, M.; Nakamura, K. Dietary soy and natto intake and cardiovascular disease mortality in Japanese adults: The Takayama study. *Am. J. Clin. Nutr.* **2017**, *105*, 426–431.
- 47. Xu, M.; Li, J.; Neves, M.A.D.; Zhu, G.; Carrim, N.; Yu, R.; Gupta, S.; Marshall, J.; Rotstein, O.; Peng, J.; et al. GPIbα is required for platelet-mediated hepatic thrombopoietin generation. *Blood* **2018**, *132*, 622–634.
- 48. Ware, J.; Russell, S.; Ruggeri, Z.M. Generation and rescue of a murine model of platelet dysfunction: The Bernard-Soulier syndrome. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 2803–2808.
- 49. Ware, J.; Russell, S.R.; Marchese, P.; Ruggeri, Z.M. Expression of human platelet glycoprotein Ib alpha in transgenic mice. *J. Biol. Chem.* **1993**, *268*, 8376–8382.
- 50. Lei, X.; Reheman, A.; Hou, Y.; Zhou, H.; Wang, Y.M.; Marshall, A.H.; Liang, C.F.; Dai, X.R.; Li, B.X.; Vanhoorelbeke, K.; et al. Anfibatide, a novel GPIb complex antagonist, inhibits platelet adhesion and thrombus formation in vitro and in vivo in murine models of thrombosis. *Thromb. Haemost.* **2014**, *111*, 279–289.
- Li, J.; Van Der Wal, D.E.; Zhu, G.; Xu, M.; Yougbare, I.; Ma, L.; Vadasz, B.; Carrim, N.; Grozovsky, R.; Ruan, M. Desialylation is a mechanism of Fc-independent platelet clearance and a therapeutic target in immune thrombocytopenia. *Nat. Commun.* 2015, 6, 7737.
- 52. Long, C.; Liu, M.; Tian, H.; Li, Y.; Wu, F.; Mwangi, J.; Lu, Q.; Mohamed Abd El-Aziz, T.; Lai, R.; Shen, C. Potential Role of Platelet-Activating C-Type Lectin-Like Proteins in Viper Envenomation Induced Thrombotic Microangiopathy Symptom. *Toxins* **2020**, *12*, 749.

- Jin, L.; Guo, X.M.; Shen, C.B.; Hao, X.; Sun, P.; Li, P.P.; Xu, T.; Hu, C.M.; Rose, O.; Zhou, H.N.; et al. Salivary factor LTRIN from Aedes aegypti facilitates the transmission of Zika virus by interfering with the lymphotoxin-beta receptor. *Nat. Immunol.* 2018, 19, 342–353.
- 54. Shen, C.; Liu, M.; Tian, H.; Li, J.; Xu, R.; Mwangi, J.; Lu, Q.; Hao, X.; Lai, R. Conformation-Specific Blockade of αIIbβ3 by a Non-RGD Peptide to Inhibit Platelet Activation without Causing Significant Bleeding and Thrombocytopenia. *Thromb. Haemost.* 2020, 120, 1432–1441.