

Review

# An Update on Phytochemicals and Pharmacological Activities of the Genus *Persicaria* and *Polygonum*

Gisela Seimandi <sup>1,†</sup>, Norma Álvarez <sup>1,†</sup>, María Inés Stegmayer <sup>1</sup>, Laura Fernández <sup>1</sup>, Verónica Ruiz <sup>1,\*</sup>,  
María Alejandra Favaro <sup>1</sup> and Marcos Derita <sup>1,2,\*</sup>

<sup>1</sup> ICiAgro Litoral, CONICET, Facultad de Ciencias Agrarias, Universidad Nacional del Litoral, Kreder 2805, Esperanza 3080HOF, Argentina; giselaseimandi@hotmail.com.ar (G.S.); nalvarez@fca.unl.edu.ar (N.Á.); mistegmayer@gmail.com (M.I.S.); laurafernandez1@gmail.com (L.F.); mfavaro@fca.unl.edu.ar (M.A.F.)

<sup>2</sup> Farmacognosia, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, Rosario S2002LRK, Argentina

\* Correspondence: vruiz@fca.unl.edu.ar (V.R.); mderita@fbioyf.unr.edu.ar (M.D.);  
Tel.: +54-93415317769 (M.D.)

† These authors contributed equally to this work.

**Citation:** Seimandi, G.; Álvarez, N.; Stegmayer, M.I.; Fernández, L.; Ruiz, V.; Favaro, M.A.; Derita, M. An Update on Phytochemicals and Pharmacological Activities of the Genus *Persicaria* and *Polygonum*. *Molecules* **2021**, *26*, 5956. <https://doi.org/10.3390/molecules26195956>

Academic Editor: Claudio Ferrante

Received: 1 September 2021

Accepted: 25 September 2021

Published: 1 October 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

**Abstract:** The discovery of new pharmaceutical identities, particularly anti-infective agents, represents an urgent need due to the increase in immunocompromised patients and the ineffectiveness/toxicity of the drugs currently used. The scientific community has recognized in the last decades the importance of the plant kingdom as a huge source of novel molecules which could act against different type of infections or illness. However, the great diversity of plant species makes it difficult to select them with probabilities of success, adding to the fact that existing information is difficult to find, it is atomized or disordered. *Persicaria* and *Polygonum* constitute two of the main representatives of the Polygonaceae family, which have been extensively used in traditional medicine worldwide. Important and structurally diverse bioactive compounds have been isolated from these genera of wild plants; among them, sesquiterpenes and flavonoids should be remarked. In this article, we firstly mention all the species reported with pharmacological use and their geographical distribution. Moreover, a number of tables which summarize an update detailing the type of natural product (extract or isolated compound), applied doses, displayed bioassays and the results obtained for the main bioactivities of these genera cited in the literature during the past 40 years. Antimicrobial, antioxidant, analgesic and anti-inflammatory, antinociceptive, anticancer, antiviral, antiparasitic, anti-diabetic, antipyretic, hepatoprotective, diuretic, gastroprotective and neuropharmacological activities were explored and reviewed in this work, concluding that both genera could be the source for upcoming molecules to treat different human diseases.

**Keywords:** natural products; *Persicaria*; *Polygonum*; infectious diseases; bioactive compounds; pharmacological activities

## 1. Introduction

Infectious diseases are caused by pathogenic microorganisms such as bacteria, viruses, parasites or fungi and can be transmitted, directly or indirectly, from one person to another. In the last decades, the rapid population growth, poverty, urban migration, international travels and environment changes have increased the exposure to several infectious agents [1]. According to the World Health Organization's estimations, infectious diseases constitute one-third of all deaths in the world. Despite the pharmaceutical efforts to design new antibiotics for the treatment of these diseases, the acquisition of microorganisms' resistance represents a worldwide concern, and this is attributed to the indiscriminate and improper use of current antimicrobial drugs [2]. For this reason, there is a

need to search for alternative anti-infective substances, for example, antimicrobials of plant origin.

Plants have developed different active principles for defense towards the pathogens' attacks, called secondary metabolites. Phenolic compounds (e.g., coumarins and phytoalexins), terpenoids (e.g., monoterpenes, saponins, steroids) and nitrogen compounds (e.g., alkaloids and lectins) are some secondary metabolites with biocidal capacity against human pathogens [3,4]. It has been estimated that 14–28% of higher plant species are used medicinally [5]. The use of antimicrobials of plant origin has many advantages [6]: they have no secondary effects, better patient tolerance, are less expensive, increase the bioavailability of free agents and demonstrate adequate therapeutic effect with relatively small doses.

The genus *Polygonum* comprises 250 species (20 represented in Argentina) of annual and perennial herbs distributed from the northern temperate to tropical and subtropical regions, preferably in low and humid areas, on the banks of streams and rivers. Nowadays, these species are classified into the *Persicaria* and *Polygonum* genera, according to botanical and phytochemical characteristics. Particularly, flavonoids and sesquiterpenes have played an important role in the systematics of *Polygonum* species as a chemotaxonomic marker and contributed to the regrouping of these species into *Persicaria* and *Polygonum* genera [7]. Species of *Persicaria* and *Polygonum* show different biological properties for, e.g., antiseptic, antibiotic [8], antinociceptive [9], antifungal [8,10,11], diuretic, antirheumatic, astringent [12,13] uses, and for the treatment of external infections such dermatoses, ulcers and sores [13,14].

The selection of the manuscripts for this review was based on the following inclusion criteria: articles published in English in the last three decades, with the keywords *Persicaria*, *Polygonum* and anti-infective in the title, abstract or full text. The authors firstly selected articles according to the title, then to the abstract and then through an analysis of the publication full text. The resulting articles were manually reviewed with the goal of identifying and excluding the works that did not fit the criteria described above. For scientific names of species, The Plant List updated database nomenclature was followed ([www.theplantlist.org](http://www.theplantlist.org)).

## 2. Pharmacological Uses and Phytochemical Composition of *Persicaria* and *Polygonum* Species

The *Persicaria* and *Polygonum* genera are known for their wide spectra of properties to treat different diseases (Table 1). This is possible due to the capacity to produce a great variety of secondary metabolites such sesquiterpenes containing dialdehydes as functional groups [15]; flavonoids [7,16]; neoflavonoids [17]; triterpenes [18,19]; lignans [20]; other types of phenolic compounds not included among flavonoids subclass [21]; phenylpropanoids [22,23]; phenolic compounds belonging to the tannins type [24]; coumarins [17,18]; and anthraquinones [23].

**Table 1.** Origin, distribution and pharmacological uses of *Persicaria* and *Polygonum* species.

Species	Origin/Distribution	Pharmacological Activity	Ref.
<i>Persicaria</i> genus			
<i>Persicaria acuminata</i> (Kunth) M.Gómez	From Mexico to South America	Antifungal; Antinociceptive; Anti-malarial	[9,25,26]
<i>Persicaria alpina</i> (All.) H.Gross	Native to Europe and temperate Asia	Anti-inflammatory; Anti-helminthic	[27]
<i>Persicaria amphibia</i> (L.) Delarbre	Native to Europe, Asia, North America, and parts of Africa	Antifungal; Antibacterial; Anti-cancer	[28,29]
<i>Persicaria amplexicaulis</i> (D.Don) Ronse Decr.	Native to China, the Himalayas and Pakistan	Antioxidant; Anti-cancer; Hepatoprotective	[30–34]
<i>Persicaria barbata</i> (L.) H.Hara	Native to Southeast Asia	Anti-inflammatory; Antinociceptive; Anti-cancer; Diuretic; Gastroprotective	[35–38]
<i>Persicaria bistorta</i> (L.) Samp	Native to Europe and Central and West Asia	Antioxidant; Anti-inflammatory; Anti-cancer; Antipyretic; Hepatoprotective; Gastroprotective	[39–47]
<i>Persicaria capitata</i> (Buch.Ham. ex D.Don) H.Gross	Native to China, India, Nepal, Malaysia, Thailand, Vietnam and Sri Lanka	Antibacterial; Anti-inflammatory	[48]
<i>Persicaria chinensis</i> (L.) H. Gross	Native to South Asian regions with sub-tropical and warm climate	Antifungal; Antibacterial; Antioxidant; Anti-inflammatory; Anti-cancer; Antiviral; Anti-helminthic; Gastroprotective	[49–57]
<i>Persicaria decipiens</i> (R.Br.) K.L.Wilson	Native to Australia	Anti-cancer	[58,59]
<i>Persicaria ferruginea</i> (Wedd.) Soják	Native to temperate climates of South America countries	Antifungal; Antibacterial; Antiviral; Anti-trypanocide	[8,26,60,61]
<i>Persicaria glabra</i> (Willd.) M.Gómez	Native to North America and Eurasia	Antibacterial; Antioxidant; Anti-inflammatory; Anti-cancer; Antiviral; Anti-malarial; Anti-leishmanial; Antipyretic; Hepatoprotective; Neuropharmacological	[62–70]
<i>Persicaria hydropiper</i> (L.) Delarbe	Distributed in the northern hemisphere	Antifungal; Antibacterial; Antioxidant; Anti-inflammatory; Antinociceptive; Anti-cancer; Anti-trypanocide; Anti-helminthic; Anti-diabetic; Antipyretic; Neuropharmacological; Gastroprotective	[71–93]

<i>Persicaria hydropiperoides</i> (Michx.) Small	Native to America, from Canada to Argentina and Chile (introduced in Europe)	Antifungal; Antibacterial; Anti-leishmanial	[19,73,94,95]
<i>Persicaria lapathifolia</i> (L.) Delarbre	Native to Europe and Asia (introduced in America)	Antifungal; Antibacterial; Antioxidant; Anti-inflammatory; Anti-cancer; Antiviral; Anti-helminthic; Anti-diabetic; Diuretic	[96–101]
<i>Persicaria limbata</i> (Meisn.) H.Hara	Distributed in the southwest of Africa, Egypt and tropical Asia	Anti-cancer	[102,103]
<i>Persicaria maculosa</i> Gray	Native to Europe and Asia (introduced in North America)	Antifungal; Antibacterial; Antioxidant; Neuropharmacological	[11,104–109]
<i>Persicaria minor</i> (Huds.) Opiz	Native to Europe (introduced in Australia and America)	Antibacterial; Antioxidant; Anti-inflammatory; Anti-cancer; Antiviral; Hepatoprotective; Neuropharmacological; Gastroprotective	[110–124]
<i>Persicaria orientalis</i> (L.) Spach	Native to India (naturalized in America)	Antioxidant; Anti-inflammatory; Anti-cancer; Anti-diabetic; Hepatoprotective; Neuropharmacological	[125–133]
<i>Persicaria pulchra</i> (Blume) Soják	Distributed in India and Africa	Hepatoprotective	[134,135]
<i>Persicaria punctata</i> (Elliott) Small	Native to America (from Canada to Argentina and Chile)	Antifungal; Antibacterial; Anti-inflammatory; Antiviral	[136–141]
<i>Persicaria sagittata</i> (L.) H.Gross	Distributed in southeastern North America	Antioxidant	[142]
<i>Persicaria senegalensis</i> (Meisn.) Soják	Distributed in south-central Africa	Antifungal; Antibacterial; Antioxidant; Anti-cancer; Anti-diabetic	[58,108,143,144]
<i>Persicaria stagnina</i> (Buch.Ham. ex Meisn.) Qaiser	Distributed in Pakistan, India, Bangladesh and Myanmar	Anti-inflammatory; Anti-cancer	[37,145]
<i>Persicaria tinctoria</i> (Aiton) H.Gross	Native to Southeast Asia	Antibacterial; Antioxidant; Anti-inflammatory; Anti-cancer; Antiviral	[146–155]
<i>Persicaria vivipara</i> (L.) Ronse Decr.	Distributed in the High Arctic	Anti-inflammatory	[156]
<i>Polygonum arenastrum</i> Boreau	Distributed in Europe, North Africa, Southwest Asia and North America	Antifungal	[157]

<i>Polygonum aviculare</i> L.	Native to Europe; It is distributed all over the world	Antifungal; Antibacterial; Antioxidant; Anti-cancer; Anti-diabetic; Neuropharmacological	[158–166]
<i>Polygonum bellardii</i> All.	Distributed in central-northern Europe, North Africa and Southwest Asia	Antifungal; Antibacterial; Antioxidant; Anti-inflammatory; Anti-cancer	[167,168]
<i>Polygonum cognatum</i> Meisn.	Distributed from central Asia to occidental Asia	Antifungal; Antibacterial; Antioxidant; Anti-cancer; Anti-diabetic	[169,170]
<i>Polygonum equisetiforme</i> Sm.	Distributed in southern Europe, North Africa and Southwest Asia	Antioxidant; Hepatoprotective	[171,172]
<i>Polygonum jucundum</i> Meisn.	Distributed in the Chinese provinces	Anti-inflammatory	[173]
<i>Polygonum maritimum</i> L.	Native to Europe. It is distributed in North Africa, Southwest Asia and North America	Antifungal; Antibacterial; Antioxidant; Anti-inflammatory; Anti-diabetic; Neuropharmacological	[174–178]
<i>Polygonum muricatum</i> Meisn.	Distributed in India, Malaysia and Nepal	Anti-helminthic	[179]
<i>Polygonum paleaceum</i> Wall.	Distributed in China and India	Antioxidant; Anti-inflammatory	[24,180]
<i>Polygonum perfoliatum</i> L.	Native to East Asia (China, Japan, Indonesia, Malaysia, Nepal, Korea and Philippines)	Antibacterial; Anti-inflammatory; Anti-cancer; Antiviral; Hepatoprotective	[20,181–189]
<i>Polygonum plebeium</i> R.Br.	Native to Madagascar, South Asia and New Zealand (introduced in United States and Australia)	Hepatoprotective	[190]
<i>Polygonum pubescens</i> Blume	Native to central-south Asia	Anti-inflammatory; Diuretic	[191]
<i>Polygonum thunbergii</i> Siebold & Zucc.	Native to Southeast Asia (China, India, Japan, Korea and Taiwan)	Anti-cancer	[192]
<i>Polygonum verticillatum</i> Biroli ex Colla		Antinociceptive	[193]
<i>Polygonum viscosum</i> Buch.Ham. ex D. Don	Native to Nepal and widely distributed in Bangladesh, northeast India, Japan and China	Anti-cancer; Antiviral; Anti-helminthic; Neuropharmacological	[194–196]

### 3. Pharmacological Activities of *Persicaria* and *Polygonum* Products: Different Type of Extracts and Compounds Responsible for the Bioactivities

The following paragraphs will summarize, mainly in table format, the major biological activities depicted in the literature for these plant genera, not only for the different type of extracts generated from them but also for the pure compounds isolated. The parts of each species used to obtain the bioactive phytochemicals and a summary of the results, thrown by the displayed bioassays, will be also comment.

#### 3.1. Antimicrobial Activity

Many authors all over the world have investigated the ability of *Persicaria* and *Polygonum* species to treat human fungal and bacterial pathogens. Table 2 summarizes the plants which show antimicrobial capacities against different fungal and bacterial strains.

**Table 2.** Antimicrobial activity of *Persicaria* and *Polygonum* species against human fungal and bacterial pathogens. Extracts, compounds, part of plant used, antimicrobial activity, standard drug and references are shown. Extracts: DCM (dichloromethane); EtOH (ethanol); MeOH (methanol); Chl (chloroform); Hex (hexane); EtOAc (ethyl acetate); But (butanol). Part used: L (leaves); F (flowers); R (roots); Sp (sprouts); S (seeds); St (stems); WP (whole plant); AP (aerial parts).

Species	Extract/Isolated Compounds	Part use	Pathogen	Growth Inhibition (mm or %)/MIC/IC <sub>50</sub>	Standard Drug	Ref.
<i>P. acuminata</i>	Polygodial (isolated from DCM extract)	L	<i>Candida albicans</i> and <i>Cryptococcus neoformans</i>	MIC 3.9–62.5 µg/mL	Amphotericin B (MIC 0.25–0.78 µg/mL)	[25]
	Polygodial, isopolygodial and drimenol (isolated from DCM extract)		<i>Microsporium gypseum</i> , <i>Trichophyton rubrum</i> and <i>T. mentagrophytes</i>	MIC 62.5 µg/mL	Amphotericin B (MIC 0.075–0.12 µg/mL)	
<i>P. amphibia</i>	Aqueous and EtOH	F, L	<i>Staphylococcus aureus</i>	11–14 mm	Chloramphenicol (inhibition of 27 mm)	[28]
<i>P. capitata</i>	Aqueous and EtOH (tannin-enriched and flavonoid-enriched fractions)	WP	<i>S. aureus</i> , <i>Escherichia coli</i> , <i>Neisseria gonorrhoeae</i> , <i>Klebsiella pneumoniae</i> and <i>Proteus mirabilis</i>	MIC 0.0022–1.37 mg/mL (Aqueous extract); 0.375–15 mg/mL (other fractions)	Ciprofloxacin (MIC 0.125–0.625 µg/mL)	[48]
	Aqueous and EtOH (gallic acid, triterpenoid and steroid-enriched fractions)		<i>N. gonorrhoeae</i>	MIC 0.375–4 mg/mL		
<i>P. chinensis</i>	MeOH, aqueous, Chl and petroleum ether extract	L	<i>C. albicans</i> and <i>C. krusei</i>	7–18.67 mm	Fluconazole, 30 µg/disc (19.67–20.33 mm)	[49]
			<i>K. pneumoniae</i> , <i>Bacillus cereus</i> , <i>Streptococcus viridians</i> , <i>Corynebacterium diphtheriae</i> , <i>Enterobacter aerogenes</i> , <i>Pseudomonas aeruginosa</i> and <i>Corynebacterium diphtheriae</i>	7.33–22.33 mm	Amoxicillin, 10 µg/disc (13–30.67 mm)	
	MeOH, aqueous, Chl and petroleum ether extract	WP	<i>C. albicans</i> , <i>T. rubrum</i> , <i>Aspergillus niger</i> , <i>A. flavus</i> and <i>Cryptococcus neoformans</i>	11–21 mm (MIC 250–500 µg/mL)	Amphotericin B (22–28 mm, MIC 30 µg/mL)	
<i>K. pneumoniae</i> , <i>P. aeruginosa</i> , <i>Bacillus coagulans</i> , <i>B. subtilis</i> , <i>B. megaterium</i> , <i>B. aerogenes</i> , <i>Lactobacillus leichmanii</i> and <i>Salmonella typhi</i>	12–15 mm (MIC 250–500 µg/mL)	Tetracycline (25–38 mm, MIC 15.5–31.5 µg/mL)				

	Cardamonin (isolated from DCM extract)		<i>Epidermophyton floccosum</i>	MIC 6.2 µg/mL		[8]
<i>P. ferruginea</i>	Crude MeOH extract, sub-extracts ( <i>n</i> -Hex and DCM) and Pashanone isolated from DCM extract	AP	<i>E. floccosum</i> , <i>M. gypseum</i> , <i>T. mentagrophytes</i> and <i>T. rubrum</i>	MIC 25–125 µg/mL	Amphotericin B (MIC 0.4–0.75 µg/mL); Terbinafine (MIC 0.004–0.04 µg/mL)	
	Pashanone (isolated from DCM)		<i>C. albicans</i> , <i>C. neoformans</i> and <i>Saccharomyces cerevisiae</i>	MIC 25–50 µg/mL		
	MeOH and EtOAc extracts	WP	<i>B. subtilis</i> and <i>Proteus vulgaris</i>	4–7 mm (MIC of 0.5–1 mg/mL)		[62]
<i>P. glabra</i>	2-methoxy-5-oxo-2,5-dihydrofuran-3-yl (2 <i>E</i> )-(-)-3-phenylprop-2-enoate, 3-hydroxy-5-methoxystilbene and (-)-pinocembrin (isolated from MeOH extract)	AP	<i>Mycobacterium tuberculosis</i>	IC <sub>50</sub> values of 2.27, 3.33 and 1.21 µg/mL, respectively		[63]
	Confertifolin (isolated from essential oils)		<i>E. floccosum</i> and <i>Curvularia lunata</i>	MIC 7.81 µg/mL	For fungi: Ketoconazole (MIC < 12.5 µg/mL); for bacteria: Streptomycin (MIC 25 µg/mL)	[71]
	Drimenol (isolated from essential oils)	L	<i>T. mentagrophytes</i> , <i>T. rubrum</i> , <i>T. simii</i> and <i>A. niger</i>	MIC 31.25 µg/mL MIC < 12.5 µg/mL		[72]
			<i>A. niger</i> , <i>A. flavus</i> , <i>A. fumigatus</i> and <i>T. rubrum</i>	17–20 mm		
<i>P. hydropiper</i>	Chl extract	R	<i>B. subtilis</i> , <i>B. megaterium</i> , <i>S. aureus</i> , <i>E. aerogenes</i> , <i>E. coli</i> , <i>P. aeruginosa</i> , <i>S. typhi</i> and <i>Shigella sonnei</i>	22–25 mm (MIC 16–64 µg/mL)	Kanamycin, 30 µg/disc (32.7–35 mm, MIC 2–8 µg/mL)	
	Polygodial	Sp	<i>C. albicans</i> , <i>C. krusei</i> , <i>C. neoformans</i> , <i>S. cerevisiae</i> , <i>T. mentagrophytes</i> , <i>T. rubrum</i> and <i>Penicillium marneffeii</i>	MIC 0.39–6.25 µg/mL	Amphotericin B (MICs 0.2–1.56 µg/mL).	[73]
	Crude MeOH extract and its fractions (saponins, chloroform and ethyl acetate sub-extract)	WP	<i>Proteus mirabilis</i>	20–30 mm (MIC 25–40.5 µg/mL)	Ceftriaxone (35 mm, MIC 10 µg/mL)	[74]
	Acetone and EtOH extracts	L, St	<i>K. pneumoniae</i> , <i>Haemophilus influenzae</i> , <i>Morganella morganii</i>	10–19 mm	Ampicillin 10 µg (17–20 mm)	[75]

<i>P. hydropiperoides</i>	Polygodial (isolated from MeOH extract)	F, Sp	<i>C. albicans, C. krusei, C. neoformans, C. utilis, S. cerevisiae, T. mentagrophytes, T. rubrum, P. marneffeii and P. chrysogenum</i>	MIC 0.78–12.5 µg/mL	Amphotericin B (MIC 0.2–1.56 µg/mL)	[73]
	EtOH extract	L	<i>S. aureus</i>	9 mm		[94]
	Triterpenoids, tannins and flavonoids (isolated from MeOH extract)	F, L	<i>Salmonella typhimurium</i>	16–19 mm	Thymol	[19]
<i>P. lapathifolia</i>	Pinostrobin chalcone and Pashanone (isolated from crude extract)	S	<i>Trichoderma sp., Fusarium sp., Aspergillus sp. and Penicillium sp.</i>	10–22 mm	Clotrimazole (15–23 mm)	[96]
	Flavokawin and Pashanone (isolated from crude extract)		<i>E. coli</i>	12–18 mm		
	Flavokawin and Pashanone (isolated from crude extract)		<i>S. aureus</i>	9–13 mm	Gentamycin (17–19 mm)	
<i>P. maculosa</i>	DCM extract and isolated compounds (polygodial, isopolygodial and pinostrobin)	AP	<i>M. gypseum, T. rubrum and T. mentagrophytes</i>	MIC 7.8–62.5 µg/mL	Ketoconazole (MIC 0.02–0.25 µg/mL) and Amphotericin (MIC 0.07–0.5 µg/mL)	[11]
	Persilben	WP	<i>Trychophyton sp.</i>	MIC 125–250 mg/L		[104]
	MeOH extract	L	<i>E. coli</i> <i>S. typhi</i> and <i>P. aeruginosa</i>	100% 42–49%	Ampicillin, 1 mg/mL	[105]
	Quinic, gallic and chlorogenic acid and quercetin 3-O-β-D-glucopyranoside (isolated from EtOH extract)	AP	<i>P. aeruginosa</i> and <i>Salmonella enterica</i>	Inhibition of biofilm formation of 50%	Dimethyl sulfoxide 0.1% (DMSO)	[106]
	MeOH and EtOH extracts (50 and 70%)		<i>E. coli, B. subtilis</i> and <i>S. aureus</i>	47%		[110]
<i>P. minor</i>	MeOH, Chl and petroleum ether extracts		<i>Helicobacter pylori</i>	11.9–16.2 mm		[111]
	Aqueous-EtOH (30%) and aqueous (100%) extracts	L	<i>E. faecalis, E. coli</i> and <i>S. aureus</i>	16.45–19.5 mm (concentration of 200 mg/mL)	Penicillin (20.7–25.5 mm at concentration of 10 mg/mL)	[112]
	Hex, DCM and MeOH extracts		<i>B. cereus</i>	12.5–14.5 mm (MIC of 1.25–2.5 mg/mL)	Ampicillin (17.5 mm; MIC of 0.1 mg/mL)	[113]
<i>P. punctata</i>	AP	<i>C. albicans, A. niger</i> and <i>Mucor sp.</i>			[136,137]	

	Polygodial (isolated from DCM extract) and DCM extract		<i>B. subtilis</i> , <i>S. aureus</i> and <i>Micrococcus luteus</i>			
	MeOH extract		<i>E. faecalis</i> , <i>S. aureus</i> , <i>B. subtilis</i> and <i>Mycobacterium phlei</i>			[138]
	Isotadeonal and ethyl ether extract		<i>P. aeruginosa</i> and <i>S. aureus</i>	75% (concentration of 100 µg/mL)		[139]
<i>P. senegalensis</i>	Pyrazolines derivates of chalcones	AP	<i>C. krusei</i> , <i>C. neoformans</i> , <i>S. aureus</i> and <i>C. glabrata</i> ,	IC <sub>50</sub> 7.56–13.74 µg/mL	Amphotericin B (IC <sub>50</sub> 0.37–1.38 µg/mL) and Ciprofloxacin (IC <sub>50</sub> 0.09 µg/mL)	[143]
	Hydroalcoholic (50%) extract	L	<i>E. faecalis</i> , <i>B. subtilis</i> and <i>S. aureus</i>	MIC 1.25–5 mg/mL		[144]
	Extract and tryptanthrin isolated from this extract	L	<i>Streptococcus mutans</i> , <i>S. sobrinus</i> , <i>Porphyromonas gingivalis</i> , <i>Campylobacter rectus</i> , <i>Prevotella intermedia</i> and <i>Actinobacillus actinomycetemcomitans</i>	MIC 1.74–3.48 µg/mL (for extract) and 6.25–25 µg/mL (for tryptanthrin)		[146]
<i>P. tinctoria</i>	Kaempferol (isolated from leaves extract)		<i>S. mutans</i> , <i>S. sobrinus</i> , <i>P. gingivalis</i> and <i>P. intermedia</i>	MIC 25–50 µg/mL		
	Tryptanthrin	WP	<i>H. pylori</i>	Inhibited 100% colony formation (dose of 10 µg/mL)	Amoxicillin, clarithromycin and omeprazole	[147]
			<i>C. albicans</i>	MIC 250 µg/mL	Chloramphenicol (MIC 0.156–1.25 µg/mL), amphotericin B and ketoconazole (MIC 0.04–0.31 µg/mL)	[157]
<i>P. arenastrum</i>	MeOH extract	L, St	<i>C. krusei</i>	MIC 62.5–15.63 µg/mL		
	Chl extract	St	<i>A. niger</i> , <i>A. flavus</i> and <i>A. fumigatus</i>	14–18 mm (MIC 1–5 mg/mL)	Cotrimoxazole antibiotic, 10 mg/g (18–34 mm)	[158]
<i>P. aviculare</i>			<i>E. coli</i> , <i>P. mirabilis</i> , <i>P. aeruginosa</i> , <i>S. typhi</i> , <i>S. aureus</i> and <i>B. subtilis</i>	24–28 mm (MIC 8–15 mg/mL)		
	EtOH extract	AP	<i>P. aeruginosa</i> , <i>S. aureus</i> and <i>Acinetobacter baumannii</i>	74–100% (at a concentration 1 mg/mL)		[159]
			<i>C. albicans</i>	11–20 mm (MIC 1–25 mg/mL)	Nystatin antibiotics, 25 µg/disc	[167]
<i>P. bellardii</i>	MeOH and EtOAc extracts, <i>n</i> -Hex, Chl and <i>n</i> -But fractions	AP	<i>S. aureus</i> , <i>B. subtilis</i> , <i>E. coli</i> and <i>P. aeruginosa</i>	11–30 mm (MIC 1–5 mg/mL)	Cefotax, 15–30 µg/disc	

<i>P. cognatum</i>	EtOH extract	L	<i>C. albicans</i>	MIC 2.5 mg/mL		[169]
		WP	<i>S. aureus</i> , <i>P. aeruginosa</i> and <i>E. coli</i>	MIC 0.156–0.625 mg/mL		
<i>P. maritimum</i>	Phenolic compounds (isolated from EtOH extract)	WP	<i>K. pneumoniae</i> , <i>S. aureus</i> , <i>E. coli</i> , <i>B. megatarium</i> , <i>C. albicans</i>	8–10 mm	Ceftriaxone and Nystatin 30 µg/disk (10–11 mm)	[170]
			<i>Penicillium</i> sp., <i>Aspergillus</i> sp., <i>Alternaria alternata</i> and <i>Fusarium semitectum</i>	19–34% (concentration of 1–5 mg/mL)		[174]
		AP	<i>B. cereus</i> , <i>B. subtilis</i> , <i>S. aureus</i> , <i>A. baumannii</i> , <i>E. faecalis</i> , <i>P. mirabilis</i> and <i>Citrobacter freundii</i>	MIC 0.12–4.02 mg/mL		[175]
<i>P. perfoliatum</i>	Crude extract		<i>E. coli</i> and <i>P. aeruginosa</i>	MIC 16.08–64.35 mg/mL		
			<i>S. aureus</i> and <i>Cutibacterium acnes</i>	MIC of 0.25%,	Quercetin (MIC of 0.06–0.3%) and methyl paraben (MIC 0.13–0.25%)	[181]
	Water extract	AP	<i>S. aureus</i> , <i>E. coli</i> , <i>Streptococcus</i> sp., <i>Salmonella</i> sp. and <i>Pasteurella</i> sp.	0.56–21.86 mm (MIC 0.031–0.063 mg/mL)		[182]
	EtOH (75%)		<i>S. aureus</i> , <i>B. subtilis</i> and <i>P. aeruginosa</i>	MIC 5–10 mg/mL		[183]

From the analysis of Table 1, it could be stated that the three main bioassays displayed for detecting antimicrobial activities of different type of extracts of *Polygonum* or *Persicaria* species, as well as their bioactive compounds, include the percentage of microorganism growth inhibition, the determination of IC<sub>50</sub> or the Minimum Inhibitory Concentration of each tested sample (extract or isolated compound). The last one constitutes a standard method for detecting antimicrobial susceptibility, so that the results obtained from carrying out these bioassays should be more convincing. In this sense, *P. acuminata*, *P. ferruginea*, *P. hydropiperoides*, *P. lapathifolia* and *P. arenastrum* were tested only as antifungals against yeasts or filamentous fungi, resulting in *P. acuminata* and *P. ferruginea* being the most active ones (MICs between 3.9 and 125 µg/mL). Authors correlated these promising activities with the presence of the sesquiterpene polygodial in *P. acuminata* extracts [25] and the chalcones cardamomin and pashanone present in *P. ferruginea* extracts [8]. Moreover, *P. chinensis*, *P. hydropiper*, *P. maculosa*, *P. punctata*, *P. senegalensis*, *P. aviculare* and *P. cognatum* resulted in both antifungals and antibacterials, highlighting the MICs values obtained for isolated compounds confertifolin and drimenol from *P. hydropiper* essential oil and polygodial from its chloroformic extract (MICs between 0.39 and 125 µg/mL comparable with standard drugs) [71–75]. Finally, *P. capitata*, *P. glabra*, *P. minor*, *P. tinctoria* and *P. perfoliatum* were reported in the literature only as antibacterials, remarking the high activity against *S. aureus*, *E. coli*, *K. pneumoniae* and *N. gonorrhoeae* of flavonoid-enriched fractions of aqueous extracts of *P. capitata* (MICs comparables to ciprofloxacin) [48]. These results are promising but limited to the in vitro evaluation, and thus, more studies regarding solubilities, absorption, blood distribution, pharmacodynamics, pharmacokinetics and tissue toxicity should be performed before these extracts or compounds could become remedies.

### 3.2. Antioxidant Activity

Species of the *Persicaria* and *Polygonum* genera have a remarkable antioxidant activity, as some compounds can remove the excess of free radicals in bodies to maintain normal metabolisms. Table 3 summarizes the most interesting results found in the literature for the following antioxidant bioassays performed: DPPH (2,2-diphenyl-2-picryl hydroxyl); TEAC (Trolox Equivalent Antioxidant Capacity); CUPRAC (Cupric Reducing Antioxidant Capacity); ABTS ((2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)); NBT (Nitroblue tetrazolium); FRAP (Ferric Reducing Antioxidant Power); ORAC (Oxygen Radical Absorbance Capacity assay); and CCA (Copper Chelating Activity).

**Table 3.** Antioxidant activity of *Persicaria* and *Polygonum* species. Extracts, compounds, part of plant used, antioxidant activity and references are shown. Extracts: DCM (dichloro-methane); EtOH (ethanol); MeOH (methanol); EtOAc (ethyl acetate); But (butanol). Part used: L (leaves); F (flowers); R (roots); Sp (sprouts); Sh (shoots); S (seeds); St (stems); Rh (rhizomes); WP (whole plant); AP (aerial parts). Antioxidant assays: DPPH (2,2-diphenyl-2-picryl hydroxyl); TEAC (Trolox Equivalent Antioxidant Capacity); CUPRAC (Cupric Reducing Antioxidant Capacity); ABTS ((2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)); NBT (Nitroblue tetrazolium); FRAP (Ferric Reducing Antioxidant Power); ORAC (Oxygen Radical Absorbance Capacity assay); CCA (Copper Chelating Activity).

Species	Extract/Isolated Compounds	Part used	Summarized Bioactivity	Ref.
<i>P. equisetiforme</i>	MeOH extract	Sh	DPPH scavenging activity from 12 to 51 mM TRE/g DW. EC <sub>50</sub> of reducing power = 68–210 µg/mL. This capacity was attributed to different compounds such as quinic acid, gallic acid, (+)-catechin, epicatechin, quercetin-3-O-β-D-galactoside, quercetin-3-O-α-L-rhamnoside and cirsiolol.	[171]
<i>P. glabra</i>	Flavonoids, phenols, tannins, terpenoids and reducing sugars (isolated from MeOH extract)	L	DPPH free radicals with inhibitory concentration (IC <sub>50</sub> ) of 79.81 µg/mL.	[64]
<i>P. lapathifolia</i>	Isoquercitrin, hyperoside, quercitrin and taxifolin, gallic acid (isolated from MeOH extract)	AP	At a dose of 5 µM, compounds showed higher antioxidant activity than the standard quercetin (TEAC value of 1.16 µM for compounds and 1.1 µM for standard quercetin). Gallic acid was the most potent scavenger of hydroxyl radicals (inhibition of 70.8%) compared to the standards quercetin and Superoxide Dismutase (SOD) (inhibition of 66.2 and 77.2%, respectively).	[97]
	Flavonol O-glycosides with TMF as an aglycone	AP, S, Sp	DPPH scavenging activity from 500 µmol/g DW, (AP), 100 µmol/g DW (Sp) and 50 µmol/g DW (S).	[148]
<i>P. tinctoria</i>	EtOAc fraction	St	Displayed high antioxidant activity (IC <sub>50</sub> 7.17 µg/mL) with respect to L-ascorbic acid (IC <sub>50</sub> 5.5 µg/mL).	[149]
	MeOH extract	L, F	ABTS radical scavenging of 99.12 and 96.35 M TE/g DW (L and F, respectively). CUPRAC values of 78.37 and 86.22 M TE/g DW (L and F, respectively). DPPH and FRAP inhibition greater than 75% at a concentration of 50 µg/mL.	[150]
<i>P. aviculare</i>	Lyophilized EtOH extract	WP	NBT greater than 90% at a concentration of 5 µg/mL. Extract showed it can protect DNA in hydroxyl-radical-induced DNA strand scission assays.	[160]
<i>P. cognatum</i>	EtOH extract	L	DPPH radical scavenging maximum of 18% and ABTS radical scavenging of 70%. Positive control BHT (DPPH and ABTS scavenging of 80%).	[169]

	EtOH extract	AP	SC <sub>50</sub> (concentration that scavenges the free radicals by 50%) of 12.5 µg/mL. FRAP value of 1.6 mmol TE/g extract.	[107]
<i>P. maculosa</i>	Persilben	AP	A 2.7 µM solution of persilben induced neutralization of DPPH radical by 40% (increase in content of persilben did not cause further reduction of DPPH).	[104]
	MeOH extract	AP	DPPH = 93.02% and FRAP = 7.3 mg/g. Standard BHT = 86.5%.	[108]
<i>P. senegalensis</i>	MeOH extract	AP	DPPH = 68.13% and FRAP = 6.2 mg/g. Standard BHT = 86.5%.	[108]
	Hydroalcoholic (50%) extract	L	IC <sub>50</sub> value for DPPH radical scavenging activities = 6.8 µg/mL. Positive control: L-ascorbic acid (IC <sub>50</sub> value 1.25 µg/mL).	[144]
	MeOH and EtOH extracts	R	IC <sub>50</sub> value of 49.20 µg/mL (MeOH) and 61.14 µg/mL (EtOH).	[39]
<i>P. bistorta</i>	MeOH extract and zinc oxide nanoparticles (ZnO-NPs)	R	MeOH extract showed higher DPPH and diammonium salt radical scavenging activity. Moreover, ZnO-NPs synthesized from root can inhibit the ABTS radicals, with IC <sub>50</sub> value of 40 µg/mL and a dose-dependent activity.	[40]
<i>P. amplexicaulis</i>	Nine known compounds and a previously undetermined one (5, 6-dihydropyranobenzopyrone), amplexicine and gallic acid (isolated from EtOH extract)	AP	IC <sub>50</sub> values of 10.2 µmol/L (5, 6-dihydropyranobenzopyrone), 12.2 µmol/L (amplexicine) and 14.4 µmol/L (gallic acid).	[30]
	Crude MeOH extract and their fractions (But, EtOH, EtOAc and aqueous)	Sh, L, Rh	All fractions and parts of plant displayed antioxidant activity (IC <sub>50</sub> between 1.03 and 58.2 µg/mL), but leaf crude MeOH and EtOAc fraction were the most effective for radical scavenging activity DPPH, with IC <sub>50</sub> = 1.03 and 3.1 µg/mL, respectively. Potent antioxidant activity respect to the standard L-ascorbic acid and Rutin:	[31]
<i>P. chinensis</i>	MeOH extract	WP, St, L	IC <sub>50</sub> from DPPH of 7.03–19.13 µg/mL. IC <sub>50</sub> from lipid peroxidation of 16.32–25.31 µg/mL. IC <sub>50</sub> from hydrogen peroxide method of 28.12–60.01 µg/mL.	[50]
<i>P. chinense</i> var. <i>chinense</i> and <i>P. chinense</i> var. <i>hispidum</i>	EtOAc fraction	AP	ORAC value of 0.965 µmol Trolox/mg.	[51]
	Aqueous extract		Exhibited moderate antioxidant activity, with IC <sub>50</sub> values from 180.87 to 255.69 µg/mL (from <i>chinense</i> variety) and 182.96 to 250.84 µg/mL (from <i>hispidum</i> variety). Gallic acid, chlorogenic acid, ellagic acid, quercitrin and brevifolin carboxylic acid isolated from these species could explain the antioxidant ability.	[52]
<i>P. minor</i>	MeOH extract	L	DPPH inhibition of 80.3% and FRAP value of 377 µMol Fe (II) g <sup>-1</sup> .	[110]

	Aqueous and EtOH extracts	L	DPPH radical scavenging of 81.88 and 89.5% and FRAP value of 849.33 and 11,220 mmol/g, from aqueous and EtOH extracts, respectively. [114] Authors have hypothesized that high levels of phenolic compounds (TPC 55.5–207 mg/g) may be the reason for the high antioxidant activity of this species. EtOAc extract showed at DPPH inhibition between 29.9 to 82.5%.
<i>P. bellardii</i>	EtOAc extract and isolated compounds	AP	Gallic acid and quercetin showed a minimum inhibition of 42.0–43.1% and maximum inhibition of 97.1–100%, values closely related to the reference's L-ascorbic acid and quercetin (45–99.6%). [167]
	EtOAc fraction	L	Strongly inhibited free radicals with an IC <sub>50</sub> value of 13.3 mg/mL. [76]
<i>P. hydropiper</i>	Flavonoids belonging to the quercetin family	L	TEAC values of 3.46–6.14. [77]
	Essential oil	L, F	Significantly DPPH free radical scavenging, ABTS and H <sub>2</sub> O <sub>2</sub> , with IC <sub>50</sub> values of 20, 180 and 45 µg/mL (for essential oil obtained from leaves) and 200, 60 and 50 µg/mL (for essential oil obtained from flowers), respectively. [78]
<i>P. paleaceum</i>	Crude extract, EtOAc and But fractions	Rh	A good capacity on DPPH, with SC <sub>50</sub> values of 16.72 µg/mL (crude extract), 10.64 µg/mL (EtOAc fraction) and 30.65 µg/mL (But fraction). [24] Gallic acid, caffeic acid derivatives and procyanidin are the main compounds that play an important role for the antioxidant activity of this plant.
<i>P. sagittata</i>	Gallic acid, methyl gallate, vanicoside A, quercetin, protocatechuic acid and vanicoside B (isolated from acetone extract)	St	Gallic acid showed the most potent DPPH scavenging activity (IC <sub>50</sub> 8.88 µM), followed by methyl gallate, vanicoside A, quercetin, protocatechuic acid and vanicoside B, with IC <sub>50</sub> values of 15.37, 26.82, 29.18, 32.38 and 35.06 µM, respectively. [142] Positive control: L-ascorbic acid (IC <sub>50</sub> = 30.49 µM).
	Acetone extract	AP	High O <sub>2</sub> radical dot scavenging (RSA of O <sub>2</sub> ) (IC <sub>50</sub> = 40.4 µg/mL) and a moderate total antioxidant capacity and anti-lipid peroxidation (IC <sub>50</sub> of 647 and 784 µg/mL). [176] IC <sub>50</sub> from DPPH of 26 µg/mL, IC <sub>50</sub> from FRAP of 48 µg/mL and IC <sub>50</sub> from CCA of 770 µg/mL.
<i>P. maritimum</i>	MeOH extract	L	Benzoic acid, phloroglucinol, phytol and linolenic acid were identified as possible compounds responsible for these bioactivities. [177]
	Crude extracts	AP	Good antioxidant scavenging effects on DPPH radical (7.71 µg/mL). [175] Positive controls: BHA, L-ascorbic acid and quercetin (DPPH 2.59–2.61 µg/mL).
<i>P. orientalis</i>	Taxifolin	L, S	Inhibitory effect on DPPH radical of 100% at a concentration of 7.5 µmol/L and the IC <sub>50</sub> value for taxifolin was 4.11 mmol/L. [125]

From the analysis of this table, it could be remarked that gallic, chlorogenic and ellagic acids may be the responsible for the strong antioxidant activities showed by *P. equisetiforme*, *P. lapathifolia*, *P. amplexicaulis*, *P. chinense* var. *chinense*, *P. chinense* var. *hispidum*, *P. bellardii*, *P. paleaceum* and *P. sagittata*, which in many experiments resulted to be even more potent than the standard drugs. On the other hand, some specific compounds such as persilben, amplexicine, quercitrin and brevifolin carboxylic acids, vanicoside A and B and taxifolin, were associated to the antioxidant response during the different bioassays carried out with *P. maculosa*, *P. amplexicaulis*, *P. chinense* var. *chinense*, *P. chinense* var. *hispidum*, *P. sagittata* and *P. orientalis*, respectively. Conspicuously, a study performed with MeOH extract and zinc oxide nanoparticles (ZnO-NPs) of *P. bistorta* inhibited the ABTS radicals with an IC<sub>50</sub> value of 40 µg/mL, and it was observed that the activity was dose-dependent. Moreover, it is well known that phenolic compounds are widespread in the plant kingdom, acting as antioxidants offering plant protections, so the *Persicaria* and *Polygonum* genera are not exceptional.

### 3.3. Analgesic and Anti-Inflammatory Activity

Many species of the *Persicaria* and *Polygonum* genera present anti-inflammatory properties, which are described in Table 4.

**Table 4.** Analgesic and anti-inflammatory activity of *Persicaria* and *Polygonum* species. Extracts, compounds, part of plant used, anti-inflammatory and analgesic activity with their references are shown. Extracts: DCM (dichloromethane); EtOH (ethanol); MeOH (methanol); Hex (hexane); EtOAc (ethyl acetate); But (butanol). Part used: L (leaves); St (stems); Rh (rhizomes); Sta (stalks); WP (whole plant); AP (aerial parts).

Species	Extract/Isolated Compounds	Part used	Summarized Bioactivity	Ref.
<i>P. chinensis</i>	MeOH extract	AP	At a concentration of 300 µg/mL, extract significantly inhibited regulation of nitric oxide (NO) at 72% in RAW264.7 cells and prostaglandin (PGE2) production was strongly suppressed up to 53%. These authors tested a murine HCl/EtOH-induced gastric ulcer model to evaluate the anti-inflammatory activity in vivo and found that the extract exhibited a significant anti-gastric activity, compared with the standard anti-ulcer ranitidine (40 mg/kg).	[53]
<i>P. chinensis</i> var. <i>hispidum</i>	Aqueous extract	AP	Ellagic acid and quercitrin inhibited the development of xylene-induced ear edema, with significant inhibition at a dose of 400 mg/kg.	[52]
<i>P. alpina</i>	MeOH extract	Rh	HRBC (Human Red Blood Cell) membrane stabilization method and percentage of inhibition protein denaturation method were used for tested in vitro anti-inflammatory activity of MeOH and aqueous extracts which showed a good anti-inflammatory ability, with a membrane stabilizing activity of 81.29% and an inhibition of protein denaturation of 72.70%, compared to the standard Indomethacin (95.56 and 88.26%, respectively).	[27]
<i>P. hydropiper</i>	MeOH extract	L	Extract blocked the production of NO, PGE2 and tumour necrosis factor on RAW264.7 cells and peritoneal macrophages.	[79]
	Aqueous extract	Sta	Extract attenuated the weight and length ratio of the colon, ameliorated the activity of MPO and the GSH content and regulated Cox-2, TNF-α and IL-1β levels in rats with TNBS-induced intestinal inflammation.	[80]
<i>P. pubescens</i>	α-Santalone (isolated from MeOH extract)	AP	The compound showed the most potent analgesic activity at a dose of 40 mg/kg b.w. (body weight), with an inhibition of acetic-acid-induced writhing response of 48.9%, compared to 62.2% for standard aminopyrine.	[191]
<i>P. lapathifolium</i> var. <i>lanatum</i>	Hex and MeOH extracts	WP	Anti-inflammatory activity: extracts inhibited carrageenan induced inflammation in rat paw at 41.09% (Hex) and 30.15% (MeOH), with a dose of 300 mg/kg b.w. Standard drug: phenylbutazone (42.15% inhibition; at a dose of 100 mg/kg b.w.). Analgesic activity: MeOH extract showed the highest inhibition of acetic acid-induced writhing reflex (62.29%; dose of 300 mg/kg b.w.), Standard drug: aminopyrine (69.94%).	[98]

<i>P. bellardi</i>	MeOH extract, their fractions (EtOAc and But) and isolated compounds (quercetin/its derivatives and myricetin/its derivatives)	AP	But extract was the most inhibitor of 5-lipoxygenase (5-LOX) (IC <sub>50</sub> 14.20 mg/mL), followed by EtOAc extract, MeOH extract, myricetin and quercetin (23.16, 24, 34.25 and 43.81 mg/mL, respectively). In addition, EtOAc and But extracts, myricetin and its glycosylated derivatives showed a significant inhibition of PGE <sub>2</sub> release (15.23–42.81%). At dose-dependent, extract exhibited a moderate inhibition percentage of haemolysis (50.37%). Positive control: hydrocortisone (inhibition of 86.56%).	[168]
<i>P. orientalis</i>	MeOH extract	L	Extract showed a good percentage of inhibition protein denaturation (79.22%). Positive control: diclofenac Na (inhibition of 86.85%). All doses tested (3.75, 5 and 7.5 g/kg) of the two extracts showed anti-inflammatory and analgesic activity, significantly inhibiting ear edema and significantly decreasing writhing in mice.	[126]
<i>P. stagnina</i>	EtOAc and ethyl ether extracts	St, L	The most potent analgesic activity was observed with the EtOAc extract (writhing inhibition of 50.3% at a dose of 400 mg/kg b.w.), while Hex extract showed the highest levels of anti-inflammatory activity (carrageenan-induced edema inhibition of 60.1% at a dose of 200 mg/kg b.w.), a much better effect than that of the conventional anti-inflammatory agent phenylbutazone (maximum 38.3% after 4 h). Extracts showed significant inhibition of NO production by LPS-stimulated RAW 264.7, at a concentration of 100 µg/mL.	[127]
<i>P. maritimum</i>	Hex, EtOAc and MeOH extracts	AP	$\beta$ -sitosterol, stigmasterol, 1-octacosanol and linolenic acid were identified as the possible compounds responsible for these bioactivities.	[145]
<i>P. maritimum</i>	MeOH and DCM extracts	L	Extract showed a potent ability to reduce NO production on LPS-stimulated RAW 264.7 macrophages (IC <sub>50</sub> of 22.0 µg/mL). Positive control: L-NAME (IC <sub>50</sub> of 27.6 µg/mL). Extract inhibited inflammatory reactions that cause instant irritation of the mouse ear, significantly inhibiting inflammatory mediators such as RAW264.7 cells (amurine macrophage cell line), production of NO, tumour necrosis factor TNF- $\alpha$ and IL-6 production in a dose-dependent manner.	[177]
<i>P. jucundum</i>	Acetone extract	AP	Flavonoids and sesquiterpene lactones may be responsible for the anti-inflammatory effect.	[176]
<i>P. jucundum</i>	EtOH extract	AP	At a dose of 30 µg/mL, EtOH extract inhibited the activities of lipoxygenase and cyclooxygenase-1, while the aqueous extract completely reduced paw edema induced by $\lambda$ -carrageenan at doses of 100 or 300 mg/kg b.w.	[173]
<i>P. minor</i>	Aqueous and EtOH extracts	AP	Exhibited anti-inflammatory activity against the carrageenan-induced pedal edema/Gastric intubation in vivo.	[115]
<i>P. punctata</i>	Decoction and EtOH-water extract	WP		[140]

	Tryptanthrin (isolated from EtOH extract)	L	Compound significantly inhibited the protein expression of iNOS and COX-2, suppressed the activation of p38 MAPK pathway and inhibited the TLR4 and MyD88 protein expression in LPS-stimulated BV2 microglial cells.	[151]
<i>P. tinctoria</i>	Polyphenolic fraction	L	Reduced NO synthesis in murine RAW264 macrophage cells stimulated with LPS, which showed a good analgesic activity.	[148]
	Extracts	St	Exhibited protective effects of DNA damage against oxidative stress and anti-inflammatory effects by its capacity for NO suppression in LPS-induced RAW264.7 cells.	[149]
<i>P. barbata</i>	Petroleum ether extract	AP	Inhibition of 39.3% paw edema after 2 h at a dose of 400 mg/kg b.w. (inhibitory ability slightly higher than that of the conventional anti-inflammatory agent phenylbutazone, 38.3% after 4 h).	[35]
<i>P. vivipara</i>	2-propanol extract		Exhibited anti-inflammatory activity against LPS-induced inflammation in RAW264.7 macrophages ( $IC_{50} = 270 \mu\text{g/mL}$ ) by inhibiting NO, prostaglandin, interleukin and tumour necrosis factor (TNF)- $\alpha$ release at similar levels as positive control.	[156]
<i>P. bistorta</i>	But extract		Extract exhibited analgesic effect, as it could reduce the writhing times of the mice induced by acetic acid and raised the threshold of pain induced by hot and electric stimulation.	[41]
<i>P. paleaceum</i>	Extracts		Extracts reduced malondialdehyde (MDA) content in inflamed paws, inhibited NO synthase and $\beta$ -NAG activities and significantly reduced the content of NO, IL-1 $\beta$ and TNF $\alpha$ in exudates.	[180]
<i>P. glabra</i>	Quercetin and quercetin glycosides	L	Compounds showed maximum effect at 90 min with latency time of 18.78 and 15.07 min, respectively (at doses of 200 mg/kg); the chemically induced writhing tests (for the evaluation of peripheral analgesic activity) showed maximum inhibition of writhing = 74.18% (quercetin) and 61.73% (quercetin glycosides) at a dose of 200 mg/kg in comparison with aspirin (dose of 50 mg/kg; inhibition of writhing = 78.41%).	[65]
<i>P. capitata</i>	Aqueous and EtOH extracts	WP	Significant inhibition of edema in animal models (76.19% and 71.13%, respectively) at a dose of 0.30 g/kg.	[48]
<i>P. perfoliatum</i>	Quercetin-3-O- $\beta$ -D-glucuronide		At a concentration of 8 mg/kg, compound suppressed ear edema induced by dimethyl benzene and peritoneal permeability induced by acetic acid in mice (45.96 and 40.10%, respectively), showing higher inhibition percentage respect to aspirin (24.62 and 34.38%, concentration 100 mg/kg).	[184]

From the analyses in Table 4, it could be observed that unlike the bioactivities described in the previous tables, for the evaluation of analgesic and anti-inflammatory effects, the whole extract of each species was evaluated more than the isolated compounds. Oppositely, more *in vivo* studies were informed for these bioactivities. The MeOH extract of *P. chinensis* exhibited anti-gastric activity compared with the standard ranitidine, but this effect was not correlated with the presence of any compound. The same fact occurred with the MeOH extract of *P. alpina*, which showed anti-inflammatory abilities compared to the standard Indomethacin, and with the MeOH extract of *P. lapathifolium* var. *lanatum*, which offered analgesic activity at the same level of the standard aminopyrine. These results could be explained by the synergistic effects of the compounds present in the MeOH extracts instead of the specific action of any compound present in them. On the other hand,  $\alpha$ -Santalone (isolated from the MeOH extract of *P. pubescens*) demonstrated to be the responsible for the potent analgesic activity of the extract, while flavonoids and sesquiterpene lactones may be responsible for the anti-inflammatory effect of *P. jucundum*. Finally, quercetin-3-O- $\beta$ -D-glucuronide isolated from *P. perfoliatum* suppressed ear edema and peritoneal permeability in mice showing higher inhibition percentage respect to aspirin.

### 3.4. Antinociceptive Activity

Antinociceptive activities of the *n*-Hex, EtOAc and MeOH extracts from *P. hydropiper* were tested by acetic-acid-induced writhing method in Swiss albino mice of either sex [81]. Ethyl acetate extract showed a moderate dose-dependent effect, with writhing inhibition of 54.95% at a dose of 500 mg/kg compared with the standard aminopyrine (inhibition of 73.62% at dose of 50 mg/kg). Additionally, Oany et al. [82] reported that crude EtOH extract of *P. hydropiper* leaves exhibited a good antinociceptive activity compared to the standard drug aspirin, displaying a reduction in the abdominal writhing's number of 41.02% and 69.23%, respectively, at a dose of 400 mg/kg b.w. Petroleum ether and chloroform (Chl) extracts of *P. barbata* showed antinociceptive properties, with an inhibition of writhing response of 46.8% and 44.8%, respectively (at a dose of 400 mg/kg b.w.), compared to 62.2% for the positive control aminopyrine [35]. The authors concluded that the compounds responsible for this activity were of apolar nature, possibly sterols or terpenoids. With respect to *P. acuminata*, the EtOH extracts of the leaves and stems exhibited antinociceptive activities with a percentage of writhing inhibition of 53.57% and 50%, respectively, at a dose of 500 mg/kg, similar values compared to the standard diclofenac (57.7% at a dose of 25 mg/kg) [9]. The antinociceptive activity of the crude MeOH extract of *P. verticillatum* rhizomes was tested by Khan et al. [193]. They found that at dose-dependency and through an opioid system, the extract could significantly reduce (72%) the number of writhes induced by acetic acid. The extract contained a high content of saponins and alkaloids.

### 3.5. Anticancer, Antitumoral and Cytotoxic Activity

Anticancer and antitumoral activities displayed by *Persicaria* and *Polygonum* species were studied in over the world. Many of these species have inhibitory abilities against different cancerogenic cell lines (Table 5): Jurkat, HL60, THP-1, CCRF-CEM, K562, U-937, K562 and P338 (human leukaemia); HepG2, Huh-7, SMMC-7721, HCCLM3, Hep3B (hepatocellular carcinoma); CaCo-2, HCT116, HT-29, CT-26, RKO, Colo320 and SW620 (colon cancer); MCF-7, HTB-26, MDA-MB-231-pcDNA3, MDA-MB-435 and Bcap-37 (breast cancer); SiHa, HeLa and PC-3 (cervical carcinoma); PC3 and PC3 (prostate carcinoma); LL2, A549, 95D and NCI H460 (lung carcinoma); SNU-601 and SGC-7901 (gastric cancer); PANC-1 (pancreatic adenocarcinoma); H22 (murine H22 ascitic hepatoma); EAC (Ehrlich ascites carcinoma); EBV-EA (Epstein-Barr virus early antigen); J82 (bladder transitional carcinoma); HEK 293 (renal cancer); NU-1066 (laryngeal cancer); OVCAR-3 (ovarian cancer); U87MG and BT-325 (glioblastoma); CAL 27 (oral adenosquamous carcinoma); Smur180 (sarcoma); and NIH3T3 (fibroblast).

**Table 5.** Anticancer and antitumoral activities of *Persicaria* and *Polygonum* species. Extracts, compounds, part of plant used, cell line and references are shown. Extracts: EtOH (ethanol); MeOH (methanol); Chl (chloroform); Hex (hexane); EtOAc (ethyl acetate); But (butanol). Part used: L (leaves); F (flowers); Fr (fruits); R (roots); Sp (sprouts); S (seeds); St (stems); Rz (rhizomes); T (tubers); B (barks); WP (whole plant); AP (aerial parts).

Species	Extract/Isolated Compounds	Part use	Cell Line	Summarized Bioactivity	Ref.
<i>P. amphibia</i>	Flavonoid glucosides: quercetin 3- <i>O</i> - $\beta$ -D-glucopyranoside and quercetin-3- <i>O</i> - $\alpha$ -rhamnosyl-(1-2)- $\beta$ -glucoside (isolated from butanoic fraction)	AP	Jurkat	EC <sub>50</sub> 1.2 and 0.12 $\mu$ M.	[29]
			HL60	EC <sub>50</sub> 0.98 and 0.01 $\mu$ M.	
<i>P. amplexicaulis</i>	Flavonoids (isolated from EtOH extract)  Amplexicaule A (isolated from EtOH extract)	Rz	HepG2, Huh-At 7, H22, SMMC-7721	At a dose-time dependence: reduced cell viability, induced cell apoptosis and increased expression of SHP-1 (tyrosine phosphatase catalysing STAT3 dephosphorylation protein).	[32]
			MCF-7 and MDA-MB-435	At a concentration of 150 mg/kg: suppressed tumour mass in 0.6–0.7 g (respect to positive control Capecitabine 10 mg/kg, 0.5–0.6 g), induced apoptosis in cancer cells, increased caspase-3, -8, -9 and PARP (enzymes and proteins that catalyse apoptosis) levels and suppressed MCL-1 and BCL-2 expression.	[33]
<i>P. barbata</i>	Methyl (2 <i>S</i> ,3 <i>S</i> )-2-(3,4-dimethoxyphenyl)-4-(( <i>E</i> )-3-ethoxy-3-oxoprop-1-en-1-yl)-7-methoxy-2,3 dihydrobenzo-furan-3-carboxylate (1) and ( <i>E</i> )-3-((2 <i>S</i> ,3 <i>S</i> )-2-(3,4-dimethoxyphenyl)-7-methoxy-3-(methoxy carbonyl)-2,3-dihydro-benzofuran-4-yl) acrylic acid (2) (isolated from EtOAc fraction)  Petroleum ether extract	WP	CAL-27 and NCI H460	IC <sub>50</sub> of 48.52 and 53.24 $\mu$ M (for 1) and 86.95 and 93.34 $\mu$ M (for 2). Standard drugs: 5-Fluorouracil, IC <sub>50</sub> 97.76 $\mu$ M (for CAL-27) and Cisplatin, IC <sub>50</sub> 19 $\mu$ M (for NCI H460). Compound 1 induced apoptosis in CAL-27 cell line after 24 to 48 h treatment.	[36]
			Potato disc assay	Moderate activity, with inhibition of tumour growth of 57.1% at a concentration of 400 $\mu$ g/disc and IC <sub>50</sub> value of 290 $\mu$ g/disc. Positive control: vincristine sulphate 3.125 $\mu$ g/disc, 100% inhibition tumour).	[37]
<i>P. bistorta</i>	Gallic acid, protocatechuic acid, syringic acid, catechol, syringol, 4-methyl catechol, myristic acid (isolated from MeOH-water extracts)	Rz	HCCLM3	Cell viability < 30% (at a dose of 200 $\mu$ g/mL) and GI <sub>50</sub> values between 86.5 to 126.8 $\mu$ g/mL.	[42]

	Aqueous extract	WP	Hep3B	Inhibited autophagosome and proteasome activity, resulting in restriction of cell motility and apoptosis induction in Hep3B cells.	[43]
	Zinc oxide nanoparticles (ZnO-NPs) (synthesized using <i>P. bistorta</i> extract)	R	MCF-7	Effective dose-dependent activity, with a percentage of cells viability < 10% (at a concentration of 125 µg/mL).	[40]
	<i>n</i> -Hex and Chl fractions and its sub-fractions	Rz	P338, HepG2, J82, HL60, MCF7, LL2	Fractions were effective against all cell lines, but showed the highest cytotoxicity against P338, HL60 and LL2 (IC <sub>50</sub> < 10–62.4 µg/mL). Sub-fractions showed cytotoxicity against all cell lines, with IC <sub>50</sub> values between < 10–91.2 µg/mL.	[44]
<i>P. chinensis</i>	Corilagin and ellagic acid	WP	SiHa	Effective dose-dependent activity, with cell inhibition of 59 and 81% (at a concentration of 100 µM) and IC <sub>50</sub> values of 21.5 and 28.7 µM, respectively.	[54]
<i>P. decipiens</i>	Phenolic acids and flavonoids (isolated from MeOH extract)	L, S	CaCo-2, PC3	IC <sub>50</sub> = 0.5–1.1 µg/mL.	[58]
<i>P. glabra</i>	(-)-pinocembrin (isolated from MeOH extract)	AP	THP-1, A549, PANC-1, HeLa, MCF7	IC <sub>50</sub> values between 1.88 to 11.00 mg/mL.	[63]
	MeOH extract and its fractions (saponins, Chl and EtOAc)	WP	Potato disc anti-tumour assay	Assays were performed on <i>Agrobacterium tumefaciens</i> containing tumour-inducing plasmid. Tumour inhibitions between 80 to 90% (at a dose of 1000 µg/mL) and IC <sub>50</sub> values between 18.39 and 342.53 µg/mL.	[83]
	β-sitosterol and stigmasterol (isolated from Chl and EtOAc fractions)	AP	NIH/3T3, HeLa, MCF-7	Effective dose-dependent activity, with percentage of cytotoxicity between 67.05 and 87.5% and IC <sub>50</sub> of 170–425 µg/mL (for β-sitosterol) and 60–170 µg/mL (for stigmasterol).	[84]
<i>P. hydropiper</i>		AP		Cell growth inhibition of 84.54% (at a concentration of 50 mg/kg/day) and improved at a 68% the survival of mice.	[85]
	MeOH extract	L	EAC	Significantly decreased tumour volume, packed cell volume and viable tumour cell and increased non-viable tumour cell. At a dose of 100 mg/kg, the median survival time (MST) was 37.21, respect to reference bleomycin (46.60).	[86]
<i>P. lapathifolia</i>	Pinostrobin (isolated from petroleum ether extract)	AP	Jurkat and HL60	Dose-dependent effects, with a percentage of apoptotic cells > 70% (at a dose of 1 µM) and a percentage of necrotic cells > 80% (at a dose of 10 µM).	[99]

	Vanicoside B and lapathoside A (extracted with MeOH)		EBV-EA	Carcinogenesis was induced by 7,12-dimethylbenz[a]anthracene (DMBA, as initiator), ( <i>E</i> )-methyl-2-[( <i>E</i> )-hydroxyamino]-5-nitro-6-methoxy-3-hexenamamide (NOR-1) and 12- <i>O</i> -tetradecanoylphorbol-13-acetate (TPA) as a promoter. The number of papillomas per mouse were reduced at 3.4 and 2.6 after 15 weeks compared to the control (DMBA + TPA, 9.1 papillomas per mouse). In the NOR-1 + TPA treatment, vanicoside B reduced the number of papillomas from 7.2 (control) to 3.	[100]
<i>P. limbata</i>	Cardamomin and 2',4'-dihydroxy-3',6'-dimethoxychalcone (isolated from MeOH extract)	AP	MCF-7, THP-1, PC-3, HeLa	Proliferation inhibition > 50%. The best activity was observed against THP-1 cell line, with IC <sub>50</sub> < 4 µg/mL.	[102]
	Flavonoids	L	CCRF-CEM, MDA-MB-231-pcDNA3, HCT116, U87MG, HepG2	The flavonoid 4-hydroxy-2,6-dimethoxychalcone, showed the best activity against all cell lines, with IC <sub>50</sub> values of 9.37, 19.58, 6.8, 35.25 and 58.63 µM. The flavonoids cardamomin and 2,4-dihydroxy-3,6-dimethoxychalcone were effective against CCRF-CEM cell line (IC <sub>50</sub> of 8.59 and 10.67 µM, respectively).	[103]
	EtOAc (100%), aqueous-EtOH (50%), MeOH (100%), EtOH (70%) extracts and aqueous solution (100%)	L	HT-29, HCT-116, CT-26	EtOAc extract (100%) showed the highest cytotoxic effect against HCT-116 and CT-26 and Aq-EtOH extract (50%) was the most effective against HT-29 cell line (IC <sub>50</sub> of 7.0 and 24.0 µg/mL, respectively). Aq solution (100%), MeOH (100%) and EtOH (70%) extracts showed moderate activity, with IC <sub>50</sub> of 34.0–78.0 µg/mL (HT-29), 13.0–33.0 µg/mL (HCT-116) and 20.0–29.0 µg/mL (CT-25), respectively. IC <sub>50</sub> values for standard agent Doxorubicin: 0.63 (HT-29), 0.46 (HCT-116) and 0.14 µg/mL (CT-26).	[116]
<i>P. orientalis</i>	Polygonumins A (extracted with MeOH)	St	K562, MCF7, HCT116	Good activity compared to the positive control Doxorubicine (IC <sub>50</sub> = 2.25–3.24 and 0.52–2.97 µg/mL, respectively).	[117]
	EtOAc and <i>n</i> -But extracts	F	HeLa, SMMC-7721	Proliferation inhibition about 40–60% at doses between 50 to 450 µg/mL and after 48 h.	[128]
	EtOAc and Acetone extracts	Fr	95D	IC <sub>50</sub> values = 199.1 mg/L (for EtOAc extract) and 261.2 mg/L (for acetone extract).	[129]
<i>P. senegalensis</i>	Phenolic acids and flavonoids (isolated from MeOH)	L, S	CaCo-2, PC3	IC <sub>50</sub> 1.5–3.5 µg/mL.	[58]

<i>P. stagnina</i>	EtOAc and <i>n</i> -Hex extracts	AP	Potato assay disc	Moderate activity, with inhibition of tumour growth between 50 to 78.6% at a concentration of 200–400 µg/disc and IC <sub>50</sub> values = 180 and 200 µg/disc (for EtOAc and <i>n</i> -Hex extracts, respectively). Positive control: vincristine sulphate 3.125 µg/disc, 100% inhibition tumour).	[37]
<i>P. tinctoria</i>	MeOH and EtOH extracts	F, L, St, S	HEK 293, HCT-116, HeLa, Hep3B, MCF-7, SNU-1066, SNU-601	EtOH extract from flowers was the most effective against all cell lines, except for MCF-7 and SNU-601, with survival rate of cancer cells = 5.10–25.27%. Leaves' MeOH extract showed a good activity against HCT-116, HeLa, Hep3B and SNU-1066 (survival cancer cells = 6.89–26.47%); stems' MeOH extract was effective against Hep3B and SNU-1066 (survival cancer cells = 26.94–29.28%); and seeds' MeOH extract showed a good activity against HEK 293, HeLa, Hep3B and SNU-601 (survival cancer cells = 22.89–29.85%).	[150]
	Tryptanthrin (isolated from EtOAc extract)	L	U-937, HL-60	Compound showed 100% of cytotoxic effects on both cell lines (at a concentration of 6.3 mg/mL) and inhibited DNA synthesis at dose-dependency.	[152]
<i>P. aviculare</i>	MeOH extract	AP	MCF-7 Hela-S	The extract induced cytotoxicity in MCF-7 cell line, with a 99% of cell death at the concentration of 400 ng/µl after 24 hrs. Showed cytotoxic effect at IC <sub>50</sub> values between 0.27 and 0.41 mg/mL and caused complete apoptosis at 24 h of treatment.	[161] [162]
<i>P. bellardii</i>	MeOH, EtOAc and <i>n</i> -But extracts and isolated compounds (gallic acid, quercetin/its derivatives and myricetin/its derivatives)	AP	HeLa, MCF-7, HepG-2	At a concentration between 130 and 170 µg/mL, all extracts and fractions inhibited cell viability at 20% of all cell lines tested. <i>n</i> -But extract was the most potent against three cell lines (IC <sub>50</sub> = 15.26, 50.66 and 30.09 µg/mL, respectively); MeOH and EtOAc extracts showed a good activity against HeLa cells (IC <sub>50</sub> = 48.6 and 44.14 µg/mL, respectively); and quercetin derivatives and myricetin and its derivatives were effective against HepG-2 (IC <sub>50</sub> between 41.03 to 70.77 µg/mL).	[168]
<i>P. cognatum</i>	EtOH extract	L	MDA-MB-231	MDA-MB-231 cell viability < 50% and IC <sub>50</sub> = 0.053 mg/mL.	[169]
<i>P. perfoliatum</i>	8-oxo-pinoresinol (isolated from MeOH extract)	T	Bcap-37, SMMC-7721,	IC <sub>50</sub> values from 8.32 to 30.1 µg/mL (positive control Mitomycin, IC <sub>50</sub> values from 1.75 to 6.24 µg/mL).	[20]

			K562, RKO, PC3 Smur180, SGC-7901, AP Colo320, PC-3, HL60	Smur180 inhibition cells = 58.46% (at a dose of 200 mg/kg) and inhibited the growth and proliferation of other cell lines, with IC <sub>50</sub> < 50 µg/mL.	[185]
		EtOAc extract	PANC-1, PC-3, SGC-7901, AP BT-325, HepG2, A549, Hela	Inhibition cell lines = 70.1–90% and IC <sub>50</sub> values between 20.6 to 40.7 µg/mL. Furthermore, the extract arrested cells at G2 phase, increased the proliferation of T and B lymphocytes, promoted the activities of NK and cytotoxic T lymphocytes (CTLs) and induced cell apoptosis.	[186]
<i>P. thunbergii</i>		Isorhamnetin	AP NIH3T3, SW620	The compound decreased the percentage of cell proliferation with IC <sub>50</sub> values of 4.1 µg/mL (for NIH3T3) and 22.4 µg/mL (for SW620).	[192]
		MeOH crude extract	B Brine shrimp lethality test	The extract showed a good cytotoxic ability respect to the standard Vincristine sulphate (IC <sub>50</sub> = 6.34 and 0.825 µg/mL, respectively).	[194]
<i>P. viscosum</i>		Quercetin 3-O-(6-feruloyl)-β-D-galactopyranoside	WP OVCAR-3	IC <sub>50</sub> = 13.33 µg/mL. Authors considered that this effect could be attributed to the presence of moieties such as quercetin, galactosyl and principally feruloyl in this compound.	[195]

According to this information, flavonoids glucosides provide the antitumoral activities to *P. amphibia*, *P. amplexicaulis*, *P. decipiens*, *P. glabra*, *P. limbata* and *P. senegalensis* with IC<sub>50</sub> values varying from 0.01 to 100 µM depending on the cellular type under study. Other compounds such as gallic acid, ellagic acid and steroids seem to be responsible for the antitumoral activities of *P. bistorta*, *P. chinensis*, *P. hydropiper* and *P. bellardii* but show lower IC<sub>50</sub> than flavonoids. Some specific compounds such as amplexicaule A, vanicoside B, lapathoside A and polygonumin A isolated from *P. amplexicaulis*, *P. lapathifolia* and *P. minor*, respectively, showed promising anticancer activities with IC<sub>50</sub> values comparable to standard drugs.

### 3.6. Antiviral Activity

Some *Persicaria* and *Polygonum* species have shown antiviral abilities against different types of viruses (Table 6): HHV-1 and HSV-1 (human herpes virus 1); EBV-EA (Epstein-Barr virus); H1N1 A/PR/8/34, H1N2 A/HK/8/68 and B/Lee/40 (influenza A y B); HIV-1<sub>VB59</sub> and HIV-1UG070 (immunodeficiency human virus); VACV-WR (vaccinia virus strain Western Reserve); VV (vaccinia virus); DEN-2 (dengue virus 2); VSV (vesicular stomatitis virus); RSV (respiratory syncytial virus); MCMV (cytomegaly virus); and HBV (hepatitis B virus). It could be highlighting the inhibition of HIV-1 protease (56%) of polygonumins A isolated from *P. minor* and the HIV-1 reverse transcriptase inhibitory activity of viscozulone isolated from *P. viscosum*.

**Table 6.** Antiviral activities of *Persicaria* and *Polygonum* species. Extracts, compounds, part of the plant used, target virus and references are shown. Extracts: EtOH (ethanol); MeOH (methanol); Hex (hexane); EtOAc (ethyl acetate); But (butanol). Part used: L (leaves); F (flowers); Fr (fruits); R (roots); Sp (sprouts); S (seeds); St (stems); Rz (rhizomes); T (tubers); B (barks); WP (whole plant); AP (aerial parts).

Species	Extract/Isolated Compounds	Part use	Pathogen	Summarized Bioactivity	Ref.
<i>P. chinensis</i>	MeOH, But and EtOAc extracts. Ellagic acid, methyl gallate and caffeic acid (isolated from EtOAc extract)	WP	H3N2-HK, H1N1-PR8, Lee	EC <sub>50</sub> = 18.3–38.4 µg/mL (for H3N2) and 45.9–70.1 µg/mL (for H1N1 and Lee). Significantly inhibited viral replication (EC <sub>50</sub> = 14.7–81.1 µg/mL) by suppressing virus replication in cells.	[55]
<i>P. ferruginea</i>	EtOH extract	AP	HHV-1, DEN-2, VACV-WR	EC <sub>50</sub> values = 21.1, 24.6 and 34.2 µg/mL, respectively.	[60]
<i>P. glabra</i>	2-methoxy-5-oxo-2,5-dihydrofuran-3-yl (2E)-(-)-3-phenylprop-2-enoate (isolated from MeOH extract)	AP	HIV-1VB59, HIV-1UG070	IC <sub>50</sub> values = 15.68–22.43 mg/mL.	[63]
<i>P. lapathifolia</i>	Lapathoside A and D, vanicoside B and hydro-piperoside (isolated from MeOH extract)	AP	EBV-EA	Inhibition of activation on EBV-EA higher than 85, 60 and 30%.	[100]
<i>P. minor</i>	Polygonumins A	St	HIV-1	Inhibition of HIV-1 protease = 56.51% (positive control Pepstatin A, inhibition = 81.48%). The authors considered that the phenyl propanoid glycoside moiety present in Polygonumins A, may be the responsible for the anti-HIV protease activity.	[117]
<i>P. punctata</i>	EtOH extract	L	HSV-1, VSV	MIC values = 0.01 and 0.02 mg/mL (for HSV-1 and VSV, respectively).	[118]
	Aqueous extract	AP	HSV-1, RSV	ED <sub>50</sub> values = 169.7 and 120 µg/mL (for HSV-1 and RSV, respectively)	[141]
	MeOH	WP	HSV-1	At a MIC of 20 µg/mL, the extract caused the complete virus inactivation.	[138]
<i>P. tinctoria</i>	MeOH and EtOAc extracts	WP	H1N1-PR8, HSV-1, VV, MCMV, VSV	IC <sub>50</sub> values between 1.25 and 16.6 µg/mL (for MeOH extract) and 0.63 to 50 µg/mL (for EtOAc extract).	[153]
<i>P. perfoliatum</i>	Aqueous extract	L	HIV-1	Inhibition of HIV-1 (IIIB) infection at EC <sub>50</sub> value of 0.5 µg/mL.	[154]
	Quercetin-3-O-β-D-glucuronide	WP	influenza A virus	Inhibition of 27.94%, at a concentration of 6 mg/kg (reference drug Ribavirin, inhibition of 23.97%).	[184]

	Flavonoids (extracted with MeOH and <i>n</i> -Hex)	L	HSV-1	At a concentration of 62.5 µg/mL, flavonoids reduced more than 80% the number of plaques in infected cultures, in the same way as the control ACV. At a dose of 30 mg/kg/day, the survival and mean survival time (MST) of mice induced with encephalitis HSV-1 were 80% and 19.0 days, respectively, higher values compared with the control ACV (survival of 70% and MST = 18.5 days).	[187]
	Gallic acyl groups (isolated from <i>n</i> -But and water extracts)	WP	HBV	Inhibition up to 74% secretion of antigen (HBeAg).	[188]
<i>P. viscosum</i>	Quercetin 3- <i>O</i> -(6-feruloyl)-β- <i>D</i> -galactopyranoside and viscoazulone	WP	HIV-1	Reverse transcriptase inhibitory activity with IC <sub>50</sub> values of 33.13 and 25.61 mg/mL (for quercetin and viscoazulone, respectively).	[195]

### 3.7. Antiparasitic Activity

Extracts and isolated compounds from *Persicaria* and *Polygonum* species were tested against different parasites that cause human diseases such as *Plasmodium falciparum* (causal agent of malaria disease); *Trypanosoma cruzi* and *T. brucei* (causal agents of Chagas disease); *Leishmania amazonensis* and *L. tropica* (causal agents of leishmaniasis disease); and *Pheretima posthuma* and other parasitic earthworms (which cause helminth infections). Results of these investigations are show in Table 7.

**Table 7.** Antiparasitic activities of *Persicaria* and *Polygonum* species. Extracts, compounds, part of the plant used, antiparasitic activity and references are shown. Extracts: EtOH (ethanol); MeOH (methanol); Hex (hexane); EtOAc (ethyl acetate). Part used: L (leaves); F (flowers); Fr (fruits); R (roots); Sp (sprouts); S (seeds); St (stems); Rz (rhizomes); T (tubers); B (barks); WP (whole plant); AP (aerial parts).

Species	Extract/Isolated Compounds	Part use	Pathogen	Summarized Bioactivity	Ref.
			Anti-malarial		
<i>P. acuminata</i>	MeOH extract	L	<i>P. falciparum</i>	IC <sub>50</sub> = 8 µg/mL	[26]
<i>P. glabra</i>	EtOH extract	L	<i>Plasmodium</i> sp.	IC <sub>50</sub> = 6.6 µg/mL	[66]
			Anti-trypanocide		
<i>P. decipiens</i>	2,4-dimethoxy-6-hydroxychalcone (chalcone 1) and 2,5-dimethoxy-4,6-dihydroxychalcone (chalcone 2)	AP	<i>T. brucei</i> <i>T. congolense</i>	EC <sub>50</sub> = 1.8–8.8 µM (for chalcone 1) and 13.9–34 µM (for chalcone 2). Positive controls: Diminazene (EC <sub>50</sub> = 0.15–1.43 µM) and Pentamidine (EC <sub>50</sub> = 0.0034–0.72 µM). IC <sub>50</sub> values = 8.6–10.5 µg/mL (for Hex and dichloromethane extracts) and 50–90 µg/mL (for EtOAc extract).	[59]
	<i>n</i> -Hex, DCM and EtOAc extracts			Positive controls: Pentamidine (IC <sub>50</sub> = 6.4 and 2.2 µg/mL) and Benznidazole (IC <sub>50</sub> = 34.7 and 54.1 µg/mL).	[61]
<i>P. ferruginea</i>	2'-hydroxy-4',6'-dimethoxychalcone, flavokawin B (1), 2',6'-dihydroxy-3',4'-dimethoxychalcone, pashanone (2) and (2',4'- dihydroxy-6'-methoxychalcone, cardamomin or alpinetin chalcone (3) (isolated from Hex extract)	AP	<i>T. cruzi</i> <i>T. brucei</i>	Compounds 1, 2 and 3 were effective against <i>T. cruzi</i> (IC <sub>50</sub> between 9.5 and 32.3 µM). Against <i>T. brucei</i> , only compound 1 was active, with IC <sub>50</sub> = 6.2 (for strain 427) and 4.8 µM (for strain 29–13).	
	MeOH extract		<i>T. cruzi</i>	IC <sub>50</sub> = 37 µg/mL.	[26]
<i>P. hydropiper</i>	Cardamomin, vanicoside F, ketopinoresinol, isorhamnetin and pinosylvin (isolated from DCM soluble portion)	WP	<i>T. brucei</i>	IC <sub>50</sub> = 0.49–0.8 µg/mL (for cardamomin) and 0.49–7.77 µg/mL (for other compounds). Positive control: α-difluoromethylornithine (DFMO), IC <sub>50</sub> = 3.02 µg/mL.	[87]
			Anti-leishmanial		
<i>P. glabra</i>	Aqueous extract	WP	<i>Leishmania tropica</i>	At a minimal concentration tested (0.05 µg/mL), extract showed a 4.23% parasite mortality, while using a concentration of 50 µg/mL, the mortality raised up to 68.1%.	[67]
<i>P. hydropiperoides</i>	MeOH extract	F	<i>Leishmania amazonensis</i>	IC <sub>50</sub> = 73 µg/mL.	[95]
			Anthelmintic		

<i>P. alpina</i>	MeOH extract	Rz Earthworms	Earthworms' death times = 71.4 min at a concentration of 100 mg/mL (standard Albendazole, death time = 56.6 min).	[27]
<i>P. chinensis</i>	Aqueous and MeOH extracts	L <i>Pheretima posthuma</i>	Inhibition and death at 5.83 and 16.5 min (for Aqueous extract), respectively, and at 9.25 and 19.67 min (for MeOH extract), respectively, at a dose = 100 mg/mL.	[56]
<i>P. hydropiper</i>	MeOH extract	AP <i>P. posthuma</i>	Standard drug: Albendazole (inhibition and death at 5.33 and 6.92 min). At the extract concentration of 50 mg/mL, the time of paralysis and death of earthworms were 12.44 and 18.19 min.	[85]
	Saponins, Chl and <i>n</i> -But (fractionated from MeOH extract)	WP <i>P. posthuma</i>	Positive control: Piperazine citrate, 10 mg/mL (time of paralysis and death = 24 and 38 min, respectively). Paralysis time between 8 and 11 min and death time between 50 and 66.33 min at a concentration of 10 mg/mL.	[88]
<i>P. lapathifolia</i>	MeOH extract	St <i>P. posthuma</i>	At a concentration of 60 mg/mL, the times taken for paralysis and death were 9 and 23.66 min, respectively.	[101]
<i>P. muricatum</i>	EtOH extract	L Earthworms	Standard drug: Piperazine citrate, 10 mg/mL (time taken for paralysis and death = 9.33 and 36 min, respectively).	[179]
			At a concentration of 100 mg/mL, the paralysis and death times were 35 and 43 min, respectively. Standard drug: Albendazole (paralysis and death time = 20 and 30 min, respectively).	
<i>P. viscosum</i>	MeOH extract		Phytochemical analysis identified various constituents such as alkaloids, carbohydrates, glycosides, phytosterols, phenolic compounds, tannins, saponins, proteins and amino acids.	[194]

Few antiparasitic studies have been conducted using these two plant genera, and most of them were carried out with extracts. It could be remarked the anti-trypanocidal activities of two chalcones obtained from *P. decipiens*, which showed IC<sub>50</sub> values compared to the standard drugs diminazene and pentamidine against *T. brucei* and *T. congolense*. In addition, saponins from the butanolic extract of *P. hydropiper* displayed anti-leishmanial actions similar to the standard drug piperazine citrate.

### 3.8. Anti-Diabetic Activity

The anti-diabetic ability of EtOH extract from *P. aviculare* leaves was tested by Cai et al. [163]. The results showed that the extract is a potent inhibitor of  $\alpha$ -glucosidase levels, higher than the standard drug acarbose, with IC<sub>50</sub> values of 21.42 and 176.79  $\mu$ g/mL, respectively. Additionally, the compounds myricitrin, quercetin, polydatin and isoquercitrin (present in the phenolic portion of *P. aviculare* extracts) were primarily responsible for this activity, with IC<sub>50</sub> values of 8.65, 15.17, 35.15 and 148  $\mu$ g/mL, respectively [163]. Ethanol extract of *P. pulchra* leaves showed a good  $\alpha$ -glucosidase (involved in Type 2 diabetes mellitus) inhibition with IC<sub>50</sub> value of 22.67 mg/mL, respect to the positive control acarbose (7.77 g/mL) [134]. In addition, Oany et al. [82] tested antihyperglycemic activity of crude EtOH extracts from leaves and stems of *P. hydropiper* and found that for all the doses tested (50, 100, 200 and 400 mg/kg b.w.), leaves's extract showed higher effectiveness than standard glibenclamide, diminishing blood glucose levels between 48.8 and 58.2% for the extract and 42.1% for glibenclamide (at dose 10 mg/kg b.w.). Furthermore, the extracts of the leaves and stems (principally *n*-Hex, Chl and MeOH extracts) displayed  $\alpha$ -amylase inhibitory activities, with IC<sub>50</sub> values between 1.03 and 3.517 mg/mL [75]. This enzyme can increase the blood sugar level, as hydrolyses (1,4)- $\alpha$ -D-glycosidic linkages in polysaccharides contain three or more (1,4)- $\alpha$ -linked D-glucose units. Another study demonstrated that the EtOH extract of *P. cognatum* strongly inhibited  $\alpha$ -amylase activity (86.6%) and moderately inhibited  $\alpha$ -glucosidase activity (41.05%) with respect to the standard acarbose (58.4% and 57.56%, respectively) [169].

Kubinova et al. [97] tested the anti-diabetic activity of isolated compounds from the MeOH extract of *P. lapathifolia* aerial parts by the inhibition of AChE, BuChE and  $\alpha/\beta$ -glucosidase. At a dose of 100  $\mu$ M, kaempferol showed the highest inhibition of AChE (60.4%) and BuChE (74.5%), compared to the standard galantamine (95.7 and 47.9%, respectively), while gallic acid was only effective on AChE (inhibition of 51.2%). With respect to the  $\alpha$ -glucosidase inhibitory assay, all isolated flavonoids showed potent enzyme inhibitory activity (72.6–97.2%) and eightfold higher activity than standard acarbose (11.4%), while for the  $\beta$ -glucosidase inhibitory assay, only quercitrin and isoquercitrin inhibited the enzyme with higher efficacy than the standard quercetin (23.6, 23.4 and 16.7%, respectively). Rodrigues et al. [177] tested the anti-diabetic ability of the leaf and root extracts of *P. maritimum* on  $\alpha$ -amylase, baker's yeast ( $\alpha$ -glucosidase) and rat's  $\alpha$ -glucosidases. Both the MeOH and DCM extracts showed IC<sub>50</sub> values lower than the positive control (acarbose) on  $\alpha$ -glucosidase, but the MeOH extract had the highest capacity to inhibit the baker's yeast  $\alpha$ -glucosidase, with an IC<sub>50</sub> value significantly lower than that of acarbose (IC<sub>50</sub> value of 19 and 29  $\mu$ g/mL for roots and leaves; 3144  $\mu$ g/mL for acarbose).

The anti-hyperglycaemic effect of aqueous extract of *P. orientalis* flowers was tested on streptozotocin (STZ)-induced diabetic mice [130]. At a dose of 200 mg/kg, it was observed the most potent results: the extract significantly decreased blood glucose levels (144 mg/mL, 21 days) and serum cholesterol (53.2 mg/dL) compared to the control (210.06 and 82.4 mg/dL, respectively) and increased glycogen content in liver (30.7 mg/g for treatment and 11.86 mg/g for diabetic control). The hydroalcoholic (50%) extract of *P. senegalensis* leaves exhibited a potent anti-diabetic activity, inhibiting 100% of the  $\alpha$ -glucosidase activity at a concentration of 10  $\mu$ g/mL [144].

### 3.9. Antipyretic Activity

The antipyretic activity of the aqueous root extract of *P. bistorta* was studied in albino rats with Brewer's-yeast-induced fever [45]. At a concentration of 100 mg/kg and after 4 h of treatment, the extract exhibited similar pyrexia activities to that of the standard drug acetaminophen, with a decrease in the rectal temperature of 0.8 °C. Another study revealed that the MeOH root extract of *P. glabra* exhibited a significant dose-dependent antipyretic activity on albino rats submitted to the Brewer's-yeast-induced hyperpyrexia method [68]. Alkaloids, flavonoids and phenolic compounds could be responsible for this activity. Finally, Akhter et al. [89] tested the antipyretic activity of MeOH, EtOH, Chl, petroleum ether and *n*-Hex extracts of *P. hydropiper* leaves on Albino Swiss mice of both sexes. At doses of 200 and 400 mg/kg b.w., the MeOH, EtOH and Chl extracts showed a good activity compared to the standard drug paracetamol (50 mg/kg b.w.) by reducing temperature up to 4–7%, approximately.

### 3.10. Hepatoprotective Activity

El-Toumy et al. [172] tested the hepatoprotective activity of the MeOH extract of *P. equisetiforme*'s aerial parts on CCl<sub>4</sub>-induced hepatic illness in Sprague-Dawley rats by measuring the levels of serum marker enzyme activities (alanine amino transferase-ALT and aspartate amino transferase-AST) and the oxidative stress mediator levels (NO, malondialdehyde-MDA, glutathione-GSH, glutathione peroxidase-GPx and superoxide dismutase-SOD). The results showed that when the rats were pre-treated with the extract, they exhibited normal levels of ALT and AST (64.86 and 45.16 IU/mL, respectively, at an extract dose of 200 mg/kg) compared to the control (64.86 and 44.22 IU/mL). The GSH, GPx and SOD levels decreased, while the NO and MDA levels increased in comparison to the control. On the other hand, Christopher et al. [119] assessed the hepatotoxicity ability of the MeOH extract obtained from the leaves of *P. minor* on CCl<sub>4</sub> and paracetamol-induced hepatotoxicity in Sprague-Dawley rats. They found that this extract significantly decreased the levels of AST, ALT, ALP and TB in both models compared to the CCl<sub>4</sub> and paracetamol controls. At dose-dependency, the MeOH extract of *P. glabra* roots reduced the marker levels of hepatic injury such as serum glutamate oxaloacetate transaminase (SGOT), serum glutamate pyruvate transaminase (SGPT), alkaline phosphatase (ALKP) and total bilirubin in CCl<sub>4</sub> and paracetamol-induced rats [120].

The anti-fibrotic effects of the aqueous and organic extracts of *P. plebeium* (whole plant) on inflammatory liver disease in CCl<sub>4</sub>-induced rats were tested, and the authors found that the extracts reduced the ALT, AST and gamma-glutamyl transpeptidase ( $\gamma$ GT) levels in CCl<sub>4</sub>-induced rats and restored the CCl<sub>4</sub>-induced tissue fibrosis [190]. In addition, the aqueous extract from the roots of *P. bistorta* exhibited significant hepatoprotective effect in rats with CCl<sub>4</sub>-induced liver damage: the CCl<sub>4</sub> administration on hepatic cells showed hydropic degeneration, swelling, congestion in portal vessels and sinusoids and optically empty cytoplasmic areas and increased the AST, ALT and total bilirubin levels in rats (173 and 223 IU/L and 1.15 mg/dL, respectively) compared to the control (67.7 and 48.2 IU/L and 0.53 mg/dL, respectively). However, the cells recouped when they were treated with aqueous roots extract of *P. bistorta* at a concentration of 100 mg/kg (levels of AST, ALT and total bilirubin decreased at 88.5 and 98.3 IU/L and 0.75 mg/dL, respectively) [46].

The hepatoprotective effects of the MeOH, EtOH and aqueous extracts of *P. amplexicaulis* rhizomes on albino mice were tested [34]. After the administration of CCl<sub>4</sub> to the mice's, ALT, AST, ALP and plasma bilirubin levels increased, and body weight decreased compared to the control; these levels were recouped when administered aqueous, MeOH and EtOH extracts at a dose of 200 mg/kg. ALT, AST, ALP and plasma bilirubin decreased and body weight increased by 16.3, 12.96 and 8.08%, respectively, compared to the control silymarin (10.88% at a dose of 100 mg/kg). Furthermore, the EtOH fruit extract of *P. orientalis* exhibited significant hepatoprotective effects against CCl<sub>4</sub>-induced acute liver injury (ALI) in rats [131]. At doses of 0.5 and 1 g/kg of extract, the levels of AST, ALT, ALP, NO,

tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-1 $\beta$  (IL-1 $\beta$ ) and interleukin-6 (IL-6) decreased, while the levels of SOD, GPx and GRd increased. Protocatechuic acid, taxifolin, and quercetin identified by HPLC in the extracts, might be mainly responsible for these effects. In addition, Fan et al. [132] tested the potential inhibitory ability of OATP1B1 (anion transporting polypeptide 1B1, associated with drug-induced liver injury) of flavonoids isolated from *P. orientalis*. The results demonstrated that the compounds isoorientin and orientin showed weak inhibitory effects on OATP1B1-mediated fluvastatin consumption in OATP1B1-HEK293 cells (27.72 and 23.3%, respectively). Nguyen et al. [135] reported that the EtOAc extract from the aerial parts of *P. pulchra* and its subfractions exhibited a potent hepatoprotective activity in CCl<sub>4</sub>-induced rats since it reduced ALT activity between 50 and 68%. Finally, Xu et al. [189] reported that the total flavonoids isolated from *P. perfoliatum* exhibited potential hepatoprotective effect in CCl<sub>4</sub>-exposed mice, decreasing liver functional enzymes (ALT and AST), TNF- $\alpha$ , interleukin 6 and heat shock protein 90 (Hsp90) and increasing intrasplenic integrin  $\beta$ 1, 5'-nucleotidase and antigen KI-67 cells at doses of 60 and 120 mg/kg/day.

### 3.11. Neuropharmacological Activity

#### 3.11.1. Anti-Alzheimer's

The potential of  $\beta$ -sitosterol isolated from *P. hydropiper* for the management of Alzheimer's disease was tested [90]. Authors observed a significant decline in free radical's load in the brain tissues of the  $\beta$ -sitosterol-treated animals, with IC<sub>50</sub> values of 140, 120, and 280  $\mu$ g/mL from DPPH, ABTS and H<sub>2</sub>O<sub>2</sub> assays, respectively. In addition, they performed memory assessment and coordination tasks (SWM, Y-maze and balance beam tests) and found that  $\beta$ -sitosterol-treated animals had gradual improvement in working memory and in motor coordination [90]. Previously, Ayaz et al. [78] reported that essential oil from the leaves and flowers of this plant demonstrated a significantly acetylcholinesterase (AChE) inhibitory ability (inhibition of 87 and 79.66%; IC<sub>50</sub> = 120 and 220  $\mu$ g/mL, respectively) and a butyrylcholinesterase (BChE) inhibition of 82.66 and 77.5%; IC<sub>50</sub> of 225  $\mu$ g/mL, respectively. Caryophyllene oxide and decahydronaphthalene were identified as the major components of the leaves' and flower's essential oils, respectively. Ahmad et al. [113] tested the AChE inhibition ability of different extracts (MeOH, EtOH, aqueous, DCM and *n*-Hex) and the essential oil of the leaves, stem and root of *P. minor*. Aqueous and MeOH leaf extracts showed the higher AChE inhibitory activity, with IC<sub>50</sub> values of 234 and 342.77  $\mu$ g/mL, respectively; root extracts had the lowest AChE inhibitory activity (IC<sub>50</sub> > 1000  $\mu$ g/mL). The presence of terpenoids could explain AChE activity of MeOH and aqueous extracts, as it can readily cross the blood-brain barrier by its small size and lipophilicity [113]. In addition, a study revealed that the aqueous extract of this plant could reverse scopolamine-induced memory deficits in a Barnes maze model (at a dose of 100 mg/kg) and could inhibit AChE activity by 68% with an IC<sub>50</sub> of 0.04 mg/mL [121]. Regarding *P. glabra*, the MeOH and EtOH extracts of its aerial parts exhibited neuroprotective activity against AlCl<sub>3</sub>-induced (aluminium chloride) toxicity in rats, improving learning and memory and decreasing AChE levels at similar values of the standard rivastigmine at a concentration of 400 mg/kg [69]. Another study stated that the MeOH extracts from roots and aerial parts of *P. maritimum* could be a good alternative for the treatment of neurodegenerative disorders such as Alzheimer's disease due to the ability for the inhibition of AChE and tyrosinase (TYRO), with IC<sub>50</sub> values of 0.17 and 0.27 mg/mL for AChE and 0.59 and 0.6 mg/mL for TYRO [178].

#### 3.11.2. Anti-Depressant and Sedative

The anti-depressant activity of the aqueous extract of *P. glabra* was tested by Nizar et al. [70]. The results showed that the extract, in a dose-dependent manner (50, 100 and 200 mg/kg), induced a significant decrease in the immobility time of mice during behavioural despair test (BDT) and tail suspension test (TST) and increased the hyperactivity scores in

an L-dopa-induced hyperactivity test, with similar values to the reference Imipramine (15 mg/kg). The depressant activity of four sesquiterpenes (viscosumic acid, viscozulenic acid, viscoazucine and viscoazulone) and the flavonoid glycoside quercetin-3-O-(6-feruloyl)- $\beta$ -D-galactopyranoside isolated from the aerial parts of *P. viscosum* was tested by open-field test [196]. Viscoazucine and viscoazulone were the most potent depressants, showing gradual decreases in the movement of mice (No. of movements at 0 min = 143.97 and 137.95 and at 240 min = 23.92 and 27.93, respectively), while the other compounds showed a moderate depressant activity.

*P. hydropiper* showed anti-depressant and sedative activity: Sharif et al. [91] tested the antidepressant activity of the MeOH, EtOH and Chl leaf extracts by open field test and swimming test (mice were forced to swim and scored immobility). At a dose of 150 mg/kg, the Chl and EtOH extracts significantly decreased movements and exerted immobile phase in mice similar to the standard Imipramine (10 mg/kg). Devarajan et al. [92] examined the depressor effect of extracts of *P. hydropiper* leaves in salt-induced hypertension in mice and found that both could reduce blood pressure and heart rate in a dose-dependent manner. Shahed-Al-Mahmud and Lima [93] revealed that the MeOH leaf extract presented a good sedative and anxiolytic activity (at a concentration between 50 and 500 mg/kg) compared to a positive control of diazepam (1 mg/kg) using different test types such as open field test and thiopental sodium-induced sleeping time test in mice (for sedative activity) and elevated-plus maze and light-dark box (for anxiolytic activity). Finally, the ion channel-blocking activity K<sup>+</sup> of G protein-activated inwardly (responsible for maintaining the resting membrane potential and cell excitability) of different extracts of *P. maculosa* were tested through the automated patch-clamp method [109]. They found that Chl extract and its HPLC eluate fractions, at a concentration of 0.1 mg/mL, showed a potent K<sup>+</sup> channel-inhibitory activity compared to the reference compound Propafenone (between 62 and 76% and 71 and 81%, respectively).

### 3.11.3. Neuroprotective Activity

Won and Ma [164] tested the neuroprotective activity of the aqueous-MeOH extract of *P. aviculare* by glutamate-induced neurotoxicity assay in primary cultures of rat cortical cells assay. The results showed that at a concentration of 100  $\mu$ g/mL, a good neuroprotective potency of 50.1% compared to the standards CNQX (59.2%) and MK-801 (70.8%) was observed. Additionally, the juglandin extracted from crude *P. aviculare* exhibited a good neuroprotective activity in mice with LPS-induced Parkinson's disease, attenuating memory impairments, promoting the expression of synaptic markers (SYP, PSD-95 and SNAP-25), decreasing production of pro-inflammatory cytokines (IL-1 $\beta$ , TNF- $\alpha$ , IL-18 and COX-2) and blocking TLR4/NF- $\kappa$ B pathway [165]. In addition, the extract of *P. aviculare* was tested for its neuro-inflammatory properties, and it was found that the extract could decrease lethargy-like behaviour and the compounds corticosterone, serotonin, and catecholamines (fatigue-related) in the brain and inhibited the production protein TNF- $\alpha$  (tumour necrosis factor) [166]. Finally, the neuroprotector effect of orientin (pyrone glucoside extracted from *P. orientalis*) was tested in pheochromocytoma cell line (PC12) stimulated by H<sub>2</sub>O<sub>2</sub> in mice [133]. The results demonstrated that orientin was not toxic for PC12 cells and could reduce the H<sub>2</sub>O<sub>2</sub>-induced viability of PC12 cells at higher concentrations than 40  $\mu$ g/mL. Moreover, orientin decreased H<sub>2</sub>O<sub>2</sub>-induced phosphorylation of signaling proteins (MAPKs, AKT and Src) and inhibited ROS (causing neurodegenerative diseases) accumulation in cells [133].

### 3.12. Diuretic Activity

In a Lipschitz test, the petroleum ether, Chl and EtOAc extracts of *P. barbata*'s aerial parts exhibited dose-dependent diuretic activity, but EtOAc extract (at a dose of 400 mg/kg) showed the most significant effect after 2 h of administration (diuretic activity = 1.77) compared to the standard Furosemide (diuretic activity = 1.72) [35]. In addition, the EtOAc extract of *P. lapathifolium* var. *lanatum* demonstrated a moderate to good diuretic

activity, with values of 1.422 and 1.87 at doses of 150 mg/kg and 300 mg/kg b.w., respectively [98]. Finally, the diuretic activity of  $\alpha$ -Santalone isolated from *P. pubescens* aerial parts was tested by the Lipschitz methods, and it was found that at a concentration of 40 mg/kg, this compound presented a good activity after the first hour of administration (diuretic activity = 1.24) compared to the standard Furosemide 3 mg/kg (diuretic activity = 1.81) [191].

### 3.13. Gastroprotective Activity

The anti-ulcer activity of the alcoholic and aqueous extracts of *P. barbata* were tested by Pylorus ligation models and ethanol-induced gastric mucosal injury in rats [38]. They found that both extracts, in a dose-dependent manner, could significantly reduce the number and index of ulcers and total acidity, as well as increasing the pH index. The compounds present in both extracts such as saponins, sterols, glycosides and alkaloids could explain this activity [38]. In addition, the aqueous extract from the leaves of *P. chinensis* showed a gastroprotective effect against ethanol-induced gastric mucosal injuries in Sprague-Dawley rats, as it reduced gastric lesions and malondialdehyde levels (MDA) and increased superoxide dismutase level (SOD) [57]. The anti-ulcer activity of the aqueous extract from the leaves of *P. minor* against ethanol-induced gastric ulcers in rats was tested [122]. The extract showed a significant anti-ulcer activity compared to the standard omeprazole (20 mL/kg): at dose-dependency (250 and 500 mL/kg), the extract increased pH and gastric mucous, suppressed areas of gastric ulcer formation (35.33–188.17 mm) and inhibited 78.25–95.92% of the gastric ulcer. In addition, Qader et al. [123] obtained five fractions (hexane:ethyl acetate 1:1 v/v (F1), ethyl acetate:methanol 1:1 v/v (F2), methanol:acetonitrile 1:1 v/v (F3), acetonitrile:distilled water 1:1 v/v (F4) and distilled water 1:1 v/v (F5)) from the EtOH extract of the leaves of *P. minor* and tested its gastroprotective activity using the ethanol induction method in rats. All the fractions exhibited gastroprotective activity, but F2 showed the best result at a dose-dependency (very similar to Omeprazole values), inhibiting 90% of ulcer lesions and increasing mucus content (120 mg/g), SOD, hexosamine and PGE2 synthesis levels in the stomach wall mucosa.

The gastroprotective effect of the hydro-alcoholic root extract of *P. bistorta* was tested by indomethacin-induced gastric ulcer in rats [47]. At a dose-dependency (500 and 1000 mg/kg), this extract significantly increased mucus, SOD and catalase levels and decreased the ulcer index and thiobarbituric acid (TBARS), with similar values shown by the standard drug ranitidine (20 mg/kg), compared to the ulcer control group. Ayaz et al. [74] tested the gastroprotective ability of the crude MeOH extract from *P. hydropiper* and its fractions by aspirin-induced ulcerogenesis in rats. At a dose dependency (100, 200 and 400 mg/kg), the extract exhibited a good gastroprotective activity compared to the standard ranitidine (50 mg/kg), as it could decrease gastric juice volume, free acidity, total acidity and pepsin levels, as well as increased gastric juice pH levels. Additionally, the essential oil obtained from the leaves and saponins (fractions) showed the highest urease inhibition (> 70%, IC<sub>50</sub> = 90 and 98  $\mu$ g/mL, respectively) compared to the control drug thiourea (urease inhibition > 80% and IC<sub>50</sub> = 80  $\mu$ g/mL) [74].

### 3.14. Other Activities

The acetone extract from the aerial parts of *P. maritimum* is considered an interesting anti-melanogenic agent, as it was demonstrated to inhibit tyrosinase and melanin production in B16 4A5 melanoma cells, with IC<sub>50</sub> values of 64.1 and 77.7  $\mu$ g/mL, respectively [176]. Myricitrin, catechin and monogalloyl-hexose isolates could be responsible for this activity. Another study conducted by George et al. [124] explored the immunomodulatory properties of aqueous extract of *P. minor*, and they showed that this extract displayed significant phagocytic index (K) at doses of 200 and 400 mg/kg b.w. (K = 0.045 and 0.062, respectively) compared with the standard levamisole (K = 0.060). Finally, Kimura et al. [155] tested the inhibitory activity of flavonol O-glycosides isolated from 80% MeOH extracts of *P. tinctoria* leaves against HMG-CoA reductase. The fraction eluted with 100%

MeOH showed the most potent inhibitory activity (56.7%), while five compounds isolated from this fraction were effective dose-dependently, inhibiting HMG-CoA reductase activity by 50 to 67.6%.

#### 4. Conclusions

This *Polygonum* and *Persicaria* genera revision revealed the great variety of chemical constituents present in these plants, highlighting the bioactive groups of sesquiterpenes, flavonoids and phenolic acids. Methanolic, ethanolic, hexanic, ethyl acetate and water extracts and their fractions and chemical phytoconstituents have demonstrated different pharmacological activities such as antifungal, antibacterial, antioxidant, anti-inflammatory, anticancer and neuropharmacological activities in many publications, which were updated and revised here. These findings revealed that the species of the *Persicaria* and *Polygonum* genera could be properly developed as good candidates for clinical assays in the future, allowing for the expansion of knowledge for the treatment of new diseases. *P. glabra*, *P. hydropiper*, *P. minor*, *P. lapathifolia* and *P. chinensis* were the species that showed the highest number of medicinal properties. Native species of Asia and Europe are ones with the largest number of studies in the world, followed by African species. However, American species are poorly studied or do not present any study (e.g., *P. ferruginea*, *P. hydropiperoides*, *P. punctata* and *P. paraguayense*), which invites the investigation of these species in the future.

The wide range of pharmacological properties of *Polygonum* and *Persicaria* species may offer a new therapeutic promise to cure different diseases and health complications. So, an accelerated progress should be made through experimental research including robust clinical trials, in order to generate natural medicines that allow to counter the negative effects on human health.

**Author Contributions:** The individual contributions of each author are as follows: Conceptualization, G.S., N.Á. and M.D.; formal analysis, G.S., L.F., M.I.S. and M.D.; investigation, G.S., N.Á., L.F. and M.A.F.; resources, M.A.F., V.R. and M.D.; writing—original draft preparation, G.S. and N.Á.; writing—review and editing, V.R. and M.D.; supervision, M.A.F., V.R. and M.D.; project administration, M.A.F., V.R. and M.D.; funding acquisition, V.R. and M.D. All authors have read and agreed to the published version of the manuscript.

**Funding:** Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, PIP N° 524), Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT, PICT 2015–2259) and ASaCTel.

**Acknowledgments:** G.S., N.Á., M.I.S. and L.F. thank CONICET for their fellowships.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

#### References

1. Centers for Disease Control and Prevention. *Addressing emerging infectious disease threats: A prevention strategy for the United States (Executive Summary)*. GPO: Washington DC., USA, 1994; 43, 1–18.
2. Usha, P.T.A.; Jose, S.; Nisha, A.R. Antimicrobial Drug Resistance—A global concern. *Vet. World*. **2010**, *3*, 138–139.
3. Lizcano González, M.C. Evaluación de la actividad antifúngica del extracto de tomillo (*Thymus vulgaris*) contra *Botrytis cinerea*, *Fusarium oxysporum* y *Sclerotinia sclerotiorum*. Doctoral Thesis, Pontificia Universidad Javeriana, Bogotá, 2007.
4. Freiesleben, S.; Jäger, A. Correlation between plant secondary metabolites and their antifungal mechanisms—A review. *Med. Aromat. Plants*. **2014**, *3*, 1–6.
5. Farnsworth, N.R.; Soejarto, D.D. *Global importance of medicinal plants*. In *Conservation of Medicinal Plants*; Akerele, O., Heywood, V., Synge, H., Eds.; Cambridge University Press: Cambridge, England, 1991; Volume 1, pp. 25–51.
6. Chanda, S.; Rakholiya, K. *Combination therapy: Synergism between natural plant extracts and antibiotics against infectious diseases*. In *Science against microbial pathogens: Communicating current research and technological advances*; Méndez-Vilas, A., Ed.; Formatex Research Center: Badajoz, Spain, 2011; Volume 1, pp. 520–529.
7. Derita, M.; Zacchino, S. Chemotaxonomic importance of sesquiterpenes and flavonoids in Argentinian species of *Polygonum* genus. *J. Essent. Oil Res.* **2011**, *23*, 11–14.

8. López, S.N.; Furlan, R.L.E.; Zacchino, S.A. Detection of antifungal compounds in *Polygonum ferrugineum* Wedd. extracts by bioassay-guided fractionation. Some evidences of their mode of action. *J. Ethnopharmacol.* **2011**, *138*, 633–636, doi:10.1016/j.jep.2011.09.038.
9. Saifuzzaman, M.; Shamim, M.S.J.; Mahaldar, K.; Ali, E.S.; Islam, M.A. Antinociceptive activity of the ethanolic extract of *Persicaria acuminata* Sach. *J. Pharm. Res.* **2013**, *6*, 753–755, doi:10.1016/j.jopr.2013.07.005.
10. Derita, M.; Leiva, M.; Zacchino, S. Influence of plant part, season of collection and content of the main active constituent, on the antifungal properties of *Polygonum acuminatum* Kunth. *J. Ethnopharmacol.* **2009**, *124*, 377–383, doi:10.1016/j.jep.2009.05.029.
11. Derita, M.; Zacchino, S. Validation of the ethnopharmacological use of *Polygonum persicaria* for its antifungal properties. *Nat. Prod. Commun.* **2011**, *6*, 931–933.
12. Agüero, E. *Plantas medicinales silvestres del centro de Argentina: Guía para su reconocimiento y uso terapéutico*; Ecoval Ediciones: Córdoba, Argentina, 2017; pp. 443.
13. Lahitte, H.B.; Hurrell, J.A.; Belgrano, M.J.; Jankowski, L.S.; Haloua, M.P.; Mehlreter, K. *Plantas medicinales rioplatenses: Plantas nativas y naturalizadas utilizadas en medicina popular en la región del Delta del Paraná, Isla Martín García y Ribera Platense*; L.O.L.A.: Buenos Aires, Argentina, 1998; pp. 240.
14. Gupta, P.M. *270 plantas medicinales iberoamericanas, Programa Iberoamericano de Ciencia y Tecnología para el desarrollo*; Subprograma de Química Fina Farmacéutica: Bogotá, Colombia, 1994; pp. 567.
15. Baharum, S.N.; Bunawan, H.; Abd Ghani, M.; Mustapha, W.A.W.; Noor, M.N. Analysis of the chemical composition of the essential oil of *Polygonum minus* Huds. using two-dimensional gas chromatography–time-of-flight mass spectrometry (GC–TOF MS). *Molecules* **2010**, *15*, 7006–7015, doi:10.3390/molecules15107006.
16. Huda-Faujan, N.; Norriham, A.; Norrakiah, A.S.; Babji, A.S. Antioxidative activities of water extracts of some Malaysian herbs. *ASEAN Food J.* **2007**, *14*, 61–68.
17. Sun, X.; Sneden, A.T. Neoflavonoids from *Polygonum perfoliatum*. *Planta Med.* **1999**, *65*, 671–673, doi:10.1055/s-2006-960846.
18. Sun, X.B.; Zhao, P.H.; Xu, Y.J.; Sun, L.M.; Cao, M.A.; Yuan, C.S. Chemical constituents from the roots of *Polygonum bistorta*. *Chem. Nat. Compd.* **2007**, *43*, 563–566, doi:10.1007/s10600-007-0193-z.
19. Bouzada, M.L.; Fabri, R.L.; Nogueira, M.; Konno, T.U.; Duarte, G.G.; Scio, E. Antibacterial, cytotoxic and phytochemical screening of some traditional medicinal plants in Brazil. *Pharm. Biol.* **2009**, *47*, 44–52, doi:10.1080/13880200802411771.
20. Wang, K.W.; Zhu, J.R.; Shen, L.Q. A new lignan with anti-tumour activity from *Polygonum perfoliatum* L. *Nat. Prod. Res.* **2013**, *27*, 568–573, doi:10.1080/14786419.2012.682993.
21. Maizura, M.; Aminah, A.; Wan Aida, W.M. Total phenolic content and antioxidant activity of kesum (*Polygonum minus*), ginger (*Zingiber officinale*) and turmeric (*Curcuma longa*) extract. *Int. Food Res. J.* **2011**, *18*, 526–531.
22. Takasaki, M.; Kuroki, S.; Kozuka, M.; Konoshima, T. New phenylpropanoid esters of sucrose from *Polygonum lapathifolium*. *J. Nat. Prod.* **2001**, *64*, 1305–1308, doi:10.1021/np010222q.
23. Zhang, X.; Liu, F.; Feng, Z.M.; Yang, Y.N.; Jiang, J.S.; Zhang, P.C. Bioactive phenylpropanoid esters of sucrose and anthraquinones from *Polygonum cuspidatum*. *Fitoterapia* **2020**, *146*, 104673, doi:10.1016/j.fitote.2020.104673.
24. Wang, K.J.; Zhang, Y.J.; Yang, C.R. Antioxidant phenolic compounds from rhizomes of *Polygonum paleaceum*. *J. Ethnopharmacol.* **2005**, *96*, 483–487, doi:10.1016/j.jep.2004.09.036.
25. Derita, M.; Montenegro, I.; Garibotto, F.; Enriz, R.; Cuellar Fritis, M.; Zacchino, S. Structural requirements for the antifungal activities of natural drimane sesquiterpenes and analogues, supported by conformational and electronic studies. *Molecules* **2013**, *18*, 2029–2051, doi:10.3390/molecules18022029.
26. Calderón, Á.I.; Romero, L.I.; Ortega-Barría, E.; Solís, P.N.; Zacchino, S.; Gimenez, A.; Pinzón, R.; Cáceres, A.; Tamayo, G.; Guerra, C.; Espinosa, A.; Correa, M.; Gupta, M.P. Screening of Latin American plants for antiparasitic activities against malaria, Chagas disease, and leishmaniasis. *Pharm. Biol.* **2010**, *48*, 545–553, doi:10.3109/13880200903193344.
27. Mir, P.A.; Mohi-u-Din, R.; Dar, M.A.; Bader, G.N. Anti-inflammatory and anti-helminthic potential of methanolic and aqueous extract of *Polygonum alpinum* rhizomes. *J. Drug Deliv. Ther.* **2019**, *9*, 455–459, doi:10.22270/jddt.v9i3.2884.
28. Borchardt, J.R.; Wyse, D.L.; Sheaffer, C.C.; Kauppi, K.L.; Ehlke, N.J.; Biesboer, D.D.; Bey, R.F. Antimicrobial activity of native and naturalized plants of Minnesota and Wisconsin. *J. Med. Plants Res.* **2008**, *2*, 98–110, doi:10.5897/JMPR.9000165.
29. Smolarz, H.D.; Budzianowski, J.; Bogucka-Kocka, A.; Kocki, J.; Mendyk, E. Flavonoid glucuronides with anti-leukaemic activity from *Polygonum amphibium* L. *Phytochem. Anal.* **2008**, *19*, 506–513, doi:10.1002/pca.1076.
30. Tantry, M.A.; Radwan, M.M.; Akbar, S.; Khan, I.A. 5,6-Dihydropyranobenzopyrone: A previously undetermined antioxidant isolated from *Polygonum amplexicaule*. *Chin. J. Nat. Med.* **2012**, *10*, 28–31, doi:10.1016/S1875-5364(12)60006-5.
31. Batool, S.; Gulfranz, M.; Naqvi, S.M.; Mirza, B.; Ahmad, M.S. Evaluation of antioxidant potential and HPLC based identification of phenolics in *Polygonum amplexicaule* extract and its fractions. *Pak. J. Pharm. Sci.* **2015**, *28*, 431–435.
32. Xiang, M.; Su, H.; Hong, Z.; Yang, T.; Shu, G. Chemical composition of total flavonoids from *Polygonum amplexicaule* and their pro-apoptotic effect on hepatocellular carcinoma cells: Potential roles of suppressing STAT3 signaling. *Food Chem. Toxicol.* **2015**, *80*, 62–71, doi:10.1016/j.fct.2015.02.020.
33. Xiang, M.; Su, H.; Shu, G.; Wan, D.; He, F.; Loaec, M.; Ding, Y.; Li, J.; Dovat, S.; Yang, G.; Song, C. Amplexicaule A exerts anti-tumor effects by inducing apoptosis in human breast cancer. *Oncotarget* **2016**, *7*, 18521–18530, doi:10.18632/oncotarget.7848.
34. Maqsood, F.; Ibrahim, T.; Farooqi, A.A.; Ahmad, M.S. *Polygonum amplexicaule* extract: An effective herbal cure to CCL. *J. Rare Disord. Diagn. Ther.* **2017**, *3*, 4, doi:10.21767/2380-7245.100057.

35. Mazid, M.A.; Datta, B.K.; Nahar, L.; Bashar, S.A.M.; Bachar, S.C.; Sarker, S.D. Antinociceptive, anti-inflammatory and diuretic properties of *Polygonum barbatum* (L.) Hara var. *barbata*. *Rev. Bras. Farmacogn.* **2009**, *19*, 749–754.
36. Farooq, U.; Naz, S.; Shams, A.; Raza, Y.; Ahmed, A.; Rashid, U.; Sadiq, A. Isolation of dihydrobenzofuran derivatives from ethnomedicinal species *Polygonum barbatum* as anticancer compounds. *Biol. Res.* **2019**, *52*, 1, doi:10.1186/s40659-018-0209-0.
37. Mazid, A.M.; Nahar, L.; Datta, B.K.; Bashar, K.S.; Sarker, S.D. Potential antitumor activity of two *Polygonum* species. *Arch. Biol. Sci.* **2011**, *63*, 465–468, doi:10.2298/ABS1102465M.
38. Kinger H.K.; Gupta M.K. Evaluation of anti-ulcer activity of *Polygonum barbatum* Linn. (whole plant). *J. Biomed. Pharmac. Res.* **2012**, *1*, 34–37.
39. Munir, N.; Ijaz, W.; Altaf, I.; Naz, S. Evaluation of antifungal and antioxidant potential of two medicinal plants: *Aconitum heterophyllum* and *Polygonum bistorta*. *Asian Pac. J. Trop. Biomed.* **2014**, *4*, S639–S643, doi:10.12980/APJTB.4.201414B182.
40. Safavi, E.; Homayouni-Tabrizi, M.; Karimi, E.; Mohammad, G.R.K.S. Biosynthesis of zinc oxide nanoparticles using anjbar (root of *Persicaria bistorta*) extract and their cytotoxic effects on human breast cancer cell line (MCF-7). *IET Nanobiotechnol.* **2019**, *13*, 736–741, doi:10.1049/iet-nbt.2018.5060.
41. Huang, Y.S., H.; Zeng, J.; Ye, H.; Huang, X.; Lai, F. A study on the analgesic effect of *Polygonum Bistorta* L. *n*-butyl alcohol. *J. Gannan Med. Coll.* **2004**, *1*, 12–13.
42. Intisar, A.; Zhang, L.; Luo, H.; Kiazolu, J.B.; Zhang, R.; Zhang, W. Anticancer constituents and cytotoxic activity of methanol-water extract of *Polygonum bistorta* L. *Afr. J. Tradit. Complement. Altern. Med.* **2013**, *10*, 53–59, doi:10.4314/ajtcam.v10i1.9.
43. Liu, Y.H.; Weng, Y.P.; Lin, H.Y.; Tang, S.W.; Chen, C.J.; Liang, C.J.; Ku, C.Y.; Lin, J.Y. Aqueous extract of *Polygonum bistorta* modulates proteostasis by ROS-induced ER stress in human hepatoma cells. *Sci. Rep.* **2017**, *7*, 41437, doi:10.1038/srep41437.
44. Pillai Manoharan, K.; Yang, D.; Hsu, A.; Tan Kwong Huat, B. Evaluation of *Polygonum bistorta* for anticancer potential using selected cancer cell lines. *Med. Chem.* **2007**, *3*, 121–126.
45. Kumar, M.D.; Deepmala, J.; Sangeeta, S. Antioxidant, antipyretic and choleric activities of crude extract and active compound of *Polygonum Bistorta* (Linn.) in albino rats. *Int. J. Pharm. Bio. Sci.* **2012**, *2*, 25–31.
46. Mittal, D.K.; Joshi, D. Evaluation of DNA damage by comet and mtt-assay in liver treated with *Polygonum bistorta* and tannic acid. *Int. J. Pharm. Bio. Sci.* **2012**, *3*, 375–382.
47. Khushtar, M.; Ahmad, A.; Rahman, M.A. Gastroprotective effect of hydro-alcoholic extract of *Polygonum bistorta* Lin root in indomethacin-induced gastric ulcers in sprague dawley rats. *Indian J. Pharm. Educ. Res.* **2018**, *52*, 618–625, doi:10.5530/ijper.52.4.72.
48. Liao, S.G.; Zhang, L.J.; Sun, F.; Zhang, J.J.; Chen, A.Y.; Lan, Y.Y.; Dong, L. Antibacterial and anti-inflammatory effects of extracts and fractions from *Polygonum capitatum*. *J. Ethnopharmacol.* **2011**, *134*, 1006–1009, doi:10.1016/j.jep.2011.01.050.
49. Lai, S.M.; Sudhahar, D.; Anandarajagopal, K. Evaluation of antibacterial and antifungal activities of *Persicaria chinensis* leaves. *Int. J. Pharm. Sci. Res.* **2012**, *3*, 2825–2830, doi:10.13040/IJPSR.0975-8232.3(8).2825-30.
50. Srividya, A.R.; Shalom, A.; Chandrasekhar, R.; Vijayan, P.; Vishnuvarthan, V.J. Cytotoxic, antioxidant and antimicrobial activity of *Polygonum chinensis* Linn. *Int. J. Pharm. Sci. Nanotechnol.* **2012**, *4*, 1569–1574, doi:10.37285/ijpsn.2011.4.4.8.
51. Zhang, H.W.; Wang, Y.; Du, X.W.; Ding, G.; Zou, Z.M. Chemical constituents of whole plant of *Polygonum chinense* L. var *hispidum* hook. f. *Chin. Pharm J.* **2015**, *50*, 1012–1016.
52. Wu, Y.; Zhang, Z.; Chen, T.; Cheng, C.; Zhang, Z.; Zhou, H.; Luo, P. Comparison of two *Polygonum chinense* varieties used in Chinese cool tea in terms of chemical profiles and antioxidant/anti-inflammatory activities. *Food Chem.* **2020**, *310*, 125840, doi:10.1016/j.foodchem.2019.125840.
53. Hossen, M.J.; Baek, K.S.; Kim, E.; Yang, W.S.; Jeong, D.; Kim, J.H.; Kweon, D.H.; Yoon, D.H.; Kim, T.W.; Kim, J.H.; Cho, J.Y. In vivo and in vitro anti-inflammatory activities of *Persicaria chinensis* methanolic extract targeting Src/Syk/NF- $\kappa$ B. *J. Ethnopharmacol.* **2015**, *159*, 9–16, doi:10.1016/j.jep.2014.10.064.
54. Chen, W.; Shen, X.; Ma, L.; Chen, R.; Yuan, Q.; Zheng, Y.; Li, C.; Peng, G. Phenolic compounds from *Polygonum chinense* induce growth inhibition and apoptosis of cervical cancer SiHa cells. *BioMed Res. Int.* **2020**, doi:10.1155/2020/8868508.
55. Tran, T.T.; Kim, M.; Jang, Y.; Lee, H.W.; Nguyen, H.T.; Nguyen, T.N.; Park, H.W.; Le Dang, Q.; Kim, J.C. Characterization and mechanisms of anti-influenza virus metabolites isolated from the Vietnamese medicinal plant *Polygonum chinense*. *BMC Complement. Alter. Med.* **2014**, *17*, 162, doi:10.1186/s12906-017-1675-6.
56. Lai, S.M.; Sudhahar, D.; Anandarajagopal, K. In vitro anthelmintic potential of *Persicaria chinensis* leaves. *Int. J. Biol. Pharm. Res.* **2012**, *3*, 400–404.
57. Ismail, I.F.; Golbabapour, S.; Hassandarvish, P.; Hajrezaie, M.; Abdul Majid, N.; Kadir, F.A.; Al-Bayaty, F.; Awang, K.; Hazni, H.; Abdulla, M.A. Gastroprotective activity of *Polygonum chinense* aqueous leaf extract on ethanol-induced hemorrhagic mucosal lesions in rats. *Evid. Based Complement. Alternat. Med.* **2012**, *2012*, 1–9, doi:10.1155/2012/404012.
58. Youssef, A.M.M.; El-Swaify, Z.A.S. Anti-Tumour Effect of two *Persicaria* species seeds on colon and prostate cancers. *Biomed. Pharmacol. J.* **2018**, *11*, 635–644, doi:10.13005/bpj/1416.
59. Zheoat, A.M.; Alenezi, S.; Elmahallawy, E.K.; Ungogo, M.A.; Alghamdi, A.H.; Watson, D.G.; Igoli, J.O.; Gray, A.I.; Koning, H.P.; Ferro, V.A. Antitrypanosomal and antileishmanial activity of chalcones and flavanones from *Polygonum salicifolium*. *Pathogens* **2021**, *10*, 175, doi:10.3390/pathogens10020175.
60. Brandão, G.C.; Kroon, E.G.; Duarte, M.G.R.; Braga, F.C.; de Souza Filho, J.D.; de Oliveira, A.B. Antimicrobial, antiviral and cytotoxic activity of extracts and constituents from *Polygonum spectabile* Mart. *Phytomedicine* **2010**, *17*, 926–929, doi:10.1016/j.phymed.2010.03.004.

61. Rodrigues, D.F.; Maniscalco, D.A.; Silva, F.A.; Chiari, B.G.; Castelli, M.V.; Isaac, V.L.; Cicarelli, R.M.B.; Lopez, S.N. Trypanocidal activity of flavokawin B, a component of *Polygonum ferrugineum* Wedd. *Planta Med.* **2017**, *83*, 239–244, doi:10.1055/s-0042-112031.
62. Jani, M.; Shah, S.; Prajapati, S. Antibacterial screening and qualitative phytochemical estimation of selected aquatic plants. *Adv. Biol. Res.* **2012**, *6*, 19–23, doi:10.5829/idosi.abr.2012.6.1.56177.
63. Said, M.S.; Chinchansure, A.A.; Nawale, L.; Durge, A.; Wadhvani, A.; Kulkarni, S.S.; Sarkar, D.; Joshi, S.P. A new butenolide cinnamate and other biological active chemical constituents from *Polygonum glabrum*. *Nat. Prod. Res.* **2015**, *29*, 2080–2086, doi:10.1080/14786419.2015.1004674.
64. Palani, R.; Karunakaran, D.; Rajesh, V.; Mathivanan, K.; Jayaraman, P. Analysis of antioxidant, antimicrobial activity and phytochemical potential of *Cleistanthus collinus* Roxb., *Polygonum glabrum* Willd. and *Melia azedarach* Linn. *Asian J. Med. Pharm. Sci.* **2014**, *2*, 149–159.
65. Rajamanickam, M.; Rajamohan, S. Analgesic activity of flavonoids isolated from *Persicaria glabra* (wild). *Adv. Tradit. Med.* **2020**, *20*, 71–76, doi:10.1007/s13596-019-00404-x.
66. Ali, A.M. Phytochemical and pharmacological studies of selected Sudanese medicinal plants with emphasis on *Polygonum glabrum*. Ph.D. Thesis, University of Khartoum, Khartoum, Sudan, 2003.
67. Rahman, H.U.; Rehman, T.U.; Ali, A.; Shah, S.A.A.; Ismail, M. In vitro anti-leishmanial activity of *Polygonum glabrum* stem extract on *Leishmania tropica* (KWH23) Strain. *J. Adv. Biol. Biotechnol.* **2015**, *3*, 23–28, doi:10.9734/JABB/2015/15204.
68. Jamal, B.D.; Avinash, K.R.G.; Naganjenulu, R.; Jyothi, M.J.; Kalishwari, E.; Anvesh, M. Phytochemical screening and antipyretic activity of root stocks of *Polygonum glabrum* Willd in rats. *Int. J. Pharmacother.* **2011**, *1*, 1–4.
69. Suneesha, Y.; Vinay Kumar, T. Evaluation of anti-alzheimer activity of ethanolic and methanolic extracts of *Polygonum Glabrum* against Aluminum chloride induced Alzheimer's in experimental rats. *Asian J. Pharm. Clin. Res.* **2021**, *4*, 118–125.
70. Nizar, K.; Sanghamitra, M.; Tiwary, M.P.; Singh, P.N.; Kumar, V. Antidepressant activity and brain neurotransmitters study of *Polygonum glabrum* Willd in rodents. *J. Herb. Med. Toxicol.* **2007**, *1*, 73–79.
71. Duraipandiyar, V.; Indwar, F.; Ignacimuthu, S. Antimicrobial activity of confertifolin from *Polygonum hydropiper*. *Pharm. Biol.* **2010**, *48*, 187–190, doi:10.3109/13880200902902471.
72. Hasan, M.F.; Das, R.; Khan, A.; Hossain, M.S.; Rahman, M. The determination of antibacterial and antifungal activities of *Polygonum hydropiper* (L.) root extract. *Adv. Biol. Res.* **2009**, *3*, 53–56.
73. Lee, S.H.; Lee, J.R.; Lunde, C.S.; Kubo, I. In vitro antifungal susceptibilities of *Candida albicans* and other fungal pathogens to polygodial, a sesquiterpene dialdehyde. *Planta Med.* **1999**, *65*, 204–208, doi:10.1055/s-1999-13981.
74. Ayaz, M.; Junaid, M.; Ullah, F.; Sadiq, A.; Shahid, M.; Ahmad, W.; Ullah, I.; Ahmad, A.; Syed, N.I.H. GC-MS analysis and gastroprotective evaluations of crude extracts, isolated saponins, and essential oil from *Polygonum hydropiper* L. *Front. Chem.* **2017**, *5*, 1–11, doi:10.3389/fchem.2017.00058.
75. Nasir, A.; Khan, M.; Rehman, Z.; Khalil, A.A.K.; Farman, S.; Begum, N.; Irfan, M.; Sajjad, W.; Parveen, Z. Evaluation of alpha-amylase inhibitory, antioxidant, and antimicrobial potential and phytochemical contents of *Polygonum hydropiper* L. *Plants* **2020**, *9*, 852, doi:10.3390/plants9070852.
76. Hashim, N.H.N.; Abas, F.; Shaari, K.; Lajis, N.H. LC–DAD–ESIMS/MS characterization of antioxidant and anticholinesterase constituents present in the active fraction from *Persicaria hydropiper*. *LWT-Food Sci. Technol.* **2012**, *46*, 468–476, doi:10.1016/j.lwt.2011.11.021.
77. Peng, Z.F.; Strack, D.; Baumert, A.; Subramaniam, R.; Goh, N.K.; Chia, T.F.; Tan, S.N.; Chia, L.S. Antioxidant flavonoids from leaves of *Polygonum hydropiper* L. *Phytochemistry* **2003**, *62*, 219–228, doi:10.1016/S0031-9422(02)00504-6.
78. Ayaz, M.; Junaid, M.; Ullah, F.; Sadiq, A.; Khan, M.A.; Ahmad, W.; Shah, M.R.; Imran, M.; Ahmad, S. Comparative chemical profiling, cholinesterase inhibitions and anti-radicals' properties of essential oils from *Polygonum hydropiper* L: A Preliminary anti-Alzheimer's study. *Lipids Health Dis.* **2015**, *14*, 1–12, doi:10.1186/s12944-015-0145-8.
79. Yang, Y.; Yu, T.; Jang, H.J.; Byeon, S.E.; Song, S.Y.; Lee, B.H.; Rhee, M.H.; Kim, T.W.; Lee, J.; Hong, S.; Cho, J.Y. In vitro and in vivo anti-inflammatory activities of *Polygonum hydropiper* methanol extract. *J. Ethnopharmacol.* **2012**, *139*, 616–625, doi:10.1016/j.jep.2011.12.003.
80. Zhang, W.; Pan, Y.; Qu, S.; Wang, D.; Cheng, S.; Liu, X. Anti-inflammatory effects of an extract of *Polygonum hydropiper* stalks on 2, 4, 6-trinitrobenzenesulphonic acid-induced intestinal inflammation in rats by inhibiting the NF- $\kappa$ B pathway. *Mediat. Inflamm.* **2018**, *2018*, doi:10.1155/2018/6029135.
81. Rahman, E.; Goni, S.A.; Rahman, M.T.; Ahmed, M. Antinociceptive activity of *Polygonum hydropiper*. *Fitoterapia* **2002**, *73*, 704–706, doi:10.1016/S0367-326X(02)00239-3.
82. Oany, A.R.; Hossain, M.U.; Islam, R.; Emran, A.A. A preliminary evaluation of cytotoxicity, antihyperglycemic and antinociceptive activity of *Polygonum hydropiper* L. ethanolic leaf extract. *Clin. Phytosci.* **2017**, *2*, 2, doi:10.1186/s40816-016-0016-5.
83. Ayaz, M.; Junaid, M.; Ullah, F.; Sadiq, A.; Subhan, F.; Khan, M.A.; Ahmad, W.; Ali, G.; Imran, M.; Ahmad, S. Molecularly characterized solvent extracts and saponins from *Polygonum hydropiper* L. show high anti-angiogenic, anti-tumor, brine shrimp, and fibroblast NIH/3T3 cell line cytotoxicity. *Front. Pharmacol.* **2016**, *7*, 74, doi:10.3389/fphar.2016.00074.
84. Ayaz, M.; Sadiq, A.; Wadood, A.; Junaid, M.; Ullah, F.; Khan, N.Z. Cytotoxicity and molecular docking studies on phytosterols isolated from *Polygonum hydropiper* L. *Steroids* **2019**, *141*, 30–35, doi:10.1016/j.steroids.2018.11.005.
85. Raihan, M.O.; Khalequezzaman, M.; Brishti, A.; Tareq, S.M.; Hossain, A.; Rana, S. Anthelmintic and antiproliferative activity of aerial parts of *Persicaria hydropiper*. *Der Pharmacia Sinica.* **2012**, *3*, 104–110.

86. Munira, S.; Sultana, F.; Mollika, S.; Nesa, L.; Ria, J.; Islam, M.M. Anti-neoplastic activities of *Polygonum hydropiper* Linn. against ehrlich ascites carcinoma in Swiss Albino mice. *Adv. Biol. Res.* **2013**, *7*, 09–14, doi:10.5829/idosi.abr.2013.7.1.71160.
87. Xiao, H.; Rao Ravu, R.; Tekwani, B.L.; Li, W.; Liu, W.B.; Jacob, M.R.; Khan, S.I.; Cai, X.; Peng, C.Y.; Khan, I.A.; Li, X.C.; Wang, W. Biological evaluation of phytoconstituents from *Polygonum hydropiper*. *Nat. Prod. Res.* **2017**, *31*, 2053–2057, doi:10.1080/14786419.2016.1269094.
88. Ayaz, M.; Junaid, M.; Subhan, F.; Ullah, F.; Sadiq, A.; Ahmad, S.; Imran, M.; Kamal, Z.; Hussain, S.; Shah, S.M. Heavy metals analysis, phytochemical, phytotoxic and anthelmintic investigations of crude methanolic extract, subsequent fractions and crude saponins from *Polygonum hydropiper* L. *BMC Complement. Alter. Med.* **2014**, *14*, 465, doi:10.1186/1472-6882-14-465.
89. Akhter, R.; Haque, M.A.; Bhuiyan, M.A.; Shahriar, M. In vivo Pharmacological investigation of leaf of *Polygonum hydropiper* (L.). *Dhaka Univ. J. Pharm. Sci.* **2013**, *12*, 165–169, doi:10.3329/dujps.v12i2.17619.
90. Ayaz, M.; Junaid, M.; Ullah, F.; Subhan, F.; Sadiq, A.; Ali, G.; Ovais, M.; Shahid, M.; Ahmad, A.; Wadood, A.; El-Shazly, M.; Ahmad, N.; El-Shazly, M. Anti-Alzheimer's studies on  $\beta$ -sitosterol isolated from *Polygonum hydropiper* L. *Front. Pharmacol.* **2017**, *8*, 697, doi:10.3389/fphar.2017.00697.
91. Sharif, S.; Shahriar, M.; Haque, M.A.; Chowdhury, Z.S.; Islam, M.R.; Bhuiyan, M.A. In-vitro antioxidant activities, anti-nociceptive and neuropharmacological activities of *Polygonum hydropiper*. *J. Biol. Agric. Healthcare.* **2013**, *3*, 61–71.
92. Devarajan, S.; Yahiro, E.; Uehara, Y.; Kuroda, R.; Hirano, Y.; Nagata, K.; Miura, S.; Saku, K.; Urata, H. Depressor effect of the young leaves of *Polygonum hydropiper* Linn. in high-salt induced hypertensive mice. *Biomed. Pharmacother.* **2018**, *102*, 1182–1187, doi:10.1016/j.biopha.2018.03.055.
93. Shahed-Al-Mahmud, M.; Lina, S.M.M. Evaluation of sedative and anxiolytic activities of methanol extract of leaves of *Persicaria hydropiper* in mice. *Clin. Phytosci.* **2017**, *3*, 20, doi:10.1186/s40816-017-0056-5.
94. Bussmann, R.W.; Glenn, A.; Sharon, D. Antibacterial activity of medicinal plants of Northern Peru—can traditional applications provide leads for modern science. *Indian J. Tradit. Knowl.* **2010**, *9*, 742–753.
95. Braga, F.G.; Bouzada, M.L.M.; Fabri, R.L.; Matos, M.D.O.; Moreira, F.O.; Scio, E.; Coimbra, E.S. Antileishmanial and antifungal activity of plants used in traditional medicine in Brazil. *J. Ethnopharmacol.* **2007**, *111*, 396–402, doi:10.1016/j.jep.2006.12.006.
96. Hailemariam, A.; Feyera, M.; Deyou, T.; Abdissa, N. Antimicrobial Chalcones from the Seeds of *Persicaria lapathifolia*. *Biochem. Pharmacol.* **2018**, *7*, 2167–0501, doi:10.4172/2167-0501.1000237.
97. Kubinova, R.; Pořízková, R.; Bartl, T.; Navratilova, A.; Cížek, A.; Valentova, M. Biological activities of polyphenols from *Polygonum lapathifolium*. *Bol. Latinoam. Caribe Plantas Med. Aromát.* **2014**, *13*, 506–516.
98. Saha, A.; Chowdhury, K.K.; Bachar, S.C.; Roy, S.C.; Kundu, J.K. Anti-inflammatory, analgesic and diuretic activity of *Polygonum lanatum* Roxb. *Pak. J. Pharm. Sc.* **2005**, *18*, 13–18.
99. Smolarz, H.D.; Mendyk, E.; Bogucka-Kocka, A.; Kockic, J. Pinostrobin—an anti-leukemic flavonoid from *Polygonum lapathifolium* L. ssp. nodosum (Pers.) Dans. *Z. Naturforsch. C.* **2006**, *61*, 64–68, doi:10.1515/znc-2006-1-212.
100. Takasaki, M.; Konoshima, T.; Kuroki, S.; Tokuda, H.; Nishino, H. Cancer chemopreventive activity of phenylpropanoid esters of sucrose, vanicoside B and lapathoside A, from *Polygonum lapathifolium*. *Cancer Lett.* **2001**, *173*, 133–138, doi:10.1016/S0304-3835(01)00670-X.
101. Bulbul, L.; Sushanta, S.M.; Uddin, M.J.; Tanni, S. Phytochemical and pharmacological evaluations of *Polygonum lapathifolium* stem extract for anthelmintic and antiemetic activity. *Int. Curr. Pharm. J.* **2013**, *2*, 57–62, doi:10.3329/icpj.v2i3.13582.
102. Dzoyem, J.P.; Nkuete, A.H.; Kuete, V.; Tala, M.F.; Wabo, H.K.; Guru, S.K.; Rajput, V.S.; Sharma, A.; Tane, P.; Khan, I.A.; Saxena, A.K.; Laatsch, H.; Tan, N.H. Cytotoxicity and antimicrobial activity of the methanol extract and compounds from *Polygonum limbatum*. *Planta Med.* **2012**, *78*, 787–792, doi:10.1055/s-0031-1298431.
103. Kuete, V.; Nkuete, A.H.; Mbaveng, A.T.; Wiench, B.; Wabo, H.K.; Tane, P.; Efferth, T. Cytotoxicity and modes of action of 4'-hydroxy-2', 6'-dimethoxychalcone and other flavonoids toward drug-sensitive and multidrug-resistant cancer cell lines. *Phyto-medicine* **2014**, *21*, 1651–1657, doi:10.1016/j.phymed.2014.08.001.
104. Smolarz, H.D.; Kosikowska, U.; Baraniak, B.; Malm, A.; Persona, A. Lipophilicity, antifungal and antioxidant properties of persilben. *Acta Pol. Pharm.* **2005**, *62*, 457–460, doi:10.1016/j.bjm.2016.12.008.
105. Hussain, F.; Ahmad, B.; Hameed, I.; Dastagir, G.; Sanaullah, P.; Azam, S. Antibacterial, antifungal and insecticidal activities of some selected medicinal plants of Polygonaceae. *Afr. J. Biotechnol.* **2010**, *9*, 5032–5036.
106. Jovanović, M.; Morić, I.; Nikolić, B.; Pavić, A.; Svirčev, E.; Šenerović, L.; Mitić-Ćulafić, D. Anti-virulence potential and in vivo toxicity of *Persicaria maculosa* and *Bistorta officinalis* extracts. *Molecules* **2020**, *25*, 1811, doi:10.3390/molecules25081811.
107. Quesada-Romero, L.; Fernández-Galleguillos, C.; Bergmann, J.; Amorós, M.E.; Jiménez-Aspee, F.; González, A.; Simirgiotis, M.; Rossini, C. Phenolic fingerprinting, antioxidant, and deterrent potentials of *Persicaria maculosa* Extracts. *Molecules* **2020**, *25*, 3054, doi:10.3390/molecules25133054.
108. Hussein, S.R.; Mohamed, A.A. Antioxidant activity and phenolic profiling of two Egyptian medicinal herbs *Polygonum salicifolium* Brouss ex Wild and *Polygonum senegalense* Meisn. *An. Univ. din Oradea Fasc. Biol.* **2013**, *20*, 59–63.
109. Lajter, I.; Vasas, A.; Orvos, P.; Bánsághi, S.; Tólosi, L.; Jakab, G.; Béni, Z.; Háda, V.; Forgo, P.; Hohmann, J. Inhibition of G protein-activated inwardly rectifying K<sup>+</sup> channels by extracts of *Polygonum persicaria* and isolation of new flavonoids from the chloroform extract of the herb. *Planta Med.* **2013**, *79*, 1736–1741, doi:10.1055/s-0033-1351016.
110. Hassim, N.; Markom, M.; Anuar, N.; Baharum, S.N. Solvent selection in extraction of essential oil and bioactive compounds from *Polygonum minus*. *J. Appl. Sci.* **2014**, *14*, 1440–1444, doi:10.3923/jas.2014.1440.1444.

111. Uyub, A.M.; Nwachukwu, I.N.; Azlan, A.A.; Fariza, S.S. In-vitro antibacterial activity and cytotoxicity of selected medicinal plant extracts from Penang Island Malaysia on metronidazole-resistant-*Helicobacter pylori* and some pathogenic bacteria. *Ethnobot. Res. Appl.* **2010**, *8*, 95–106.
112. Abubakar, M.A.; Zulkifli, R.M.; Hassan, W.N.; Shariff, A.H.M.; Malek, N.Á.; Zakaria, Z.; Ahmad, F. Antibacterial properties of *Persicaria minor* (Huds.) ethanolic and aqueous-ethanolic leaf extracts. *J. Appl. Pharm. Sci.* **2015**, *5*, 50–56, doi:10.7324/JAPS.2015.58.S8.
113. Ahmad, R.; Baharum, S.N.; Bunawan, H.; Lee, M.; Mohd Noor, N.; Rohani, E.R.; Zin, N.M. Volatile profiling of aromatic traditional medicinal plant, *Polygonum minus* in different tissues and its biological activities. *Molecules* **2014**, *19*, 19220–19242, doi:10.3390/molecules191119220.
114. Qader, S.W.; Abdulla, M.A.; Chua, L.S.; Najim, N.; Zain, M.M.; Hamdan, S. Antioxidant, total phenolic content and cytotoxicity evaluation of selected Malaysian plants. *Molecules* **2011**, *16*, 3433–3443, doi:10.3390/molecules16043433.
115. George, A.; Chinnappan, S.; Chintamaneni, M.; Kotak, V.; Choudhary, Y.; Kueper, T.; Radhakrishnan, A.K. Anti-inflammatory effects of *Polygonum minus* (Huds) extract (Lineminus™) in in-vitro enzyme assays and carrageenan induced paw edema. *BMC Complement. Alter. Med.* **2014**, *14*, 335, doi:10.1186/1472-6882-14-355.
116. Rohin, M.A.K.; Abd Hadi, N.; Ridzwan, N. Different polarity Extracts of *Polygonum minus* towards cytotoxic activities against colon cancer cell lines (HT-29, HCT-116, CT-26). *J. Pharm. Res. Int.* **2020**, *32*, 168–180, doi:10.9734/jpri/2020/v32i1430617.
117. Ahmad, R.; Sahidin, I.; Taher, M.; Low, C.; Noor, N.M.; Sillapachaiyaporn, C.; Chuchawankul, S.; Sarachana, T.; Tencomnao, T.; Iskandar, F.; Rajab, N.F.; Baharum, S.N. Polygonumins A, a newly isolated compound from the stem of *Polygonum minus* Huds with potential medicinal activities. *Sci. Rep.* **2018**, *8*, 4202, doi:10.1038/s41598-018-22485-5.
118. Hamidi, J.A.; Ismaili, N.H.; Ahmadi, F.B.; Lajisi, N.H. Antiviral and cytotoxic activities of some plants used in Malaysian indigenous medicine. *Pertanika J. Trop. Agric. Sci.* **1996**, *19*, 129–136.
119. Christopher, P.V.; Joe, L.S.; Tian, M.; Mohan, T.S.B.; Parasuraman, S.; Al-suede, F.S.R.; Murugaiyah, V. Evaluation of methanol extract of *Polygonum minus* Huds. leaves for its hepatoprotective activity. *Malays. J. Microbiol.* **2016**, *12*, 345–352.
120. Basha, D.J.; Joy, J.M.; Kumar, A.S.; kumar Reddy, G.A.; Dalith, M.D. Hepatoprotective activity of root stocks of *Polygonum glabrum* Willd. Family Polygonaceae. *Int. J. Phytopharm. Res.* **2011**, *2*, 22–29.
121. George, A.; Ng, C.P.; O'Callaghan, M.; Jensen, G.S.; Wong, H.J. In vitro and ex-vivo cellular antioxidant protection and cognitive enhancing effects of an extract of *Polygonum minus* Huds (Lineminus™) demonstrated in a Barnes Maze animal model for memory and learning. *BMC Complement. Alter. Med.* **2014**, *14*, 161, doi:10.1186/1472-6882-14-161.
122. Wasman, S.Q.; Mahmood, A.A.; Salehuddin, H.; Zahra, A.A.; Salmah, I. Cytoprotective activities of *Polygonum minus* aqueous leaf extract on ethanol-induced gastric ulcer in rats. *J. Med. Plants Res.* **2010**, *4*, 2658–2665, doi:10.5897/JMPR09.412.
123. Qader, S.W.; Abdulla, M.A.; Chua, L.S.; Sirat, H.M.; Hamdan, S. Pharmacological mechanisms underlying gastroprotective activities of the fractions obtained from *Polygonum minus* in Sprague Dawley rats. *Int. J. Mol. Sci.* **2012**, *13*, 1481–1496, doi:10.3390/ijms13021481.
124. George, A.; Chinnappan, S.; Choudhary, Y.; Bommur, P.; Sridhar, M. Immunomodulatory activity of an aqueous extract of *Polygonum minus* Huds on Swiss albino mice using carbon clearance assay. *Asian Pac. J. Trop. Dis.* **2014**, *4*, 398–400, doi:10.1016/S2222-1808(14)60594-6.
125. Wei, Y.; Chen, X.; Jiang, X.; Ma, Z.; Xiao, J. Determination of taxifolin in *Polygonum orientale* and study on its antioxidant activity. *J. Food Compos. Anal.* **2009**, *22*, 154–157, doi:10.1016/j.jfca.2008.08.006.
126. Ansari, P.; Uddin, M.J.; Rahman, M.M.; Abdullah-Al-Mamun, M.; Islam, M.R.; Ali, M.H.; Reza, A.A. Anti-inflammatory, anti-diarrheal, thrombolytic and cytotoxic activities of an ornamental medicinal plant: *Persicaria orientalis*. *J. Basic Clin. Physiol. Pharm.* **2017**, *28*, 51–58, doi:10.1515/jbcpp-2016-0023.
127. Gou, K.J.; Zeng, R.; Dong, Y.; Hu, Q.Q.; Hu, H.W.Y.; Maffucci, K.G.; Dou, Q.L.; Yang, Q.B.; Qin, X.H.; Qu, Y. Anti-inflammatory and analgesic effects of *Polygonum orientale* L. extracts. *Front. Pharm.* **2017**, *8*, 562, doi:10.3389/fphar.2017.00562.
128. Tong, M.M.; Zhang, Y.Y.; Chu, Z.Y.; Zhai, Y.J. Research on antitumor effect in vitro of flowers of *Polygonum orientale* L. *Chinese J. Ethnomed. Ethnopharm.* **2013**, *24*, 7–8.
129. Xie, Z.T.; Tian, L.Q. Study on antitumor activity in vitro of *Polygonum orientale* L. total extraction and its chemical fractions. *Zhong Yi Xue Bao* **2012**, *27*, 1550–1551.
130. Nigam, V. Antihyperglycaemic activity on flower of *Polygonum orientale* Linn. using streptozotocin induced diabetic mice model. *Int. J. Pharm. Teach. Pract.* **2013**, *4*, 759–764.
131. Chiu, Y.J.; Chou, S.C.; Chiu, C.S.; Kao, C.P.; Wu, K.C.; Chen, C.J.; Tsai, J.C.; Peng, W.H. Hepatoprotective effect of the ethanol extract of *Polygonum orientale* on carbon tetrachloride-induced acute liver injury in mice. *J. Food Drug Anal.* **2018**, *26*, 369–379, doi:10.1016/j.jfda.2017.04.007.
132. Fan, X.; Bai, J.; Hu, M.; Xu, Y.; Zhao, S.; Sun, Y.; Wang, B.; Hu, J.; Li, Y. Drug interaction study of flavonoids toward OATP1B1 and their 3D structure activity relationship analysis for predicting hepatoprotective effects. *Toxicology* **2020**, *437*, 152,445, doi:10.1016/j.tox.2020.152445.
133. Qi, S.; Feng, Z.; Li, Q.; Qi, Z.; Zhang, Y. Inhibition of ROS-mediated activation Src-MAPK/AKT signaling by orientin alleviates H<sub>2</sub>O<sub>2</sub>-induced apoptosis in PC12 cells. *Drug Des. Devel. Ther.* **2018**, *12*, 3973–3984, doi:10.2147/DDDT.S178217.
134. Do, N.H.; Le, T.M.; Nguyen, C.D.; Ha, A.C. Optimization of total flavonoid content of ethanolic extract of *Persicaria pulchra* (Bl.) Soják for the inhibition of  $\alpha$ -glucosidase enzyme. *Fine Chem. Technol.* **2020**, *15*, 39–50, doi:10.32362/2410-6593-2020-15-4-39-50.

135. Nguyen, H.N.; Van Vo, G.; Tran, H.; Ho, H.T.C. Hepatoprotective effect of isolated constituent from *Polygonum tomentosum* against carbon tetrachloride induced toxicity. *Sci. Technol. Dev. J.* **2012**, *15*, 78–86, doi:10.32508/stdj.v15i2.1803.
136. Alves, T.M.D.A.; Ribeiro, F.L.; Kloos, H.; Zani, C.L. Polygodial, the fungitoxic component from the Brazilian medicinal plant *Polygonum punctatum*. *Mem. Inst. Oswaldo Cruz.* **2001**, *96*, 831–833, doi:10.1590/S0074-02762001000600016.
137. Penna, C.; Marino, S.; Vivot, E.; Cruaños, M.C.; Muñoz, J.D.D.; Cruaños, J.; Ferraro, G.; Martino, V. Antimicrobial activity of Argentine plants used in the treatment of infectious diseases. Isolation of active compounds from *Sebastiania brasiliensis*. *J. Ethnopharmacol.* **2001**, *77*, 37–40, doi:10.1016/S0378-8741(01)00266-5.
138. Lopez, A.; Hudson, J.B.; Towers, G.H.N. Antiviral and antimicrobial activities of Colombian medicinal plants. *J. Ethnopharmacol.* **2001**, *77*, 189–196, doi:10.1016/S0378-8741(01)00292-6.
139. Gilabert, M.; Cartagena, E.; Escobar, G.; Bardón, A.; Arena, M. Volatile terpenoids from water pepper (*Polygonum punctatum*) against *Pseudomonas aeruginosa* and *Staphylococcus aureus* virulence strategies. *Glob. J. Agric. Innov. Res. Dev.* **2014**, *1*, 3–10, doi:10.15377/2409-9813.2014.01.01.1.
140. Oliveira-Simoes, C.M.; Ribeiro-do-Vale, R.M.; Poli, A.; Nicolau, M.; Zanin, M. Pharmacological investigation on *Polygonum punctatum* Elliott (= *P. acre* HBK) extracts. *J. Pharm. Belg.* **1989**, *44*, 275–284.
141. Kott, V.; Barbini, L.; Cruanes, M.; de D Munoz, J.; Vivot, E.; Cruaños, J.; Martino, V.; Ferraro, G.; Cavallaro, L.; Campos, R. Antiviral activity in Argentine medicinal plants. *J. Ethnopharmacol.* **1998**, *64*, 79–84, doi:10.1016/S0378-8741(98)00098-1.
142. Sahidin, I.; Bahrún, A.; Taufik, M.; Mahatva Yodha, A.W.; Sabandar, C.W.; Imran, I.; Kadidae, L.O.; Darmawan, A.; Widodo, H.; Diantini, A. Chemical constituents of *Persicaria sagittata* (L.) H. Gross: Antioxidant activity and chemotaxonomy significance. *Jundishapur J. Nat. Pharm. Prod.* **2020**; *15*, e64788; doi:10.5812/jjnpp.64788.
143. Kenanda, E.O.; Omosa, L.K. Semi-Synthetic Pyrazoline Derivatives from *Polygonum senegalense* Chalcones and their Anti-Microbial Activities. *Pharmacog. Commn.* **2017**, *7*, 53–60, doi:10.5530/pc.2017.1.8.
144. Bothon, F.T.; Debiton, E.; Avlessi, F.; Forestier, C.; Teulade, J.C.; Sohounhloue, D.K. In vitro biological effects of two anti-diabetic medicinal plants used in Benin as folk medicine. *BMC Complement. Alter. Med.* **2013**, *13*, 51, doi:10.1186/1472-6882-13-51.
145. Mazid, M.A.; Datta, B.K.; Bachar, S.C.; Bashar, S.K.; Nahar, L., Sarker, S.D. Analgesic and anti-inflammatory activities of *Polygonum stagninum*. *Pharm. Biol.* **2010**, *48*, 770–774, doi:10.3109/13880200902991557.
146. Iwaki, K.; Koya-Miyata, S.; Kohno, K.; Ushio, S.; Fukuda, S. Antimicrobial activity of *Polygonum tinctorium* Lour: Extract against oral pathogenic bacteria. *J. Nat. Med.* **2006**, *60*, 121–125, doi:10.1007/s11418-005-0025-z.
147. Kataoka, M.; Hirata, K.; Kunikata, T.; Ushio, S.; Iwaki, K.; Ohashi, K.; Ikeda, M.; Kurimoto, M. Antibacterial action of tryptanthrin and kaempferol, isolated from the indigo plant (*Polygonum tinctorium* Lour.), against *Helicobacter pylori*-infected Mongolian gerbils. *J. Gastroenterol.* **2001**, *36*, 5–9, doi:10.1007/s005350170147.
148. Tokuyama-Nakai, S.; Kimura, H.; Ishihara, T.; Jisaka, M.; Yokota, K. In Vitro anti-inflammatory and antioxidant activities of 3,5,4'-trihydroxy-6,7-methylenedioxyflavone-O-glycosides and their aglycone from leaves of *Polygonum tinctorium* Lour. *Appl. Biochem. Biotechnol.* **2018**, *184*, 414–431, doi:10.1007/s12010-017-2555-8.
149. Kim, S.J.; Jang, T.W.; Kim, D.W.; Park, J.H. Study on antioxidant and anti-inflammatory activities of *Persicaria tinctoria*. *Korea J. Herbol.* **2015**, *30*, 17–24, doi:10.6116/kjh.2015.30.6.17.
150. Heo, B.G.; Park, Y.J.; Park, Y.S.; Bae, J.H.; Cho, J.Y.; Park, K.; Jastrzebski, Z.; Gorinstein, S. Anticancer and antioxidant effects of extracts from different parts of indigo plant. *Ind. Crops Prod.* **2014**, *56*, 9–16, doi:10.1016/j.indcrop.2014.02.023.
151. Lee, S.; Kim, D.C.; Baek, H.Y.; Lee, K.D.; Kim, Y.C.; Oh, H. Anti-neuroinflammatory effects of tryptanthrin from *Polygonum tinctorium* Lour. in lipopolysaccharide-stimulated BV2 microglial cells. *Arch. Pharm. Res.* **2018**, *41*, 419–430, doi:10.1007/s12272-018-1020-8.
152. Kimoto, T.; Hino, K.; Koya-Miyata, S.; Yamamoto, Y.; Takeuchi, M.; Nishizaki, Y.; Micallef, M.J.; Ushio, S.; Iwaki, K.; Ikeda, M.; Kurimoto, M. Cell differentiation and apoptosis of monocytic and promyelocytic leukemia cells (U-937 and HL-60) by tryptanthrin, an active ingredient of *Polygonum tinctorium* Lour. *Pathol. Int.* **2001**, *51*, 315–325, doi:10.1046/j.1440-1827.2001.01204.x.
153. Tatefuji, T.; Aga, M.; Kunikata, T.; Ikeda, M.; Kurimoto, M. Antiviral effect of *Polygonum tinctorium* Lour. extracts on virus-infected cells. *Nat. Med.* **1999**, *53*, 297–301.
154. Zhong, Y.; Yoshinaka, Y.; Takeda, T.; Shimizu, N.; Yoshizaki, S.; Inagaki, Y.; Matsuda, S.; Honda, G.; Fujii, N.; Yamamoto, N. Highly potent anti-HIV-1 activity isolated from fermented *Polygonum tinctorium* Aiton. *Antiviral Res.* **2005**, *66*, 119–128, doi:10.1016/j.antiviral.2005.02.003.
155. Kimura, H.; Tokuyama, S.; Ishihara, T.; Ogawa, S.; Yokota, K. Identification of new flavonol O-glycosides from indigo (*Polygonum tinctorium* Lour) leaves and their inhibitory activity against 3-hydroxy-3-methylglutaryl-CoA reductase. *J. Pharm. Biomed. Anal.* **2015**, *108*, 102–112, doi:10.1016/j.jpba.2015.02.005Get.
156. Cheng, H.W.; Lee, K.C.; Cheah, K.P.; Chang, M.L.; Lin, C.W.; Li, J.S.; Yu, W.Y.; Liu, E.T.; Hu, C.M. *Polygonum viviparum* L. inhibits the lipopolysaccharide-induced inflammatory response in RAW264. 7 macrophages through haem oxygenase-1 induction and activation of the Nrf2 pathway. *J. Sci. Food Agric.* **2013**, *93*, 491–497, doi:10.1002/jsfa.5795.
157. Ceylan, S.; Cetin, S.; Camadan, Y.; Saral, O.; Ozsen, O.; Tutus, A. Antibacterial and antioxidant activities of traditional medicinal plants from the Erzurum region of Turkey. *Ir. J. Med. Sci.* **2019**, *188*, 1303–1309, doi:10.1007/s11845-019-01993-x.
158. Salama, H.M.; Marraiki, N. Antimicrobial activity and phytochemical analyses of *Polygonum aviculare* L. (Polygonaceae), naturally growing in Egypt. *Saudi J. Biol. Sci.* **2010**, *17*, 57–63, doi:10.1016/j.sjbs.2009.12.009.

159. Zhang, L.; Ravipati, A.S.; Koyyalamudi, S.R.; Jeong, S.C.; Reddy, N.; Bartlett, J.; Smith, P.; de la Cruz, M.; Monteiro, M.C.; Melguizo, A.; Jiménez, E.; Vicente, F. Anti-fungal and anti-bacterial activities of ethanol extracts of selected traditional Chinese medicinal herbs. *Asian Pac. J. Trop. Med.* **2013**, *6*, 673–681, doi:10.1016/S1995-7645(13)60117-0.
160. Hsu, C.Y. Antioxidant activity of extract from *Polygonum aviculare* L. *Biol. Res.* **2006**, *39*, 281–288, doi:10.4067/S0716-97602006000200010.
161. Habibi, R.M.; Mohammadi, R.A.; Delazar, A.; Halabian, R.; Soleimani, R.J.; Mehdipour, A.; Bagheri, M.; Jahanian-Najafabadi, A. Effects of *Polygonum aviculare* herbal extract on proliferation and apoptotic gene expression of MCF-7. *J. of Ethnopharmacol.* **2011**, *19*, 326–331.
162. Mohammad, R.; Hossein, B.; Davood, F.; Farnaz, T.; Ali, F.; Yusef, R. The apoptotic and cytotoxic effects of *Polygonum aviculare* extract on Hela-S cervical cancer cell line. *Afr. J. Biochem. Res.* **2011**, *5*, 373–378, doi:10.5897/AJBR.9000230.
163. Cai, Y.; Wu, L.; Lin, X.; Hu, X.; Wang, L. Phenolic profiles and screening of potential  $\alpha$ -glucosidase inhibitors from *Polygonum aviculare* L. leaves using ultra-filtration combined with HPLC-ESI-qTOF-MS/MS and molecular docking analysis. *Ind. Crops Prod.* **2020**, *154*, 112,673, doi:10.1016/j.indcrop.2020.112673.
164. Won, J.B.; Ma, C.J. Neuroprotective activities of some medicinal plants against glutamate-induced neurotoxicity in primary cultures of rat cortical cells. *Nat. Prod. Sci.* **2009**, *15*, 125–129.
165. Zhang, F.X.; Xu, R.S. Juglanin ameliorates LPS-induced neuroinflammation in animal models of Parkinson's disease and cell culture via inactivating TLR4/NF- $\kappa$ B pathway. *Biomed. Pharmacother.* **2018**, *97*, 1011–1019, doi:10.1016/j.biopha.2017.08.132.
166. Park, S.H.; Jang, S.; Son, E.; Lee, S.W.; Park, S.D.; Sung, Y.Y.; Kim, H.K. *Polygonum aviculare* L. extract reduces fatigue by inhibiting neuroinflammation in restraint-stressed mice. *Phytomedicine* **2018**, *42*, 180–189, doi:10.1016/j.phymed.2018.03.042.
167. El-Kader, A.; Adel, M.; Nafady, A.M.; Ahmed, A.S.; Ibraheim, Z.Z. Antioxidant, hepatoprotective and antimicrobial activities of the aerial parts of *Polygonum bellardii* All. *Bull. Pharm. Sci. Assiut.* **2012**, *35*, 43–53, doi:10.21608/bfsa.2012.64600.
168. Abd El-kader, A.M.; El-Readi, M.Z.; Ahmed, A.S.; Nafady, A.M.; Wink, M.; Ibraheim, Z.Z. Polyphenols from aerial parts of *Polygonum bellardii* and their biological activities. *Pharm. Biol.* **2013**, *51*, 1026–1034, doi:10.3109/13880209.2013.775160.
169. Eruygur, N.; Ucar, E.; Atas, M.; Ergul, M.; Ergul, M.; Sozmen, F. Determination of biological activity of *Tragopogon porrifolius* and *Polygonum cognatum* consumed intensively by people in Sivas. *Toxicol. Rep.* **2020**, *7*, 59–66, doi:10.1016/j.toxrep.2019.12.002.
170. Pekdemir, M.E.; Pekdemir, S.; İnci, Ş.; Kirbağ, S.; Çiftci, M. Thermal, magnetic properties and antimicrobial effects of magnetic iron oxide nanoparticles treated with *Polygonum cognatum*. *Iran. J. Sci. Technol.* **2021**, doi:10.1007/s40995-021-01167-4.
171. Mahmoudi, M.; Boughalleb, F.; Mabrouk, M.; Tlili, N.; Potter, D.; Abdellaoui, R.; Nasri, N. Chemical analysis of the antioxidants from the aerial parts of wild *Polygonum equisetiforme* from Tunisia. *Food Biosci.* **2019**, *29*, 24–29, doi:10.1016/j.fbio.2019.03.004.
172. El-Toumy, S.A.H.; Salib, J.Y.; Shafik, N.H.; Elkarim, A.S.A.; Salama, A.; Omara, E.A.A.; Micky, J. Evaluation of hepatoprotective activity of *Polygonum equisetiforme* methanolic extract. *J. Appl. Pharm. Sci.* **2019**, *9*, 54–59, doi:10.7324/JAPS.2019.91107.
173. Zhang, C.F.; Hu, Y.; Lin, Y.; Huang, F.; Zhang, M. Anti-inflammatory activities of ethyl acetate extract of *Polygonum jucundum* and its phytochemical study. *J. Med. Plants Res.* **2012**, *6*, 1505–1511, doi:10.5897/JMPR10.671.
174. Jovanović, M.; Srdić-Rajčić, T.; Svirčev, E.; Jasnić, N.; Nikolić, B.; Bojić, S.; Stević, T.; Knežević-Vukčević J.; Mitić-Ćulafić, D. Evaluation of anticancer and antimicrobial activities of the *Polygonum maritimum* ethanol extract. *Arch. Biol. Sci.* **2018**, *70*, 665–673, doi:10.2298/ABS180423028J.
175. El-Haci, I.A.; Bekkara, F.A.; Mazari, W.; Hassani, F.; Didi, M.A. Screening of biological activities of *Polygonum maritimum* L. from Algerian coast. *Asian Pac. J. Trop. Biomed.* **2013**, *3*, 611–616, doi:10.1016/S2221-1691(13)60124-0.
176. Rodrigues, M.J.; Matkowski, A.; Ślusarczyk, S.; Magne, C.; Poleze, T.; Pereira, C.; Custodio, L. Sea knotgrass (*Polygonum maritimum* L.) as a potential source of innovative industrial products for skincare applications. *Ind. Crops Prod.* **2019**, *128*, 391–398, doi:10.1016/j.indcrop.2018.11.038.
177. Rodrigues, M.J.; Custódio, L.; Lopes, A.; Oliveira, M.; Neng, N.R.; Nogueira, J.M.; Martins, A.; Rauter, A.P.; Varela, J.; Barreira, L. Unlocking the in vitro anti-inflammatory and antidiabetic potential of *Polygonum maritimum*. *Pharm. Biol.* **2017**, *55*, 1348–1357, doi:10.1080/13880209.2017.1301493.
178. Rodrigues, M.J.; Varela, J.; Barreira, L.; Custódio, L. Antioxidant and neuroprotective potential of two halophytes from the Algarve coast. *Planta Med.* **2014**, *80*, P2Y16, doi:10.1055/s-0034-1394990.
179. Goswami, P.; Islam, R.; Khongmalai, E.; Sarkar, B.R.; Sen, S.; Choudhury, A.; Dey, B.K. Pharmacognostical, phytochemical and anthelmintic activity of *Polygonum muricatum*. *J. Appl. Pharm. Res.* **2018**, *6*, 07–12, doi:10.18231/2348-0335.2018.0009.
180. Li, W.; Zhang, X.; Wu, Y.; Tian, Z. Anti-inflammatory effects and mechanism of *Polygonum paleaceum* Wall. extracts. *Pharm. Clinics Chinese Mater. Med.* **2002**, *2*, 18–20.
181. Kim, S.Y.; Won, D.H.; Park, S.N. Antibacterial activity and component analysis of *Persicaria perfoliata* extracts. *Microbiol. Biotechnol. Lett.* **2010**, *38*, 278–282.
182. Fu, Y.; He, X.; Li, J.; Liu, X. Chemical composition Analysis and Antibacterial effect of *Polygonum perfoliatum*. *Prog. Vet. Med.* **2008**, *9*, 45–49.
183. Huang, X.; Jiang, L.; Wang, S.; Jiang, X.; Huang, D. Antibacterial activity research of medical plants of *Persicaria perfoliata*. *J. Mod. Med. Health.* **2013**, *29*, 3041–3042.
184. Fan, D.; Zhou, X.; Zhao, C.; Chen, H.; Zhao, Y.; Gong, X. Anti-inflammatory, antiviral and quantitative study of quercetin-3-O- $\beta$ -D-glucuronide in *Polygonum perfoliatum* L. *Fitoterapia* **2011**, *82*, 805–810, doi:10.1016/j.fitote.2011.04.007.
185. Tao, F.; Zhang, R.S. Research on anti-tumor effect of *Polygonum perfoliatum* L. in vitro and in vivo. *Chin. Arch. Tradit. Chin. Med.* **2013**, *31*, 2019–2021.

186. Li, Q.; Fu, X.; Ge, X.; Tao, F.; Huang, P.; Ge, M.; Jin, H. Antitumor effects and related mechanisms of ethyl acetate extracts of *Polygonum perfoliatum* L. *Front. Oncol.* **2019**, *9*, 578, doi:10.3389/fonc.2019.00578.
187. Zhang, Q.G.; Wei, F.; Liu, Q.; Chen, L.J.; Liu, Y.Y.; Luo, F.; Xiong, H.R.; Yang, Z.Q. The flavonoid from *Polygonum perfoliatum* L. inhibits herpes simplex virus 1 infection. *Acta Virol.* **2014**, *58*, 368–373, doi:10.4149/av\_2014\_04\_368.
188. Wang, Q. Preliminary study on Anti-HBV and hepatoprotective activity of *Polygonum perfoliatum*. Doctoral Thesis, Academy of Military Medical Sciences, Beijing, China, 2009.
189. Xu, L.; Huang, G.; Guo, X.; Zhou, Q.; He, S. Total flavonoids, extracted from *Polygonum knotweed* L, exert beneficial hepatoprotection against liver injury. *J. Cell. Biochem.* **2019**, *120*, 12677–12683, doi:10.1002/jcb.28535.
190. Rehman, A.; Waheed, A.; Tariq, R.; Zaman, M.; Tahir, M.J. Anti-fibrotic effects of *Polygonum plebeium* r. br. in CCl4-induced hepatic damage and fibrosis in rats. *Biomed. Res. Ther.* **2018**, *5*, 2223–2234, doi:10.15419/bmrat.v5i4.435.
191. Mazid, A.; Datta, B.K.; Nahar, L.; Rashid, A.; Bachar, S.C.; Bashar, S.K.; Sarker, S.D. Analgesic and diuretic properties of  $\alpha$ -santalone from *Polygonum flaccidum*. *Phytother. Res.* **2010**, *24*, 1084–1087, doi:10.1002/ptr.3053.
192. Lee, S.K.; Kwon, B.M.; Baek, N.I.; Kim, S.H.; Lee, J.H.; Park, H.W.; Kim, J.S.; Moon, M.K.; Kim, D.K.; Kim, D.K. Anti-proliferation effects of isorhamnetin isolated from *Persicaria thunbergii* on cancer cell lines. *Nat. Prod. Sci.* **2006**, *12*, 214–216.
193. Khan, H.; Saeed, M.; Khan, M.A.; Dar, A.; Khan, I. The antinociceptive activity of *Polygonatum verticillatum* rhizomes in pain models. *J. Ethnopharmacol.* **2010**, *127*, 521–527, doi:10.1016/j.jep.2009.10.003.
194. Amin, M.N.; Majumder, M.S.; Moghal, M.M.R.; Banik, S.; Kar, A.; Hossain, M.M. Anthelmintic and cytotoxic activities of two medicinal plants: *Polygonum viscosum* and *Aphanamixis polystachya* growing in Bangladesh. *J. Sci. Res.* **2014**, *6*, 339–345, doi:10.3329/jsr.v6i2.17299.
195. Datta, B.; Datta, S.; Khan, T.; Kundu, J.; Rashid, M.; Nahar, L.; Sarker, S. Anti-cholinergic, cytotoxic and anti-HIV-1 activities of sesquiterpenes and a flavonoid glycoside from the aerial parts of *Polygonum viscosum*. *Pharm. Biol.* **2004**, *42*, 18–23, doi:10.1080/13880200490504943.
196. Datta, B.K.; Datta, S.K.; Chowdhury, M.M.; Khan, T.H.; Kundu, J.K.; Rashid, M.A.; Nahar, L.; Sarker, S.D. Analgesic, antiinflammatory and CNS depressant activities of sesquiterpenes and a flavonoid glycoside from *Polygonum viscosum*. *Pharmazie* **2004**, *59*, 222–225.