

Health potential of Clery strawberries: enzymatic inhibition and anti-*Candida* activity evaluation

Francesco Cairone^{a,†}, Giovanna Simonetti^{b,†} Anastasia Orekhova^c, Maria Antonietta Casadei^a, Gokhan Zengin^d, Stefania Cesa^a

^aDipartimento di Chimica e Tecnologie del Farmaco, Università degli Studi di Roma “La Sapienza”,

Piazzale Aldo Moro 5, 00185 Roma, Italy

^bDipartimento di Biologia Ambientale, Università degli Studi di Roma “La Sapienza” P.le Aldo Moro 5, 00185 Rome, Italy

^cDipartimento di Sanità Pubblica e Malattie infettive, Università degli Studi di Roma “La Sapienza”

P.le Aldo Moro 5, 00185 Rome, Italy

^dDepartment of Biology, Science Faculty, Selcuk University, 42130 Konya, Turkey

[†] The authors contributed equally

*Corresponding author: Stefania Cesa

Address: Department of Drug Chemistry and Technology, Università degli Studi di Roma “LaSapienza”,

Piazzale Aldo Moro 5, 00185 Rome, Italy

Tel.: + 39 0649913198; fax: + 39 0649913133

E-mail address: stefania.cesa@uniroma1.it

Table S1. Colorimetric CIEL^aa^bb^c parameters of Clery graft on *Fragaria vesca* strawberry homogenates.

Reported results represent the mean value \pm SD from four experiments.

Sample	L*	a*	b*	C* _{ab}	h _{ab}
M graft	36.0 \pm 0.3	18.5 \pm 1.2	8.8 \pm 0.7	20.5 \pm 1.4	25.5 \pm 0.2
U graft	36.5 \pm 0.3	17.9 \pm 0.3	8.1 \pm 0.2	19.6 \pm 0.3	24.4 \pm 0.3
MP graft	37.6 \pm 0.4	18.5 \pm 0.8	8.3 \pm 0.3	20.3 \pm 0.9	24.0 \pm 0.4
UP graft	37.6 \pm 0.6	17.6 \pm 0.6	7.3 \pm 0.5	19.1 \pm 0.7	22.6 \pm 0.9
BM graft	44.0 \pm 0.2	22.4 \pm 0.6	9.9 \pm 0.9	24.5 \pm 0.9	23.8 \pm 1.7
BU graft	43.0 \pm 0.7	21.9 \pm 0.4	11.0 \pm 0.3	24.5 \pm 0.5	26.5 \pm 0.3
BMP graft	41.5 \pm 0.3	18.2 \pm 1.2	7.5 \pm 0.8	19.6 \pm 1.4	22.3 \pm 0.9
BUP graft	41.1 \pm 0.9	18.0 \pm 0.4	8.0 \pm 0.2	19.7 \pm 0.4	24.1 \pm 0.1
WM graft	35.9 \pm 0.3	18.8 \pm 0.4	8.7 \pm 0.8	20.8 \pm 0.8	24.7 \pm 1.6
WU graft	38.5 \pm 0.2	21.5 \pm 1.2	9.5 \pm 1.9	23.6 \pm 1.9	24.0 \pm 3.2
PM graft	37.2 \pm 2.2	20.0 \pm 0.1	9.4 \pm 0.3	22.1 \pm 0.2	25.0 \pm 0.7
PU graft	39.2 \pm 3.1	21.7 \pm 0.8	9.5 \pm 0.6	23.7 \pm 0.9	23.5 \pm 0.8

Table S2. HPLC-DAD quantitative analysis of Clery graft samples.

	M graft	U graft	MP graft	UP graft	BM graft	BU graft	BMP graft	BUP graft	WM graft	WU graft	PM graft	PU graft
Catechin	59.7	113.2	50.3	58.3	125.7	223.7	64.1	52.8	68.1	63.4	76.4	174.9
Epicatechin	277.9	300.8	402.2	268.6	365.2	374.4	257.8	197.4	154.2	285.3	197.7	217.5
p-Coumaric acid	12.9	13.2	13.6	13.6	nd	nd	4.5	8.9	7.2	18.7	8.6	5.7
Ferulic acid and derivatives*	209.3	242.5	222.2	166.9	145.7	219.8	109.1	82.7	85.5	97.9	167.4	170.3
Flavonols**	8.6	4.2	5.4	3.9	3.1	7.6	2.1	2.9	2.4	9.3	4.8	18.1
Anthocyanins***	91.8	78.4	70.9	65.9	128.1	99.1	41.8	63.1	37.5	77.8	183.6	190.1

Results are expressed in µg/g of fresh weight. The RSD value, evaluated on triplicates, was < 5%.

*Expressed as ferulic acid. ** Expressed as quercetin-3-D-galactoside. *** Expressed as

pelargonidin-3-glucoside. nd = not detected