

Supplementary material Table S1. Main constituents and percentage content of essential oils (EOs) from seven Greek native medicinal-aromatic plants (MAPs) tested *in vitro* against *Aeromonas veronii* bv. *sobria*. *GO: Greek oregano; *S: Savoury; R: Rosemary; WC: Wild carrot; *SO: Spanish oregano; P: Pennyroyal; L: Lemon balm (for the origin of the material see Table 4). The EOs content in bold from MAPs marked with asterisk (*) were evaluated as highly effective single-note EOs against the studied fish pathogen (from Anastasiou et al. 2020 with modifications).

No	Constituents	*GO	*S	R	WC	*SO	P	L
1	α -Thujene		2.5			3.5		
2	α -Pinene		1.8	8.8	20.5	1.6		
3	Camphene			4.0	1.1			
4	Sabinene				2.3			
5	β -Pinene		1.0	7.6				
6	1-Octen-3-ol					1.0		
7	Myrcene	1.2	2.9	1.1	6.9	4.6		
8	3-Octanol						2.0	
9	α -Terpinene		3.5			3.9		
10	p-Cymene	9.6	6.5	1.4		11.9		
11	Sylvestrene					1.0		
12	Limonene			1.9	7.5		1.2	
13	Eucalyptol			45.0				
14	<i>trans</i> -Ocimene						1.6	
15	γ -Terpinene	5.3	34.0			20.5		
16	Linalool		1.4			1.1		
17	Camphor			11.5				
18	Citronellal						10.2	
19	Menthone					3.5		
20	Isomenthone					5.2		
21	Borneol			4.3				
22	Terpinen-4-ol					1.0		
23	α -Terpineol			1.9				
24	Citronellol						1.8	
25	Pulegone					47.6		
26	Neral						6.5	
27	Carvacrolmethyleneether		1.8					
28	Piperitone					2.0		
29	Methylcitronellate						2.3	
30	Geranial						8.8	
31	Bornylacetate			1.9				
32	Thymol	2.1						
33	Carvacrol	72.0	32.8			42.0		
34	α -longipinene				5.2			
35	Piperitenone						33.0	
36	α -Copaene						2.4	
37	β -Bourbonene						1.5	

No	Constituents	*GO	*S	R	WC	*SO	P	L
38	β -Caryophyllene		6.9	3.7		3.4		27.7
39	α -Caryophyllene							2.2
40	<i>trans</i> - β -Farnesene							1.2
41	γ -Muurolene							12.6
42	α -Cedrene				3.7			
43	Isoeugenol methyl ether				14.8			
44	α -(<i>E,E</i>)-Farnesene							1.8
45	γ -Cadinene							1.6
46	δ -Cadinene							2.9
47	Caryophyllene oxide							2.7
48	β -Himachalene				21.6			
49	α -Cadinol							1.0