

In Vivo and In Vitro Biological Evaluation and Molecular Docking Studies of Compounds Isolated from *Micromeria biflora* (Buch. Ham. ex D.Don) Benth

Abdullah S. M. Aljohani ^{1,*}, Fahad A. Alhumaydhi ², Abdur Rauf ^{3,*}, Essam M. Hamad ⁴ and Umer Rashid ⁵

¹ Department of Veterinary Medicine, College of Agriculture and Veterinary Medicine, Qassim University, Buraydah 52571, Saudi Arabia

² Department of Medical Laboratories, College of Applied Medical Sciences, Qassim University, Buraydah 52571, Saudi Arabia; f.alhumaydhi@qu.edu.sa

³ Department of Chemistry, University of Swabi, Anbar23561, Khyber Pakhtunkhwa, Pakistan

⁴ Department of Food Science and Human Nutrition, College of Agriculture and Veterinary Medicine, Qassim University, Buraydah 52571, Saudi Arabia; e.hamad@qu.edu.sa

⁵ Department of Chemistry, COMSATS University Islamabad, Abbottabad Campus, 22060, Pakistan; umerrashid@cuiatd.edu.pk

* Correspondence: jhny@qu.edu.sa (A.S.M.A.); mashaljcs@yahoo.com (A.R.)




Figure S1. (a) The 2D interaction diagrams of isolated salicylalazine into the binding site of COX-1 (3N8Y) (b) COX-2 (1CX2).




Figure S2. (a) The 2D interaction diagrams of isolated salicylalazine into the binding site of GABA (4COF) (b) μ-opioid (5C1M).