

Review

Agarwood—The Fragrant Molecules of a Wounded Tree

Pooja Shivanand ^{1,*}, Nurul Fadhila Arbie ¹, Sarayu Krishnamoorthy ² and Norhayati Ahmad ^{1,3}

¹ Environmental and Life Sciences Program, Faculty of Science, Universiti Brunei Darussalam, Jalan Tungku Link, Bandar Seri Begawan BE1410, Brunei; 18b2114@ubd.edu.bn (N.F.A.); norhayati.ahmad@ubd.edu.bn (N.A.)

² Department of Civil Engineering, Environmental Water Resources Engineering Division, Indian Institute of Technology Madras, Chennai 600 036, India; sarayuk1528@gmail.com

³ Institute for Biodiversity and Environmental Research, Universiti Brunei Darussalam, Jalan Tungku Link, Bandar Seri Begawan BE1410, Brunei

* Correspondence: pooja.shivanand@ubd.edu.bn or poojashivanand@outlook.com; Tel.: +673-2460922-1378; Fax: +673-2461502

Abstract: Agarwood, popularly known as oudh or gaharu, is a fragrant resinous wood of high commercial value, traded worldwide and primarily used for its distinctive fragrance in incense, perfumes, and medicine. This fragrant wood is created when *Aquilaria* trees are wounded and infected by fungi, producing resin as a defense mechanism. The depletion of natural agarwood caused by overharvesting amidst increasing demand has caused this fragrant defensive resin of endangered *Aquilaria* to become a rare and valuable commodity. Given that instances of natural infection are quite low, artificial induction, including biological inoculation, is being conducted to induce agarwood formation. A long-term investigation could unravel insights contributing toward *Aquilaria* being sustainably cultivated. This review will look at the different methods of induction, including physical, chemical, and biological, and compare the production, yield, and quality of such treatments with naturally formed agarwood. Pharmaceutical properties and medicinal benefits of fragrance-associated compounds such as chromones and terpenoids are also discussed.

Keywords: agarwood; *Aquilaria*; artificial induction; bioactive compounds; chromones; terpenoids

Citation: Shivanand, P.; Arbie, N.F.; Krishnamoorthy, S.; Ahmad, N.

Agarwood—The Fragrant Molecules of a Wounded Tree. *Molecules* **2022**, *27*, 3386. <https://doi.org/10.3390/molecules27113386>

Academic Editor: Louis Pergaud Sandjo

Received: 29 April 2022

Accepted: 19 May 2022

Published: 24 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Agarwood, eaglewood, oud, aloeswood, and *gaharu* are some of the names of the highly valuable fragrant heartwood used for nontimber purposes and produced by the *Aquilaria* species, which belongs to the *Thymelaeaceae* family. These “woods of the Gods” have been used and traded for thousands of years for making perfume, which continue to be used in cultural and religious ceremonies, and for the production of incense sticks and fragrance products. It is also used in traditional medicine and has been recorded in the Ayurvedic medicinal text, the *Susruta Samhita* of prehistoric times, and in the *Sahih Muslim* that dates to the 8th century [1]. Agarwood oil is a desirable product and is traded worldwide, with high demand in Japan, China, the United Arab Emirates, and Saudi Arabia. It has a high demand in the Middle East because the oil symbolizes wealth, culture, and hospitality [2]. The value of agarwood depends on its quality, geographical location, and uses in a culture. Agarwood chips can cost from £20 per kilo up to £6000 per kilo depending on the amount of resin in the chips, while the agarwood oil can cost as much as £20,000 depending on the purity [3].

There are 21 *Aquilaria* species, with 9 agarwood (resin)-producing species [4,5], and they are found in India, New Guinea, Hainan Island in China, and in Southeast Asian countries [6]. Akter [7] stated that the formation of agarwood is associated with wounding of the tree and fungal invasion, causing the tree to produce a fragrant resin high in volatile compounds

that helps suppress the growth of fungi. Increased damage would eventually cause more resin to be produced.

Aquilaria

Due to its importance in culture, religion, and value, agarwood is highly sought after, and the trees are often overexploited and traded illegally, which has placed it on the list of endangered species in Appendix II of the Convention on International Trade in Endangered Species of Wild Flora and Fauna (CITES) in 1995 [1]. Furthermore, Chakrabarty et al. [8] reported that natural agarwood takes 20 years onward to be produced, and Sadgopal [9], as cited in [7], suggested that the best yields of resin are produced from trees aged 50 years and older. However, not all *Aquilaria* trees produce agarwood, and it was estimated that only 10% of wild *Aquilaria* spp. are infected and produce resin [10]. Due to this, it is difficult to meet the increasing demand for agarwood.

To prevent the extinction of *Aquilaria* trees and meet demand, some countries have taken the initiative to make *Aquilaria* plantations and artificially induce the production of agarwood in the trees. In Vietnam, a nongovernmental organization known as The Rainforest Project (TRP) developed field experiments for the artificial induction of agarwood. The experiment was performed by deliberately wounding the trees by drilling the trees and keeping the wound open with a small piece of plastic pipe, and injecting chemicals to stimulate the trees' defense mechanism to produce resin. This artificial induction was able to yield agarwood 10 times faster than natural formation and has been considered 'one of the most successful findings' [4]. This is supported by a recent study by Mohammad et al. [11], in which *A. beccariana* and *A. microcarpa* in Brunei Darussalam were inoculated with fungi isolated from various locations in the country and were able to produce agarwood within a month.

This review will analyze in-depth the different induction approaches (physical, chemical, and biological) and compare whether the production, yield, and quality of artificially induced agarwood varies from naturally formed agarwood. This will provide insight into how to prevent the overexploitation of *Aquilaria* trees in the wild while trying to meet the increasing demand for agarwood in the world. Additionally, fungal interactions with *Aquilaria* trees will also be explored in this paper to provide a better understanding of why resin is produced when the trees are stressed.

2. Natural Agarwood

Agarwood is formed when *Aquilaria* trees are wounded and exposed to biotic and abiotic stresses [12]. The infection triggers the trees' defense mechanism, causing resin to be produced, which aids the trees in suppressing the growth of the microbes infecting the trees in a process known as tylosis [7,12]. From the infection, the tree undergoes a biochemical reaction that produces oleoresin, which causes the color of the wood to eventually change from a lighter to a darker color, becoming what is commonly referred to as agarwood [12]. Wild-type agarwood takes years to produce, and few traders are willing to wait so long. Furthermore, only a small number of *Aquilaria* are infected in the wild and produce agarwood, and the only way to be certain that the tree contains the desired resin is to cut down the trees [4].

3. Artificially Induced Agarwood

Methods of artificial induction of agarwood have been created to prevent *Aquilaria* trees from becoming extinct. This causes the trees to become endangered, and, therefore, researchers have produced methods to artificially induce agarwood formation. There are three methods used: biological inoculation, chemical induction, and physical wounding. Table 1 summarizes the different induction mechanisms involved in the agarwood formation along with the details of its quality.

Table 1. Types of wounds inflicted on *Aquilaria* trees and the method to inflict these wounds for agarwood production. The quality and yield of agarwood are dependent on the method of wound infliction.

Wounding Type	Method	Quality and Yield	References
Natural	Lightning strike, animal grazing, insect attack, microbial invasion on wounded parts	High quality and low yield	[13,14]
Biological	Bacterial and fungal strains cultured on agar and injected into <i>Aquilaria</i> trees	Quality and yield depend on the strain used for the inoculation—can be either high or low quality and yield	[10,15,16]
Chemical	Whole-tree agarwood-inducing technique (Agar-wit), cultivated agarwood kit (CA-Kits), agarwood inducement method (AINM), jasmonic acid, acetic acid, sulfuric acid, alcohol, phytohormones, salts, minerals, biologically derived substances	Similar quality to natural agarwood and more yield compared with biological and physical wounding; however, both yield and quality depend on the chemicals used	[15,16,17,18,19]
Physical	Axe wounds, severe bark removal, nailing on tree trunk, partial trunk pruning method, burning-chisel-drilling method	Low quality and low yield	[1,13,16]

3.1. Physical Wounding

Mechanical injuries are the common and traditional method used to induce agarwood formation, as it is cheap and is inexpensive, requiring no chemicals or reagents to be used. It is also much easier to teach methods of mechanical injuries to farmers who cultivate agarwood. In China, farmers in Hainan, Guangdong, and Yunnan provinces were taught the physical wounding method to cultivate more than 20 million *A. sinensis* trees [15]. Ponjanagroon and Kaewrak [13] have used various methods of mechanical injuries on *A. crassna* to induce the production of agarwood; they inflicted wide and narrow wounds on the trees, made holes on the trees with screws of varied sizes, severe bark removal with hatchets, inserted nails of assorted sizes into the tree trunk, and the last one is to simply beat the *Aquilaria* trunk with a hammer. All methods produced discoloration; however, when the wood is burnt, the wood with nails hammered into the trunk gave no specific agarwood scent. Nobushi and Siripatanadilok [20] suggested that air and oxygen play a role in agarwood formation. Thus, when the nails are hammered into the trunk of the trees, oxygen is not able to enter the wound, and the discoloration around the wound could be caused by the reaction of ferric oxide in the nails and wood fibers [13]. Hence, there is no aromatic scent when burning the wood, as little or no resin was formed. The study also concluded that larger objects used to injure the trees cause wider discoloration, and the holes wounded with large screws were preferred, as it produces the classic agarwood scent when burnt and the quantity of agarwood at 20 months (about 1 and a half years) harvest was still not enough for commercial purposes [13].

3.2. Biological Inoculation

Biological inoculation is also another alternative method to agarwood formation and has been proven by many researchers to help induce agarwood formation. It is necessary for the tree to first be wounded before it can be infected by microbes to induce agarwood formation. However, not all fungi can promote agarwood production; some of the species identified in agarwood-producing trees are *Fusarium*, *Lasiodiplodia*, *Penicillium*, and *Aspergillus*, amongst others [12]. Inoculation of endophytic fungi on *Aquilaria* trees has also been proven to produce resin in as fast as 6 months [21]. Chen et al. [22] studied the agarwood formation induced by fermentation liquid of different fungi, in which the fungi were isolated from a previously infected tree that produced agarwood and were inserted into the *Aquilaria* tree by using a transfusion set. It was found that the dominant fungi were *Lasiodiplodia theobromae*, which was present in all layers of the wood, followed by *Fusarium solani* [22]. This suggests that *L.theobromae* and *F.solani* have a significant role in agarwood production and are agarwood-promoting fungi.

3.3. Chemical Induction

Chemical induction is another common method of producing agarwood in many countries. It is common to use sulfuric acid, jasmonic acid, acetic acid, and alcohol to induce agarwood formation, of which jasmonic acid has been proven to induce agarwood formation by 2–3 mm (about 0.12 in) thickness in Vietnam [17]. However, some countries have used sulfuric acid and acetic acid with unsuccessful results, and some chemicals are toxic to humans, hence the importance of choosing the proper chemicals when the agarwood is intended to be used for making perfumes, tea, and medicines [17]. Methods for injecting the chemicals in agarwood are similar in many reports, in which a hole is drilled into the trunk of the tree, and the chemicals are injected into the tree using a syringe or transfusion set [15,17,18]). There are kits and techniques made by researchers to induce agarwood production, such as the cultivated agarwood kits (CA-Kits) developed by Prof. Blanchette from the University of Minnesota, Vietnam, the whole-tree agarwood-inducing technique (Agar-Wit) that was developed in China [15]), and the agarwood inducement method (AINM) developed by Nuclear Malaysia, in which small holes of about 50 cm are drilled into the xylem of a tree followed by injection of agarwood inducers into the xylem. The resin can then be harvested after 6 months [18].

4. Biological Induction and Biosynthesis of Resin

Many endophytic fungal species have been reported to play a vital role in agarwood resin production in the *Aquilaria* and *Gyrinops* species [23]. Table 2 describes the various fungal endophytes known to induce agarwood formation in the *Aquilaria* and *Gyrinops* species. These fungal species could be inoculated into the trees either by natural (naturally occurring endophytes) or through artificial inoculation (through artificially created wounds or openings) methods and as pure or mixed cultures enabling the stimulation of the plants' immune response favoring resin production [24,25,26,27]. Biological induction is always considered the efficient method in resin formation, as it is safer, healthier, and ecofriendly compared with other methods and is a continuous process compared with the natural and physical induction methods [17]. Various sequential processes catalyzed by a set of specific enzymes in the plant cells enable the formation of the resinous materials, namely chromones and sesquiterpenoids in the plants.

Table 2. Resin-inducing endophytic fungal species isolated from the *Aquilaria* and *Gyrinops* species.

Fungal Endophyte	Specific Plant Species Reported for Resin Production	References
<i>Aspergillus</i> sp.	<i>Aquilaria</i> sp.	[28,29,30]
<i>Botryodiplodis</i> sp.		

<i>Botryosphaeria dothidea</i>		
<i>Diplodia</i> sp.		
<i>Fusarium bulbigenium</i>		
<i>Fusarium lateritium</i>		
<i>Fusarium oxysporum</i>		
<i>Fusarium</i> sp.		
<i>Fusarium moniliforme</i>		
<i>Fusarium sambucinum</i>		
<i>Fusarium solani</i>		
<i>Fusarium tricinctum</i>		
<i>Epicoccum granulatum</i>		
<i>Cladosporium</i> sp.		
<i>Torula</i> asp.	<i>Aquilaria agallocha</i>	[9,31,32,33,34]
<i>Chaetomium globosum</i>		
<i>Fusarium oxysporum</i>		
<i>Melanotus flavolivens</i>		
<i>Lasiodiplodia</i> sp.		
<i>Xylaria</i> sp.		
<i>Paraconiothyrium variabile</i>		
<i>Botryosphaeria</i> sp.		
<i>Fusarium</i> sp.	<i>Aquilaria sinensis</i>	[35,36,37,38,39,40,41,42,43,44,45,46]
<i>Lasiodiplodia theobromae</i>		
<i>Fusarium oxysporum</i>		
<i>Rigidoporus vinctus</i>		
<i>Nigrospora oryzae</i>		
<i>Fusarium solani</i>		
<i>Acremonium</i> sp.	<i>Aquilaria microcarpa</i>	[47,48]
<i>Fusarium</i> sp.		
Unidentified <i>Deuteromycetes</i>		
Unidentified <i>Ascomycetes</i>		
<i>Fusarium</i> sp.	<i>Aquilaria malaccensis</i>	[49,50,51,52,53]
<i>Fusarium solani</i>		
<i>Acremonium</i> sp.		
<i>Fusarium</i> sp.	<i>Aquilaria crassna</i>	[54]
<i>Fusarium solani</i>		
<i>Lasiodiplodia theobromae</i>		
<i>Fusarium</i> sp.	<i>Aquilaria beccariana</i>	[55]
<i>Fusarium</i> sp.	<i>Gyrinops versteegii</i>	[30,53,56]
<i>Fusarium solani</i>		
<i>Aspergillus niger</i>	<i>Gyrinops walla</i>	[57]
<i>Fusarium solani</i>		

The 2-(2-phenylethyl) chromones (PECs) and sesquiterpenoids are the major secondary metabolites synthesized by the trees along with triterpenes and sterols due to the stress developed by the natural and artificial induction methods on the immune system [58]. Chromones with a remarkably high and wide range of medicinal and therapeutic values [59,60,61] are derived from benzopyrans (polycyclic organic compound) with a keto group in the oxime ring. Biosynthesis of the chromones is initiated through multiple mechanisms involving the pentaketide pathway, shikimic acid pathway, and addition of nitrogenous groups from amino acids to the chromones [59,62].

Recent studies by Wang et al. [63] have shown that the stress caused by elevated levels of salinity has improved resin production in the *Aquilaria* tree. Similarly, according to Liao et al. [64], flindersia-type 2-(2-phenylethyl) chromones, an important and major constituent of the resin, were found to be formed by the catalysis of III polyketide synthase (PKs) through condensation of dihydro-cinnamoyl-CoA analogs and malonyl-CoA with 2-hydroxy-benzoyl-CoA, and on subsequent catalysis with hydroxylases or O-methyltransferases (OMTs). As PECs are synthesized using a complex phenomenon involving a sequence of processes in the plants, it is still a challenging task for researchers worldwide to elucidate the exact pathway involved in the synthesis of PECs. Researchers Goel and Makrandi [65] and Tawfik et al. [61] have stated that the known pathways of PECs synthesis to date were found to be incomplete, with inadequate information lacking the proper understanding of the specific linkages in the pathway of the PECs synthesis. Conversely, sesquiterpenes, triterpenes, and sterols are synthesized using the isoprenoid precursor with the mevalonic acid pathway in the cytosol of the plant cells, and the methylerythritol phosphate pathway acts as a precursor that helps in the synthesis of monoterpenes, diterpenes, and carotenoids in the plastids [66,67,68]. Both the pathways discussed above initiate the synthesis of C5 homoallylic isoprenoid precursors such as isopentenyl pyrophosphate and dimethylallyl pyrophosphate (which are found to be exchanged among the two pathways in the space between the cytosol and plastids), engaging pyruvate, acetyl-CoA, and various other enzymes [67]. These C5 homoallylic isoprenoid precursors on sequential condensation in the presence of C15 farnesyl pyrophosphate synthase generates C15 farnesyl pyrophosphate [66,69]. Researchers such as Yang et al. [69] and Liu et al. [70] have stated that C15 farnesyl pyrophosphate synthase acts as the key limiting factor in the synthesis of sesquiterpenes that are encoded by the gene *Am-FaPS-1* and *AsFPS1* in *Aquilaria* species. Another set of enzymes, namely sesquiterpene synthase and cytochrome P450 dependent mono-oxygenases, play a pervasive role in the final stage of the sesquiterpene synthesis by oxidative functionalization of the C15 farnesyl pyrophosphate with the cytochrome P450 dependent mono-oxygenases, thus forming a multicyclic scaffold complex. These scaffolds are further modified by the addition of functional groups by alkylation, esterification, and the addition of sugar residues to the hydroxyl end generated by the cytochrome P450-dependent mono-oxygenases in the scaffolds, thus forming different known sesquiterpenoids [71,72]. Further, many studies are underway to better understand the synthesis pathway of the sesquiterpenoids and the role of different genes involved in their synthesis and the synthesis of various synthases involved in the process.

5. Agarwood Quality

The market value of agarwood is determined by its quality. Agarwood quality grading is often supervised by trained human graders who base the quality on color, odor, high fixative, and consumer perception [2]. Different countries have their own ways of labeling agarwood quality; some countries prefer to use the terms 'high quality' and 'low quality,' and other countries prefer to categorize agarwood into groups A, B, C, or D [73]. However, it is subjective when agarwood quality is determined by humans, in which each person might have different opinions on the physical appearance of agarwood. In addition, humans are prone to fatigue and nausea when exposed to many fragrances for a prolonged period, and this would limit the analysis and may cause wrong judgment to be made. The agarwood compound is associated with the presence of sesquiterpenes, which are the main active compounds in agarwood that gives it the fragrant odor [74]. Thus, the more sesquiterpenes the agarwood contains, the higher the quality of the oil would be.

The three methods of artificial induction come with their own advantages and disadvantages. The physical-mechanical method is by far the most common method and is also known as the traditional method. According to researchers [13,16], this method produces low-grade and low-yield agarwood and often takes a long time to produce

agarwood. However, this method is common, as it is easy to teach farmers who cultivate agarwood and is cheap cost-wise. As for the biological method, the quality and yield depend on the strains of fungi or bacteria used during the inoculation process. Some studies have mentioned *Aspergillus* sp., *L. theobromae*, and *Fusarium* sp. [12,22] as playing a significant role in the biological inoculation of agarwood. The quality of agarwood is then determined by analyzing the sesquiterpenes compounds that are present. As for the production time, agarwood can be produced in as early as 6 months [21], but the yield of agarwood from this method needs to be further evaluated when using different strains.

In terms of chemical induction, Chong et al. [18] harvested agarwood from trees that were 18 months (about 1 and a half years) old, in which some of the trees were inoculated with fungi, and the others were injected with chemical inducers using AINM. Their results showed that the yield of agarwood using chemical inducers was 12.9 times higher than using fungal strains [18]. Another study by Liu et al. [15] also showed that agarwood collected from the Agar-Wit method after 6 months was also superior compared with physical wounding and biological inoculation. High-quality agarwood had a high alcohol soluble extractive content, and the samples collected using the Agar-Wit method had an alcohol soluble content surpassing the 10% standard, which was similar to the agarwood obtained from wild *Aquilaria* trees [15]. Furthermore, when the harvest occurred after 12 or 20 months (about 1 and a half years), the agarwood produced was an even higher quality [15]. Additionally, in these studies, the quality of agarwood oil and the yield using chemical inducers compared with other methods had high levels of the sesquiterpenes compound, which shows that chemical induction is the best method to produce agarwood that has similar qualities to wild agarwood.

6. Bioactive Compounds Obtained from Agarwood, Their Pharmaceutical Properties, and Medicinal Benefits

Agarwood and its products, either as oil, smoke, or powder admixtures, are well known for their bioactivity in controlling various fungal pathogens and their unique medicinal properties globally [75]. Several chemical compounds have been reported to be identified from agarwood such as chromone derivatives, terpenoids, flavonoids, benzophenones, lignans, benzenoid derivatives, phenolic compounds, triterpenes, steroids, and other chemical compounds [58,76]. Of these, chromones and terpenoids (sesquiterpenoids) are the compounds of interest that are potentially known for their bioactivity, pharmaceutical value, and medicinal properties [76]. In contrast, the other chemical compounds are natural compounds observed in most plants and trees [77].

6.1. Chromones

More than 80 different chromones (2-(2-phenylethyl) chromones) and about 31 different terpenoids are known to date (Table 3) that are responsible for various medicinal benefits [78]. Chromones (1,4-benzopyrone) are the known isomer of coumarin, which are derivatives of benzopyran with a substituted keto group in the pyran ring, are ubiquitously present in the Plant Kingdom, and are a part of the human and animal diet [79]. The 2-(2-phenylethyl) chromones are the rare and uncommon group of chromones that possess phenylethylene at the C-2 position, which are reported to be abundantly available in agarwood resin (Table 3) [80,81]. These 2-(2-phenylethyl) chromones have been isolated only from a few plant species, making it a rare compound extracted from plants such as agarwood that are responsible for the warm, balsamic, long-lastingly sweet odor of the burnt agarwood [82]. Furthermore, they are classified into four subgroups, namely 2-(2-Phenylethyl)chromone monomers (PEC), comprising four subdivisions (Flindersia-type 2-(2-phenylethyl)chromones, the most abundant group; 5,6,7,8-Tetrahydro-2-(2-phenylethyl)chromones, highly oxidizing group; mono-epoxy-5,6,7,8-tetrahydro-2-(2-phenylethyl)chromones; diepoxy-5,6,7,8-tetrahydro-2-(2-phenylethyl)chromones), 2-(2-Phenylethyl)chromones are predominantly obtained from chemical synthesis, agarwood, cyanobacteria, and rhizomes of *Imperata cylindrica* or

Platanus x acerifolia; dimeric 2-(2-phenylethyl)chromones; sesquiterpenoid-4H-chromones and benzylacetone-4H-chromones, predominantly obtained from *Gyrinops salicifolia*; and trimeric 2-(2-phenethyl)chromones [64,83,84,85,86,87]. They act as potential and remarkable pharmacological compounds containing various bioactivities such as antimicrobial, antiviral, anticancer, antitumor, anti-inflammatory, antioxidant, enzyme inhibition, antifeedant, antidepressant, antiobesity, and antihypersensitive properties, including antagonistic activity in melanin-concentrating hormone receptor-1 [88,89,90].

Table 3. List of chromones and terpenoids known to be extracted from agarwood obtained from *Aquilaria sinensis*, *A. malaccensis*, *A. filaria*, and *Gyrinops versteegii* (as reported by Wang et al. [91] and Yuan et al. [76]).

Chemical Compounds	Bioactive Properties	References
Chromones		
6-hydroxy-7-methoxy-2-[2-(3'-hydroxy-4'-methoxyphenyl)ethyl]chromone		[60,92,93]
6,7-dimethoxy-2-[2-(3'-hydroxy-4'-methoxyphenyl)ethyl]chromone		[60,93]
7-hydroxy-6-methoxy-2-[2-(3'-hydroxy-4'-methoxyphenyl)ethyl]chromone		[60]
6,7-dimethoxy-2-[2-(4'-hydroxy-3'-methoxyphenyl)ethyl]chromone		[60,93]
6,7-dihydroxy-2-[2-(4'-methoxyphenyl)ethyl]chromone		[60]
6-hydroxy-7-methoxy-2-[2-(4'-hydroxyphenyl)ethyl]chromone		[60]
6,8-dihydroxy-2-[2-(3'-hydroxy-4'-methoxyphenyl)ethyl]chromone		[60,92,93]
6-methoxy-2-[2-(3-hydroxy-4-methoxyphenyl)ethyl]chromone		[80]
5-hydroxy-6-methoxy-2-[2-(4-methoxyphenyl)ethyl]chromone	Antioxidant, analgesic, digestive, tumor cell inhibition, enzyme inhibition, antimicrobial,	[80,94]
6-methoxy-2-[2-(4-methoxyphenyl)ethyl]chromone	antiplasmodial, antifeedant,	[80]
6-methoxy-2-[2-(4-hydroxyphenyl)ethyl]chromone	immunomodulatory, anti-inflammatory, antitubercullii	[80,92]
6-methoxy-2-[2-(3-methoxy-4-hydroxyphenyl)ethyl]chromone		[80,95,96]
6,7-dimethoxy-2-[2-phenylethyl]chromone	Treatment of gastritis, diarrhea,	[80,94,97]
6-hydroxy-2-(2-phenylethyl)chromone	stiff muscles, hypothermic disease,	[80,94]
6-hydroxy-2-[2-(4-methoxyphenyl)ethyl]chromone	diuretic disease, CNS activity,	[80]
6,8-dihydroxy-2-(2-phenylethyl)chromone	nephritis, cystitis, urethral disease,	[80]
5-hydroxy-6-methoxy-2-[2-(3-hydroxy-4-methoxyphenyl)ethyl]chromone	pyrexia, rheumatism, headache, hepatitis, cough, bronchitis, asthma	[80]
5-hydroxy-6-methoxy-2-[2-(4-methoxyphenyl)ethyl]chromone		[80]
6-hydroxy-7-methoxy-2-[2-(4-methoxyphenyl)ethyl]chromone		[97]
6-methoxy-7-hydroxy-2-[2-(4-methoxyphenyl)ethyl]chromone		[97]
6-hydroxy-2-[2-(3-methoxy-4-hydroxyphenyl)ethyl]chromone		[97]
6-hydroxy-2-[2-(3-hydroxy-4-methoxyphenyl)ethyl]chromone		[92,97]
6-hydroxy-2-[2-(3,4-dimethoxyphenyl)ethyl]chromone		[97]
6,8-dihydroxy-2-[2-(4-methoxyphenyl)ethyl]chromone		[97]
8-chloro-6-hydroxy-2-[2-(3-methoxy-4-hydroxyphenyl)ethyl]chromone		[97]
5-methoxy-6-hydroxy-2-[2-(3-methoxy-4-hydroxyphenyl)ethyl]chromone		[97]
4',6-dihydroxy-3',7-dimethoxy-2-(2-phenyl)ethylchromone		[92]

6-methoxy-2-[2-(3-methoxy-4-hydroxyphenyl)ethyl]chromone	[95]
6-hydroxy-2-[2-(4-hydroxyphenyl)ethyl]chromone	[95]
6,8-dihydroxy-2-[2-(3-methoxy-4-hydroxyphenyl)ethyl]chromone	[96]
7-hydroxy-6-methoxy-2-(2-phenylethyl)chromone	[95]
6,7-dimethoxy-2-(2-phenylethyl)chromone	[95]
6-hydroxy-2-[2-(2-hydroxyphenyl)ethyl]chromone	[98]
7-hydroxy-2-(2-phenylethyl)chromone	[98]
flindersiachromone	[94]
6-methoxy-2-(2-phenylethyl)chromone	[94]
5-hydroxy-6,7-dimethoxy-2-[2-(4'-methoxyphenyl)ethyl]chromone	[81]
8-chloro-6-hydroxy-2-(2-phenylethyl)chromen-4-one	[99]
8-chloro-6-hydroxy-2-[2-(4-methoxyphenyl)-ethyl]chromen-4-one	[99]
aquilarinoside C	[100]
6-hydroxy-2-[2-(40-hydroxy-30-methoxyphenyl)ethenyl]chromone	[60]
(5S,6S,7S,8R)-2-[2-(3-hydroxy-4-methoxyphenyl)ethyl]-5,6,7,8-tetrahydroxy-5,6,7,8-tetrahydrochromone	[92]
(5S,6S,7S,8R)-2-(2-phenylethyl)-5,6,7,8-tetrahydroxy-5,6,7,8-tetrahydrochromone	[92]
(5S,6S,7S,8R)-2-[2-(4-methoxyphenyl)ethyl]-5,6,7,8-tetrahydroxy-5,6,7,8-tetrahydrochromone	[92]
(5S,6R,7S,8R)-2-[2-(3-hydroxy-4-methoxyphenyl)ethyl]-5,6,7,8-tetrahydroxy-5,6,7,8-tetrahydrochromone	[92]
(5S,6R,7R,8S)-2-[2-(3-hydroxy-4-methoxyphenyl)ethyl]-5,6,7,8-tetrahydroxy-5,6,7,8-tetrahydrochromone	[92]
(5S,6R,7R,8S)-2-[2-(4-hydroxyphenyl)ethyl]-5,6,7,8-tetrahydroxy-5,6,7,8-tetrahydrochromone	[92]
8-chloro-2-(2-phenylethyl)-5,6,7-trihydroxy-5,6,7,8-tetrahydrochromone	[92]
6S,7R)-5,6,7,8-Tetrahydro-6,7-dihydroxy-2-(2-phenylethyl)-4H-1-benzopyran-4-one	[94]
aquilarone F	[94]
(5R,6R,7R,8S)-8-chloro-5,6,7-trihydroxy-2-(4-methoxyphenethyl)-5,6,7,8-tetrahydrochromone	[81]
(5S,6S,7S,8S)-8-chloro-5,6,7-trihydroxy-2-(2-phenylethyl)-5,6,7,8-tetrahydrochromone	[81]
(5R,6R,7R,8R)-8-chloro-5,6,7-trihydroxy-2-(4-methoxyphenethyl)-5,6,7,8-tetrahydrochromone	[81]
(5R,6S,7S)-5,6,7-trihydroxy-2-(4-hydroxy-3-methoxyphenethyl)-5,6,7,8-tetrahydrochromone	[81]
tetrahydrochromone A	[101]
tetrahydrochromone B	[101]
tetrahydrochromone C	[101]
tetrahydrochromone D	[101]
tetrahydrochromone E	[101]
tetrahydrochromone F	[101]
tetrahydrochromone G	[101]
tetrahydrochromone H	[101]
tetrahydrochromone I	[101]
tetrahydrochromone J	[101]

(5R,6R,7S,8R)-5,6,7,8-tetrahydroxy-2-(2-phenylethyl)-5,6,7,8-tetrahydrochromone	[84]
agarotetrol	[84]
qinanmer	[84]
5,6-epoxy-7 β -hydroxy-8 β -methoxy-2-(2-phenylethyl)chromone	[80]
rel-(1aR,2R,3R,7bS)-1a,2,3,7b-tetrahydro-2,3-dihydroxy-5-(2-phenylethyl)-7H-oxireno-benzopyran-7-one	[80]
rel-(1aR,2R,3R,7bS)-1a,2,3,7b-tetrahydro-2,3-dihydroxy-5-[2-(4-methoxyphenyl)ethyl]-7H-oxireno-benzopyran-7-one	[94]
oxidoagarochromone A	[80]
tetrahydrochromone L	[101]
tetrahydrochromone M	[101]
oxidoagarochromone A	[80]
oxidoagarochromone B	[80]
oxidoagarochromone C	[101]
(R)-6,7-dimethoxy-2-(2-hydroxy-2-phenylethyl)chromone	[93]
(S)-6,7-dimethoxy-2-(2-hydroxy-2-phenylethyl)chromone	[93]
AH10	[94]
AH14	[94]
(5S,6R,7S,8R)-2-[2-(4-Methoxyphenyl)ethyl]-5,6,7-trihydroxy-5,6,7,8-tetrahydro-8-{6-methoxy-2-[2-(3 ^{'''} -methoxy-4 ^{'''} -hydroxyphenyl)ethyl]chromonyl-7-oxy}chromone	[102]
(5S,6R,7S,8R)-2-[2-(4-Methoxyphenyl)ethyl]-5,6,7-trihydroxy-5,6,7,8-tetrahydro-8-{2-[2-(4 ^{'''} -methoxyphenyl)ethyl]chromonyl-6-oxy}chromone	[102]
(5S,6R,7S,8R)-2-(2-Phenylethyl)-5,6,7-trihydroxy-5,6,7,8-tetrahydro-8-[2-(2-phenylethyl)chromonyl-6-oxy]chromone	[102]
(5R,6R,7R,8S)-2-(2-Phenylethyl)-5,6,7-trihydroxy-5,6,7,8-tetrahydro-8-[2-(2-phenylethyl)chromonyl-6-oxy]chromone	[102]
aquisinenone A	[81]
(-)-4'-methoxyaquisinenone A	[81]
aquisinenone B	[81]
(-)-6''-hydroxyaquisinenone B	[81]
(+)-6''-hydroxy-4',4'''-dimethoxyaquisinenone B	[81]
aquisinenone C	[81]
(-)-aquisinenone D	[81]
4'-demethoxyaquisinenone D	[81]
(+)-aquisinenone E	[81]
(-)-aquisinenone F	[81]
(-)-aquisinenone G	[81]
(+)-4'-methoxyaquisinenone G	[81]
Terpenoids	
(+)-9 β -Hydroxyeudesma-4,11(13)-dien-12-al	[103]
(+)-Eudesma-4,11(13)-dien-8 α ,9 β -diol	[103]
(+)-8 α -Hydroxyeudesma-3,11(13)-dien-14-al	[103]
(+)-Eudesma-3,11(13)-dien-8 α ,9 β -diol	Antimicrobial, anti-inflammatory, [103]
(+)-Eudesma-4(14),11(13)-dien-8 α ,9 β -diol	anticancer, antidiabetes, [103]
(4R,5R,7S,9S,10S)-(-)-Eudesma-11(13)-en-4,9-diol	antioxidants [103]
(+)-9 β ,10 β -Epoxyeremophila-11(13)-en	Treatment of immunological [103]
(+)-11-Hydroxyvalenc-1(10),8-dien-2-one	disorders, neurological disorders, [103]
(-)-Eremophila-9-en-8 β ,11-diol	cancer [103]
1,10-Dioxo-4H-5H-7H-11H-1,10-secoguaia-2(3)-en-12,8-olide	[104]
1-Hydroxy-4H-5H-7H-11H-8,9-secoguaia-9(10)-en-8,12-olide	[104]
1-Hydroxy-4 α ,10 α -dimethyl-5H-octahydro-azulen-8-one	[104]

1 α -Hydroxy-4 α ,10 α -dimethyl-5 β H-octahydro-azulen-8-one	[104]
4-Hydroxyl-baimuxinol	[105]
7 β -H-9(10)-ene-11,12-Epoxy-8-oxoeremophilane	[105]
7 α -H-9(10)-ene-11,12-Epoxy-8-oxoeremophilane	[105]
(5S,7S,9S,10S)-(+)-9-Hydroxy-selina-3,11-dien-12-al	[106]
(5S,7S,9S,10S)-(-)-9-Hydroxy-selina-3,11-dien-14-al	[106]
(5S,7S,9S,10S)-(+)-9-Hydroxy-eudesma-3,11(13)-dien-12-methyl ester	[106]
(7S,9S,10S)-(+)-9-Hydroxy-selina-4,11-dien-14-al	[106]
(7S,8S,10S)-(+)-8,12-Dihydroxy-selina-4,11-dien-14-al	[106]
Qinanol A	[107]
Qinanol B	[107]
Qinanol C	[107]
Qinanol D	[107]
Qinanol E	[107]
Qinanol F	[107]
3-oxo-7-Hydroxylholosericin A	[97]
1,5,8,12-Diepoxy-guaia-12-one	[97]
(+)-8 β -Hydroxy-longicamphenylone	[60]
11 β -Hydroxy-13-isopropyl-dihydrodehydrocostus lactone	[60]
Aquilarabietic acid A	[69]
Aquilarabietic acid B	[69]
Aquilarabietic acid C	[69]
Aquilarabietic acid D	[69]
Aquilarabietic acid E	[69]
Aquilarabietic acid F	[69]
Aquilarabietic acid G	[69]
Aquilarabietic acid H	[69]
Aquilarabietic acid I	[69]
Aquilarabietic acid J	[69]
Aquilarabietic acid K	[69]
Aquilarin B	[108]
Aquilanol A	[109]
Aquilanol B	[109]
Daphnauranol D	[109]
Chamaejasmane E	[109]
Aquilacallane A	[110]
Aquilacallane B	[110]
Aquimavitalin	[111]
12-O-(20E,40E)-6-oxohexa-20, 40-Dienoylphorbol-13-acetate	[112]
12-Deoxy-13-O-acetylphorbol-20-(90Z)-octadecenoate	[112]
12-O-(20E,40E)-60, 70-(erythro)-dihydroxytetradeca-20, 40-dienoylphorbol-13-acetate	[112]
12-O-(20E,40E)-60, 70-(threo)-dihydroxytetradeca-20, 40-dienoylphorbol-13-acetate	[112]

Furthermore, chromones are considered naturally available products with diverse structures and functions that act as potential candidates for replacing synthetic drugs in various pharmacological therapeutics [89,113,114,115,116,117,118,119,120,121,122,123,124,125,126]. Similarly, Duan et al. [127] and Vanguru et al. [128] have reported the therapeutic properties of chromones against a wide range of cancers, especially in controlling breast cancer and ovarian cancer [129,130]. They act as intercellular adhesion molecule inhibitors, cyclooxygenase inhibitors, mast cell stabilizers, leukotriene receptor antagonists, interleukin-5 inhibitors, lipoxygenase inhibitors, and nitric oxide production inhibitors controlling inflammation as potential anti-inflammatory compounds

[92,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146]. Chromones derived from the leaf extracts of plants act as the basic structural compound in the development of various drugs that inhibit infectious diseases caused by a wide range of microbes inferring its antimicrobial property [147,148,149,150,151]. Reactive oxygen species (ROS) act as oxygen moieties impairing the DNA, lipids, proteins, and lipoproteins with their oxidative damage [152,153,154]. Chromones present in the food material (plants) act as potential antioxidants in reducing the lipid peroxidase activity of the ROS and free radicals and disease progression [155,156,157,158,159,160,161,162]. In particular, chromones have been found to be promising in controlling and treating the neurodegenerative disease Alzheimer's, which is caused due to the redox imbalance created by the free radicals and the ROS in the human brain [62,163]. Valentina et al. [164] and Wang et al. [91] have investigated and reported the inhibitory effect of the chromones on the α -amylase and α -glucosidase enzymes to manage postprandial diabetes by delaying sugar uptake in the human body. Further, the role of chromones in the treatment of various disorders such as gastritis, diarrhea, stiff muscles, hypothermic disease, diuretic disease, nephritis, cystitis, pyrexia, rheumatism, headache, hepatitis, cough, bronchitis, asthma, etc. have been reported and described by various researchers worldwide (Table 3) [88,89,90].

6.2. Terpenoids

Terpenoids are abundantly present in nature, especially in plants, and are known to contain oxygen derivatives that bear a hydroxyl group at the C-3 position [77]. To date, about three different terpenoids have been identified in agarwood and *Aquilaria* plant leaves, namely 3-oxo-22-hydroxyhopane, 3b-olean-12-ene-3,23-diol, and hederagenin [13,70,97]. Among all the terpenoids known, sesquiterpenoids are the dominant fragrant compounds observed to be present in agarwood and agarwood products with the presence of three isoprene structural units (Table 3) [82,165]. They exist as volatile compounds in the essential oils extracted from agarwood and are of several types with unique aromatic properties specific to each type [77]. Eudesmanes, otherwise known as selinanes, are bicyclic sesquiterpenoids with a decalin skeleton, diverse functional groups, and multiple chiral centers acting as significant sesquiterpenoids found in agarwood obtained from *Aquilaria agallocha*, *A. sinensis*, *A. crassna*, *A. malaccensis*, and *Gyrinops salicifolia* [13,84,86,103,166,167]. They possess a sweet, woody, honey, balsamic, minty, and fresh floral odor [82]. Guaianes are the sesquiterpenoids made of a five- and seven-membered ring-coupled structures containing 4,10-dimethyl-7-isopropenyl moiety and are observed in *Aquilaria* and *Gyrinops* species with a woody and floral odor [84,97,104,168]. Agarospiranes sesquiterpenes, also known as vetispiranes, are composed of spirocyclic sesquiterpenes present in *Aquilaria agallocha*, *A. sinensis*, *A. crassna*, and *A. malaccensis* with a spicy, peppery, woody, sweet, and balsamic odor [82,168,169,170]. Eremophilanes, known as valencanes, are similar to eudesmanes structurally and differ due to the presence of a specific methyl group in the structure with a characteristic warm woody and minty odor [13,84,86]. Acoranes, responsible for the long-lasting odor of agarwood, are the spiro sesquiterpenes that are rarely obtained from agarwood [97,103,171]. Cadinanes, a bicyclic sesquiterpene, is similar to eudesmanes with the presence of an isopropyl and methyl group in its structure obtained from *A. sinensis* and *A. crassna* [13,172]. Prezizaanes are the tricyclic sesquiterpenes found in *A. malaccensis* with a special fragrance [171,173,174]. Similarly, Zizaanes, a tricyclic, and Humulanes, a humulane-type sesquiterpenes, have also been reported to be obtained from *A. sinensis* and *A. crassna* [103,171,175]. Further, other sesquiterpenoids such as daphnauranols obtained from *A. malaccensis*, 12-Hydroxy-dihydrocyperolone from *G. salicifolia*, malacinones A and B from *A. malaccensis*, and 1,5,9-trimethyl-1,5,9-cyclododecatriene from *A. sinensis* have been reported by researchers such as Ma et al. [175], Li et al. [97], Chen et al. [176], and Ma et al. [177]. Among all the sesquiterpenoids reported, eudesmanes, eremophilanes, and guaianes are the potential sesquiterpenoids obtained

from agarwood of which guaianes are the compounds possessing high structural diversity that are specific to each plant species [77].

Furthermore, sesquiterpenes act as a potential neuroprotective agent and help in combating Alzheimer's disease [178]. Similarly, agarol obtained from *A. malaccensis*, an eudesmane sesquiterpene, and 8bH-Dihydrogmelofuran and gmelofuran, a cadinene sesquiterpenes isolated from *A. malaccensis* and *A. agallocha*, were reported to have anticancer, antioxidant and antimicrobial properties [179]. Antidiabetes activity has been observed in some sesquiterpenoids such as Prezizaane, jinkohol II, aquilarene D, jinkohol, zizaane, agarozizanol E, and isokhusenol acting as inhibitors against α -glucosidase [180].

Moreover, cucurbitacin triterpenoids were found to possess cytotoxic activities, making them a potential candidate for the treatment of cancer [181]. Similarly, β -Caryophyllene, isolated from *A. crassna*, was found to specifically help in the de-proliferation of cancerous cells. Further, β -Caryophyllene was a potential antimicrobial compound effective against various pathogenic strains that include *Bacillus cereus*, *B. subtilis*, *S. aureus*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Aspergillus niger*, *Penicillium citrinum*, *Rhizopus oryzae*, and *Trichoderma reesei* [179]. Flavonoids, benzophenones, xanthenes, lignans, phenolic compounds, degraded sesquiterpenes, alkaloids, and nucleosides are some of the other chemical constituents of the *Aquilaria* species that are available compounds in plants, and they contribute to the structural stability of the plants [58].

7. Economic Importance and Market Value of Agarwood

Aquilaria trees and the agarwood obtained from them are very popular for the wide variety of economically important products obtained from them [182]. Their products, such as wood, wood chips, oil, powder, and flakes, have been used in various applications such as medicinal, religious, cosmetics, and cultural purposes [183,184]. Antonopoulou et al. [185], Barden et al. [1], Kiet [186], Lim and Anack [187], Persoon [188], and Sitepu et al. [189] have reported the potential usage of agarwood oil and powder in Ayurvedic medicine practiced in the Indian subcontinent, and also in Tibetan, Vietnamese, Chinese, and Malaysian medicines. Moreover, it is used for religious and cultural purposes in many countries such as Northeast and Southeast Asia, Taiwan, Korea, and Japan [184,185,190]. Furthermore, Sitepu et al. [189], Barden et al. [1], and Chakrabarty et al. [8] have reported the importance of agarwood in the manufacturing of perfumes, cosmetic products, soaps, shampoos, incense, and other fragrance products worldwide.

Therefore, the market value of agarwood and the demand for its products have increased tremendously. Globally, agarwood products, such as wood pieces, wood chips, powder, oil, dust, incense, and perfumes, have been reported to have a market value of USD several thousand billion [1,182,191]. Agarwood is graded into different categories based on the resin quality; the first grade has a high global demand increasing the market value to about USD 10,000 per kg of wood, and the least grade with USD 30 per kg [192]. Further, according to a report by Nanyang Siang Pau in 2005 and Abdin [193], agarwood oil is generally sold at a cost of USD 30,000 to 40,000 per kg. Similarly, in 2013, Akter et al. [7] reported an estimated increase in the market value of agarwood and its products to reach up to USD 6–8 billion or even up to USD 36 billion by 2017. Similarly, other reports from various sources indicate that the agarwood oil market value is expected to reach about USD 201.03 million by 2026. Hence, it is essential to better understand the basic scientific concepts of agarwood induction and formation to improve its production. Furthermore, the invention of simple technology for induction would enable the easy handling of the techniques by farmers to induce the crops.

8. Conclusions

Agarwood formation in the natural environment, although mysterious, can be produced through artificial inoculation techniques. Agarwood is highly sought after due to its economic value and cultural and medicinal uses around the world. Its high value has caused indiscriminate logging of wild *Aquilaria* trees, thereby causing dominant species to become critically endangered. Therefore, methods of artificial induction are developed to prevent its extinction and reduce the burden on wild *Aquilaria*. The quality of artificially induced agarwood is close to wild agarwood, and its production in *Aquilaria* plantations may be able to meet the ever-increasing demand for this unique fragrance. Several applications, including perfumery and therapeutics, further substantiate the ongoing investigations into agarwood and the growing impetus for sustainable cultivation.

Funding: The authors gratefully acknowledge the support provided by the Universiti Brunei Darussalam through the research grants UBD/RSCH/URC/NIG/1.0/2019/003 and UBD/RSCH/1.13/FICBF(b)/2019/012.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Barden, A.; Anak, N.A.; Mulliken, T.; Song, M. *Heart of the Matter: Agarwood Use and Trade and Cites Implementation for Aquilaria Malaccensis*; TRAFFIC: Cambridge, UK, 2000.
2. Ismail, N.; Ali, N.A.M.; Jamil, M.; Rahiman, M.H.F.; Tajuddin, S.N.; Taib, M.N. A Review Study of Agarwood Oil and Its Quality Analysis. *J. Teknol.* **2014**, *68*. <https://doi.org/10.11113/jt.v68.2419>.
3. Hashim, Y.Z.H.-Y.; Kerr, P.; Abbas, P.; Mohd-Salleh, H. *Aquilaria* spp. (agarwood) as source of health beneficial compounds: A review of traditional use, phytochemistry and pharmacology. *J. Ethnopharmacol.* **2016**, *189*, 331–360. <https://doi.org/10.1016/j.jep.2016.06.055>.
4. Persoon, G.A. *Agarwood: The Life of a Wounded Tree*; In Proceedings of the 2nd International Agarwood Conference, Bangkok/Koh Chang, Thailand, 4–11 March 2007; IIAS: Leiden, The Netherlands, 2007.
5. Chhipa, H.; Kaushik, N. Fungal and Bacterial Diversity Isolated from *Aquilaria malaccensis* Tree and Soil, Induces Agarospore Formation within 3 Months after Artificial Infection. *Front. Microbiol.* **2017**, *8*, 1286. <https://doi.org/10.3389/fmicb.2017.01286>.
6. Ng, L.T.; Chang, Y.S.; Azizol, A.K. A review on agar (gaharu) producing *Aquilaria* species. *J. Trop. For. Prod.* **1997**, *2*, 272–285.
7. Akter, S.; Islam, T.; Zulkefeli, M.; Khan, S.I. Agarwood Production—A Multidisciplinary Field to be Explored in Bangladesh. *Int. J. Pharm. Life Sci.* **2013**, *2*, 22–32. <https://doi.org/10.3329/ijpls.v2i1.15132>.
8. Chakrabarty, K.; Kumar, A.; Menon, V. *Trade in Agarwood*; TRAFFIC: India, New Delhi 1994.
9. Sadgopal, V.B. Exploratory studies in the development of essential oils and their constituents in aromatic plants. Part I. Oil of agarwood. *Soap Perfum. Cosmet.* **1960**, *33*, 41–46.
10. Gibson, I.A.S. The role of fungi in the origin of oleoresin deposits (agaru) in the wood of *Aquilaria agallocha* Roxb. *Bano Biggyan Patrika* **1977**, *6*, 16–26.
11. Mohammad, Y.H.; Shivanand, P.; Metali, F.; Taha, H.; Matussin, N.B.A.; Abdul-Halim, A.M.A.-A.; Mohaimin, A.Z. Agarwood formation in *Aquilaria beccariana* and *Aquilaria microcarpa* in response to inoculation of newly isolated fungi from Brunei Darussalam. *Biodiversitas J. Biol. Divers.* **2021**, *22*. <https://doi.org/10.13057/biodiv/d221002>.
12. Sangareswari, M.; Parthiban, K.T.; Kanna, S.U.; Karthiba, L.; Saravanakumar, D. Fungal Microbes Associated with Agarwood Formation. *Am. J. Plant Sci.* **2016**, *7*, 1445–1452. <https://doi.org/10.4236/ajps.2016.710138>.
13. Pojanagaroon, S.; Kaewrak, C. Mechanical Methods to Stimulate Aloe Wood Formation in *Aquilaria Crassna* Pierre Exhlec. (Kritsana) Trees. *Acta Hort.* **2005**, 161–166. <https://doi.org/10.17660/actahortic.2005.676.20>.
14. Blanchette, R.; Heuveling, V.B.H. Cultivated Agarwood. US 7638145B2, 29 December 2009.
15. Liu, Y.; Chen, H.; Yang, Y.; Zhang, Z.; Wei, J.; Meng, H.; Chen, W.; Feng, J.; Gan, B.; Chen, X.; et al. Whole-tree Agarwood-Inducing Technique: An Efficient Novel Technique for Producing High-Quality Agarwood in Cultivated *Aquilaria sinensis* Trees. *Molecules* **2013**, *18*, 3086–3106. <https://doi.org/10.3390/molecules18033086>.
16. Zhang, X.L.; Liu, Y.Y.; Wei, J.H.; Yang, Y.; Zhang, Z.; Huang, J.Q.; Chen, H.Q.; Liu, Y.J. Production of high-quality agarwood in *Aquilaria sinensis* trees via whole-tree agarwood-induction technology. *Chin. Chem. Lett.* **2012**, *23*, 727–730.

17. Turjaman, M.; Hidayat, A.; Santoso, E. Development of Agarwood Induction Technology Using Endophytic Fungi. In *Agarwood*; Mohamed, R., Ed.; Tropical Forestry; Springer: Berlin, Germany; Singapore, 2016; pp. 57–71.
18. Chong, S.P.; Osman, M.F.; Bahari, N.; Nuri, E.A.; Zakaria, R.; Abdul Rahim, K. Agarwood inducement technology: A method for producing oil grade agarwood in cultivated *Aquilaria malaccensis*. *Lamk J. Agrobiotech.* **2015**, *6*, 1–16.
19. Van Thanh, L.; Van Do, T.; Son, N.H.; Sato, T.; Kozan, O. Impacts of biological, chemical and mechanical treatments on sesquiterpene content in stems of planted *Aquilaria crassna* trees. *Agrofor. Syst.* **2015**, *89*, 973–981. <https://doi.org/10.1007/s10457-015-9829-3>.
20. Nobuchi, T.; Siripatanadilok, S. Preliminary observation of *Aquilaria crassna* wood associated with the formation of aloeswood. *Bull. Kyoto Univ. For.* **1991**, *63*, 226–235.
21. Jong, P.L.; Tsan, P.; Mohamed, R. Gas chromatography–mass spectrometry analysis of agarwood extracts from mature and juvenile *Aquilaria malaccensis*. *Int. J. Agric. Biol.* **2014**, *16*, 644–648.
22. Chen, X.; Liu, Y.; Yang, Y.; Feng, J.; Liu, P.; Sui, C.; Wei, J. Trunk surface agarwood-inducing technique with *Rigidoporus vinctus*: An efficient novel method for agarwood production. *PLoS ONE* **2018**, *13*, e0198111. <https://doi.org/10.1371/journal.pone.0198111>.
23. Setshedi, K.J.; Mutingwende, N.; Ngqwala, N.P. The Use of Artificial Neural Networks to Predict the Physicochemical Characteristics of Water Quality in Three District Municipalities, Eastern Cape Province, South Africa. *Int. J. Environ. Res. Public Health* **2021**, *18*, 5248. <https://doi.org/10.3390/10.3390/ijerph18105248>.
24. Rasool, S.; Mohamed, R. Understanding Agarwood Formation and Its Challenges. In *Agarwood*; Mohamed, R., Ed.; Tropical Forestry; Springer: Berlin, Germany; Singapore, 2016; pp. 39–56. https://doi.org/10.1007/978-981-10-0833-7_3.
25. Chhipa, H.; Chowdhary, K.; Kaushik, N. Artificial production of agarwood oil in *Aquilaria* sp. by fungi: A review. *Phytochem. Rev.* **2017**, *16*, 835–860. <https://doi.org/10.1007/s11101-017-9492-6>.
26. Azren, P.D.; Lee, S.Y.; Emang, D.; Mohamed, R. History and perspectives of induction technology for agarwood production from cultivated *Aquilaria* in Asia: A review. *J. For. Res.* **2018**, *30*, 1–11. <https://doi.org/10.1007/s11676-018-0627-4>.
27. Novriyanti, E.; Santosa, E.; Syafii, W.; Turjaman, M.; Sitepu, I.R. Antifungal activity of wood extract of *Aquilaria crassna* Pierre ex Lecomte against agarwood-inducing fungi, *Fusarium solani*. *J. For. Res.* **2010**, *7*, 155–165.
28. Bose, S.R. The nature of ‘Agaru’ formation. *Sci. Cult.* **1938**, *4*, 89–91.
29. Santoso, E. Pembentukan gaharu dengan cara inokulasi. In Proceedings of the Makalah Diskusi Hasil Penelitian Dalam Menunjang Pemanfaatan Hutan Yang Lestari, Bogor, Indonesia, 11–12 March 1996; pp. 1–3.
30. Tabata, Y.; Widjaja, E.; Mulyaningsih, T.; Parman, I.; Wiriadinata, H.; Mandang, Y.I.; Itoh, T. Structural survey and artificial induction of aloeswood. *Wood Res.* **2003**, *90*, 10–12.
31. Bhattacharyya, B. On the formation and development of agaru in *Aquilaria agallocha*. *Sci. Cult.* **1952**, *18*, 240–243.
32. Tamuli, P.; Boruah, P.; Nath, S.C.; Leclercq, P. Essential Oil of Eaglewood Tree: A Product of Pathogenesis. *J. Essent. Oil Res.* **2005**, *17*, 601–604. <https://doi.org/10.1080/10412905.2005.9699008>.
33. Tamuli, P.; Boruah, P.; Nath, S.C.; Samanta, R. Fungi from disease agarwood tree (*Aquilaria agallocha* Roxb): Two new records. *Adv. For. Res. India* **2000**, *22*, 182–187.
34. Budi, S.; Santoso, E.; Wahyudi, A. Identification of potential types of fungi on establishment agarwood stem of *Aquilaria* spp. *J. Silvikultur Trop.* **2010**, *1*, 1–5.
35. Cui, J.; Guo, S.; Fu, S.; Xiao, P.; Wang, M. Effects of inoculating fungi on agilawood formation in *Aquilaria sinensis*. *Chin. Sci. Bull.* **2013**, *58*, 3280–3287. <https://doi.org/10.1007/s11434-013-5856-5>.
36. Cui, J.; Wang, C.; Guo, S.; Yang, L.; Xiao, P.; Wang, M. Evaluation of fungus-induced agilawood from *Aquilaria sinensis* in China. *Symbiosis* **2013**, *60*, 37–44. <https://doi.org/10.1007/s13199-013-0237-z>.
37. Lin, F.; Mei, W.-L.; Wu, J.; Dai, H.-F. GC-MS analysis of volatile constituents from Chinese eaglewood produced by artificial methods. *Zhong Yao Cai Zhongyao Cai J. Chin. Med. Mater.* **2010**, *33*, 222–225.
38. Han, X.M.; Liang, L.; Zhang, Z.; Li, X.J.; Yang, Y.; Meng, H.; Gao, Z.H.; Xu, Y.H. Sesquiterpenes induced by *Chaetoceros cocoa* on white wood incense. *Chin. J. Tradit. Chin. Med.* **2014**, *39*, 192. (In Chinese).
39. Tian, J.J.; Gao, X.X.; Zhang, W.M.; Wang, L.; Qu, L.H. Molecular identification of endophytic fungi from *Aquilaria sinensis* and artificial agarwood induced by pinholes-infusion technique. *Afr. J. Biotechnol.* **2013**, *12*, 3115–3131.
40. Zhang, Z.; Han, X.M.; Wei, J.H.; Xue, J.; Yang, Y.; Liang, L.; Li, X.J.; Guo, Q.M.; Xu, Y.H.; Gao, Z.H. Compositions and antifungal activities of essential oils from agarwood of *Aquilaria sinensis* (Lour.) Gilg induced by *Lasiodiplodia theobromae* (Pat.) Griffon. *Maubl. J. Braz. Chem. Soc.* **2014**, *25*, 20–26.
41. Huang, H.; Liu, T.; Wu, X.; Guo, J.; Lan, X.; Zhu, Q.; Zheng, X.; Zhang, K. A new antibacterial chromone derivative from mangrove-derived fungus *Penicillium aculeatum* (No. 9EB). *Nat. Prod. Res.* **2017**, *31*, 2593–2598. <https://doi.org/10.1080/14786419.2017.1283498>.
42. Chen, X.Y.; Liu, Y.Y.; Liu, P.W.; Peng, D.Q.; Wei, J.H. Study on biological characteristics of two strains of *Lasiodiplodia theobromae* promoting agarwood formation. *Acta Agric. Jiangxi* **2017**, *29*, 95–98.
43. Chen, X.; Sui, C.; Liu, Y.; Yang, Y.; Liu, P.; Zhang, Z.; Wei, J. Agarwood Formation Induced by Fermentation Liquid of *Lasiodiplodia theobromae*, the Dominating Fungus in Wounded Wood of *Aquilaria sinensis*. *Curr Microbiol.* **2017**, *74*, 460–468. <https://doi.org/10.1007/s00284-016-1193-7>.
44. Gu, L.P.; Zheng, K.; Liu, Y.R.; Ma, H.F.; Xiao, Z.Y. Effects of two fungi on xylem tissue structure of white wood incense. *West. For. Sci.* **2018**, *47*, 141–144. (In Chinese).

45. Huang, Y.Q.; Guo, H.; Chen, X.Y.; Zhong, Z.J.; Li, H.H.; Zhang, W.M.; Gao, X.X. Analysis of secondary metabolites of rice black spore A8 and its induced artificial agarwood. *Tradit. Chin. Med.* **2018**, *41*, 1662–1667. (In Chinese).
46. Zheng, K.; Gu, L.P.; Xiao, Z.Y.; Ma, H.F. Study on the effects of *Fusarium oxysporum* and *Dichromium cocoanum* on the chemical composition of balsam wood xylem. *For. Investig. Plan.* **2019**, *44*, 27–32. (In Chinese).
47. Rahayu, G.; Putridan Juliarni, A.L. Acremonium and methyl-jasmonate induce terpenoid formation in agarwood tree (*Aquilaria crassna*). In Proceedings of the Makalahdi Presenta Sikandalam 3rd Asian Conference on Crop Protection, 22–24 August 2007, Yogyakarta, Indonesia; pp. 22–24.
48. Santoso, E.; Irianto, R.S.B.; Turjaman, M.; Sitepu, I.R.; Santosa, S.; Najmulah; Yani, A.; dan Aryanto. *Gaharu-Producing Tree Induction Technology*. In *Development of Gaharu Production, Technology*; Turjaman, M., Ed.; Ministry of Forestry: Bogor, Indonesia 2011; pp. 31–46.
49. Mohamed, R.; Jong, P.L.; Kamziah, A.K. Fungal inoculation induces agarwood in young *Aquilaria malaccensis* trees in the nursery. *J. For. Res.* **2014**, *25*, 201–204. <https://doi.org/10.1007/s11676-013-0395-0>.
50. Lisdayani, L.; Anna, N.; Siregar, E.B.M. Isolation and identifying of fungi from the stem of agarwood (*Aquilaria malaccensis* Lamk.) was had been inoculation. *Peronema For. Sci.* **2015**, *4*, 1–5.
51. Faizal, A.; Esyanti, R.R.; Aulianisa, E.N.; Santoso, E.; Turjaman, M. Formation of agarwood from *Aquilaria malaccensis* in response to inoculation of local strains of *Fusarium solani*. *Trees* **2017**, *31*, 189–197.
52. Sen, S.; Dehingia, M.; Talukdar, N.C.; Khan, M. Chemometric analysis reveals links in the formation of fragrant bio-molecules during agarwood (*Aquilaria malaccensis*) and fungal interactions. *Sci. Rep.* **2017**, *7*, 44406. <https://doi.org/10.1038/srep44406>.
53. Nasution, A.A.; Siregar, U.J.; Miftahudin; Turjaman, M. Identification of chemical compounds in agarwood-producing species *Aquilaria malaccensis* and *Gyrinops versteegii*. *J. For. Res.* **2019**, *31*, 1371–1380. <https://doi.org/10.1007/s11676-018-00875-9>.
54. Triadiati, T.; Carolina, D.A.; Miftahudin. Induksi pembentukan gaharu menggunakan berbagai media tanam dan cendawan *Acremonium* sp dan *Fusarium* sp. pada *Aquilaria crassna*. *J. Sumberd. Hayati* **2016**, *2*, 1–6.
55. Iskandar, D.; Suhendra, A. Uji Inokulasi *Fusarium* Sp Untuk Produksi Gaharu Pada Budidaya A. Beccariana. *J. Sains dan Teknol. Indones.* **2013**, *14*. <https://doi.org/10.29122/jsti.v14i3.924>.
56. Faizal, A.; Azar, A.W.P.; Turjaman, M.; Esyanti, R.R. *Fusarium solani* induces the formation of agarwood in *Gyrinops versteegii* (Gilg.) Domke branches. *Symbiosis* **2020**, *81*, 15–23. <https://doi.org/10.1007/s13199-020-00677-w>.
57. Subasinghe, S.M.C.U.P.; Hitihamu, H.I.D.; Fernando, K.M.E.P. Use of two fungal species to induce agarwood resin formation in *Gyrinops walla*. *J. For. Res.* **2018**, *30*, 721–726. <https://doi.org/10.1007/s11676-018-0654-1>.
58. Tan, C.S.; Isa, N.; Ismail, I.; Zainal, Z. Agarwood Induction: Current Developments and Future Perspectives. *Front. Plant Sci.* **2019**, *10*, 122. <https://doi.org/10.3389/fpls.2019.00122>.
59. Khadem, S.; Marles, R.J. Chromone and Flavonoid Alkaloids: Occurrence and Bioactivity. *Molecules* **2011**, *17*, 191–206. <https://doi.org/10.3390/molecules17010191>.
60. Yang, L.; Qiao, L.R.; Xie, D.; Yuan, Y.H.; Chen, N.H.; Dai, J.G.; Guo, S.X. 2-(2-phenylethyl) chromones from Chinese eaglewood. *Phytochemistry* **2012**, *76*, 92–97.
61. Tawfik, H.A.; Ewies, E.F.; El-Hamouly, W.S. ChemInform Abstract: Synthesis of Chromones and Their Applications during the Last Ten Years. *ChemInform* **2015**, *46*. <https://doi.org/10.1002/chin.201513331>.
62. Reis, J.; Gaspar, A.; Milhazes, N.; Borges, F. Chromone as a Privileged Scaffold in Drug Discovery: Recent Advances. *J. Med. Chem.* **2017**, *60*, 7941–7957. <https://doi.org/10.1021/acs.jmedchem.6b01720>.
63. Wang, X.; Gao, B.; Liu, X.; Dong, X.; Zhang, Z.; Fan, H.; Zhang, L.; Wang, J.; Shi, S.; Tu, P. Salinity stress induces the production of 2-(2-phenylethyl)chromones and regulates novel classes of responsive genes involved in signal transduction in *Aquilaria sinensis calli*. *BMC Plant Biol.* **2016**, *16*, 119. <https://doi.org/10.1186/s12870-016-0803-7>.
64. Liao, G.; Dong, W.-H.; Yang, J.-L.; Li, W.; Wang, J.; Mei, W.-L.; Dai, H.-F. Monitoring the Chemical Profile in Agarwood Formation within One Year and Speculating on the Biosynthesis of 2-(2-Phenylethyl)Chromones. *Molecules* **2018**, *23*, 1261. <https://doi.org/10.3390/molecules23061261>.
65. Goel, S.; Shashi, S.; Makrandi, J.K. Synthesis of 2-(2-Phenylethyl)chromones. *ChemInform* **2006**, *37*. <https://doi.org/10.1002/chin.200623140>.
66. Rohmer, M. The discovery of a mevalonate-independent pathway for isoprenoid biosynthesis in bacteria, algae and higher plants. *Nat. Prod. Rep.* **1999**, *16*, 565–574. <https://doi.org/10.1039/a709175c>.
67. Dong, L.; Jongedijk, E.; Bouwmeester, H.; Van Der Krol, A. Monoterpene biosynthesis potential of plant subcellular compartments. *New Phytol.* **2015**, *209*, 679–690. <https://doi.org/10.1111/nph.13629>.
68. Singh, B.; Sharma, R.A. Plant terpenes: Defense responses, phylogenetic analysis, regulation and clinical applications. *3 Biotech* **2014**, *5*, 129–151. <https://doi.org/10.1007/s13205-014-0220-2>.
69. Yang, L.; Qiao, L.; Ji, C.; Xie, D.; Gong, N.-B.; Lu, Y.; Zhang, J.; Dai, J.; Guo, S. Antidepressant Abietane Diterpenoids from Chinese Eaglewood. *J. Nat. Prod.* **2013**, *76*, 216–222. <https://doi.org/10.1021/np3006925>.
70. Liu, X.-M.; Tao, T.-T.; Meng, X.-X.; Zhang, W.-W.; Chang, J.; Xu, F. Cloning and Expression Analysis of a Farnesyl Diphosphate Synthase (FPPS) Gene from *Chamaemelum nobile*. *Not. Bot. Horti Agrobot. Cluj-Napoca* **2017**, *45*, 358–364. <https://doi.org/10.15835/nbha45210858>.
71. Pateraki, I.; Heskes, A.M.; Hamberger, B. Cytochromes P450 for Terpene Functionalisation and Metabolic Engineering. *Adv. Biochem. Eng./Biotechnol.* **2015**, *148*, 107–139. https://doi.org/10.1007/10_2014_301.

72. Kumeta, Y.; Ito, M. Characterization of δ -Guaiene Synthases from Cultured Cells of *Aquilaria*, Responsible for the Formation of the Sesquiterpenes in Agarwood. *Plant Physiol.* **2010**, *154*, 1998–2007. <https://doi.org/10.1104/pp.110.161828>.
73. Hidayat, W.; Shakaff, A.Y.M.; Ahmad, M.N.; Adom, A.H. Classification of Agarwood Oil Using an Electronic Nose. *Sensors* **2010**, *10*, 4675–4685. <https://doi.org/10.3390/s100504675>.
74. Tawfeek, N.; Mahmoud Mona F.; Hamdan Dalia I.; Sobeh M.; Farrag N.; Wink M.; El-Shazly Assem M. Phytochemistry, Pharmacology and Medicinal Uses of Plants of the Genus *Salix*: An Updated Review. *Front. Pharmacol.* **2021**, *12*, 1–30. DOI=10.3389/fphar.2021.593856.
75. Hidayat, A.; Turjaman, M.; Qamyari, R.; Imanuddin, R.; Tohir, D.; Rahmanto, R.G.H.; Susilowati, A. Bioactive composition, antifungal, antioxidant, and anticancer potential of agarwood essential oil from decaying logs (*Gyrinops* spp.) of Papua Island (Indonesia). *J. Appl. Pharm. Sci.* **2021**, *11*, 70–78. <https://doi.org/10.7324/japs.2021.1101010>.
76. Yuan, H.W.; Zhao, J.P.; Liu, Y.B.; Qiu, Y.X.; Xie, Q.L.; Li, M.J.; Khan, I.A.; Wang, W. Advance in studies on chemical constituents, pharmacology and quality control of *Aquilaria sinensis*. *Digit. Chin. Med.* **2018**, *1*, 316–330.
77. Li, W.; Chen, H.-Q.; Wang, H.; Mei, W.-L.; Dai, H.-F. Natural products in agarwood and *Aquilaria* plants: Chemistry, biological activities and biosynthesis. *Nat. Prod. Rep.* **2020**, *38*, 528–565. <https://doi.org/10.1039/d0np00042f>.
78. Wang, S.L.; Hwang, T.L.; Chung, M.L.; Sung, P.J.; Shu, C.W.; Cheng, M.J.; Chen, J.J. New flavones, a 2-(2-phenylethyl)-4H-chromen-4-one derivative, and anti-inflammatory constituents from the stem barks of *Aquilaria sinensis*. *Molecules* **2015**, *20*, 20912–20925.
79. Cock, I.E. The Genus *Aloe*: Phytochemistry and Therapeutic Uses Including Treatments for Gastrointestinal Conditions and Chronic Inflammation. In *Novel Natural Products: Therapeutic Effects in Pain, Arthritis and Gastro-Intestinal Diseases*; Rainsford, K., Powanda, M., Whitehouse, M., Eds.; Springer: Basel, Switzerland, 2015; Volume, 70, pp. 179–235. https://doi.org/10.1007/978-3-0348-0927-6_6.
80. Liu, Y.Y.; Chen, D.L.; Yu, Z.X.; Can-Hong, W.; Feng, J.; Meng, Y.; Wei, J.H. New 2-(2-phenylethyl)chromone derivatives from agarwood and their inhibitory effects on tumor cells. *Nat Prod Res.* **2020**, *34*, 1721–1727. doi: 10.1080/14786419.2018.1528591.
81. Huo, H.X.; Zhu, Z.X.; Song, Y.L.; Shi, S.P.; Sun, J.; Sun, H.; Zhao, Y.F.; Zheng, J.; Ferreira, D.; Zjawiony, J.K.; et al. Anti-inflammatory dimeric 2-(2-phenylethyl) chromones from the resinous wood of *Aquilaria sinensis*. *J. Nat. Prod.* **2017**, *81*, 543–553.
82. Naef, R. The volatile and semi-volatile constituents of agarwood, the infected heartwood of *Aquilaria* species: A review. *Flavour Fragr. J.* **2011**, *26*, 73–87. <https://doi.org/10.1002/ffj.2034>.
83. Wang, S.; Yu, Z.; Wang, C.; Wu, C.; Guo, P.; Wei, J. Chemical Constituents and Pharmacological Activity of Agarwood and *Aquilaria* Plants. *Molecules* **2018**, *23*, 342. <https://doi.org/10.3390/molecules23020342>.
84. Shao, H.; Kong, F.D.; Wang, H.; Mei, W.L.; Dai, H.F. Qinanmer, a new compound from chinese agarwood ‘Qi-nan’ originating from *Aquilaria sinensis*. *J. Asian Nat. Prod. Res.* **2017**, *19*, 935–940.
85. Kuang, T.D.; Chen, H.Q.; Kong, F.D.; Cai, C.H.; Yang, L.; Mei, W.L.; Dai, H.F. UPLC-MS-guided isolation of single ether linkage dimeric 2-(2-phenylethyl)chromones from *Aquilaria sinensis* *Phytochem. Lett.* **2018**, *26*, 96–100.
86. Wu, B.; Kwon, S.W.; Hwang, G.S.; Park, J.H. Eight New 2-(2-Phenylethyl)chromone (=2-(2-Phenylethyl)-4H-1-benzopyran-4-one) Derivatives from *Aquilaria malaccensis* Agarwood. *Helvetica Chim. Acta* **2012**, *95*, 1657–1665. <https://doi.org/10.1002/hlca.201200069>.
87. Liu, J.M.; Gao, Y.H.; Xu, H.H.; Xu, Z.Q. Chemical constituents of *Lignum Aquilariae Resinatum* (II). *Chin. Tradit. Herb. Drugs* **2007**, *38*, 1138–1140.
88. Sharma, S.K.; Kumar, S.; Chand, K.; Kathuria, A.; Gupta, A.; Jain, R. An update on natural occurrence and biological activity of chromones. *Curr. Med. Chem.* **2011**, *18*, 3825–3852. <https://doi.org/10.2174/092986711803414359>.
89. Nazhand, A.; Durazzo, A.; Lucarini, M.; Romano, R.; Mobilia, M.A.; Izzo, A.A.; Santini, A. Human health-related properties of chromones: An overview. *Nat. Prod. Res.* **2019**, *34*, 137–152. <https://doi.org/10.1080/14786419.2019.1678618>.
90. Kulshrestha, A.; Rupanwal, R.; Singh, N.; Panhekar, D.; Ey, J. Therapeutic Potential of Chromones. *Int. J. Chem. Sci.* **2017**, *15*, 200.
91. Wang, G.; Chen, M.; Qiu, J.; Xie, Z.; Cao, A. Synthesis, in vitro α -glucosidase inhibitory activity and docking studies of novel chromone-isatin derivatives. *Bioorg. Med. Chem. Lett.* **2018**, *28*, 113–116. <https://doi.org/10.1016/j.bmcl.2017.11.047>.
92. Chen, D.; Xu, Z.; Chai, X.; Zeng, K.; Jia, Y.; Bi, D.; Ma, Z.; Tu, P. Nine 2-(2-Phenylethyl)chromone Derivatives from the Resinous Wood of *Aquilaria sinensis* and Their Inhibition of LPS-Induced NO Production in RAW 264.7 Cells. *Eur. J. Org. Chem.* **2012**, *2012*, 5389–5397. <https://doi.org/10.1002/ejoc.201200725>.
93. Liao, G.; Mei, W.-L.; Dong, W.-H.; Li, W.; Wang, P.; Kong, F.-D.; Gai, C.-J.; Song, X.-Q.; Dai, H.-F. 2-(2-Phenylethyl)chromone derivatives in artificial agarwood from *Aquilaria sinensis*. *Fitoterapia* **2016**, *110*, 38–43. <https://doi.org/10.1016/j.fitote.2016.01.011>.
94. Yagura, T.; Ito, M.; Kiuchi, F.; Honda, G.; Shimada, Y. Four New 2-(2-Phenylethyl)chromone Derivatives from Withered Wood of *Aquilaria sinensis*. *Chem. Pharm. Bull.* **2003**, *51*, 560–564. <https://doi.org/10.1248/cpb.51.560>.
95. Wang, S.-C.; Wang, F.; Yue, C.-H. Chemical constituents from the petioles and leaves of *Aquilaria sinensis*. *Biochem. Syst. Ecol.* **2015**, *61*, 458–461. <https://doi.org/10.1016/j.bse.2015.07.022>.
96. Liu, J.M.; Gao, Y.H.; Xu, H.H. Chemical constituents of *Aquilaria sinensis*. *Chin. Tradit. Herb. Drugs* **2006**, *37*, 325–327.
97. Li, W.; Liao, G.; Dong, W.-H.; Kong, F.-D.; Wang, P.; Wang, H.; Mei, W.-L.; Dai, H.-F. Sesquiterpenoids from Chinese Agarwood Induced by Artificial Holing. *Molecules* **2016**, *21*, 274. <https://doi.org/10.3390/molecules21030274>.
98. Wu, Y.; Liu, C.; Li, H.-F.; Sun, J.-B.; Li, Y.-Y.; Gu, W.; Wang, D.-Y.; Liu, J.-G.; Hu, Y.-L. A novel neolignan glycoside from *Aquilaria sinensis*. *Biochem. Syst. Ecol.* **2014**, *55*, 41–45. <https://doi.org/10.1016/j.bse.2014.02.030>.

99. Gao, Y.-H.; Liu, J.-M.; Lu, H.-X.; Wei, Z.-X. Two New 2-(2-Phenylethyl)chromen-4-ones from *Aquilaria sinensis* (Lour.) Gilg. *Helvetica Chim. Acta* **2012**, *95*, 951–954. <https://doi.org/10.1002/hlca.201100442>.
100. Quan, L.; Zhang, J.; Wu, Y. One Novel 2-(2-Phenylethyl)Chromone Glycoside from *Aquilaria sinensis*. *Chem. Nat. Compd.* **2017**, *53*, 635–637. <https://doi.org/10.1007/s10600-017-2078-0>.
101. Liao, G.; Mei, W.-L.; Kong, F.-D.; Li, W.; Yuan, J.-Z.; Dai, H.-F. 5,6,7,8-Tetrahydro-2-(2-phenylethyl)chromones from artificial agarwood of *Aquilaria sinensis* and their inhibitory activity against acetylcholinesterase. *Phytochemistry* **2017**, *139*, 98–108. <https://doi.org/10.1016/j.phytochem.2017.04.011>.
102. Xiang, P.; Mei, W.; Chen, H.; Kong, F.; Wang, H.; Liao, G.; Zhou, L.; Dai, H. Four new bi-phenylethylchromones from artificial agarwood. *Fitoterapia* **2017**, *120*, 61–66. <https://doi.org/10.1016/j.fitote.2017.05.012>.
103. Huo, H.X.; Zhu, Z.X.; Pang, D.R.; Li, Y.T.; Huang, Z.; Shi, S.P.; Zheng, J.; Zhang, Q.; Zhao, Y.F.; Tu, P.F.; et al. Anti-neuroinflammatory sesquiterpenes from Chinese eaglewood. *Fitoterapia* **2015**, *106*, 115–121.
104. Zhao, H.; Peng, Q.; Han, Z.; Yang, L.; Wang, Z. Three New Sesquiterpenoids and One New Sesquiterpenoid Derivative from Chinese Eaglewood. *Molecules* **2016**, *21*, 281. <https://doi.org/10.3390/molecules21030281>.
105. Yang, D.-L.; Wang, H.; Guo, Z.-K.; Li, W.; Mei, W.-L.; Dai, H.-F. Fragrant agarofuran and eremophilane sesquiterpenes in agarwood 'Qi-Nan' from *Aquilaria sinensis*. *Phytochem. Lett.* **2014**, *8*, 121–125. <https://doi.org/10.1016/j.phytol.2014.03.003>.
106. Li, W.; Cai, C.-H.; Guo, Z.-K.; Wang, H.; Zuo, W.-J.; Dong, W.-H.; Mei, W.-L.; Dai, H.-F. Five new eudesmane-type sesquiterpenoids from Chinese agarwood induced by artificial holing. *Fitoterapia* **2015**, *100*, 44–49. <https://doi.org/10.1016/j.fitote.2014.11.010>.
107. Yang, D.; Wang, J.; Li, W.; Dong, W.; Mei, W.; Dai, H. New guaiane and acorane sesquiterpenes in high quality agarwood Qi-Nan from *Aquilaria sinensis*. *Phytochem. Lett.* **2016**, *17*, 94–99. <https://doi.org/10.1016/j.phytol.2016.07.017>.
108. Peng, K.; Mei, W.-L.; Zhao, Y.-X.; Tan, L.-H.; Wang, Q.-H.; Dai, H.-F. A novel degraded sesquiterpene from the fresh stem of *Aquilaria sinensis*. *J. Asian Nat. Prod. Res.* **2011**, *13*, 951–955. <https://doi.org/10.1080/10286020.2011.598860>.
109. Ma, C.T.; Eom, T.; Cho, E.; Wu, B.; Kim, T.R.; Oh, K.B.; Han, S.B.; Kwon, S.W.; Park, J.H. Aquilanol A and B, Macrocyclic Humulene-Type Sesquiterpenoids from the Agarwood of *Aquilaria malaccensis*. *J. Nat. Prod.* **2017**, *80*, 3043–3048.
110. Cheng, J.T.; Han, Y.Q.; He, J.; De Wu, X.; Bin Dong, L.; Peng, L.Y.; Li, Y.; Zhao, Q.S. Two new tirucallane triterpenoids from the leaves of *Aquilaria sinensis*. *Arch. Pharmacol. Res.* **2013**, *36*, 1084–1089. <https://doi.org/10.1007/s12272-013-0088-4>.
111. Korinek, M.; Wagh, V.D.; Lo, I.W.; Hsu, Y.M.; Hsu, H.Y.; Hwang, T.L.; Wu, Y.C.; Cheng, Y.B.; Chen, B.H.; Chang, F.R. Antiallergic phorbol ester from the seeds of *Aquilaria malaccensis*. *Int. J. Mol. Sci.* **2016**, *17*, 398.
112. Wagh, V.D.; Korinek, M.; Lo, I.W.; Hsu, Y.M.; Chen, S.L.; Hsu, H.Y.; Hwang, T.L.; Wu, Y.C.; Chen, B.H.; Cheng, Y.B.; et al. Inflammation modulatory phorbol esters from the seeds of *Aquilaria malaccensis*. *J. Nat. Prod.* **2017**, *80*, 1421–1427.
113. Abenavoli, L.; Izzo, A.A.; Milic, N.; Cicala, C.; Santini, A.; Capasso, R. Milk thistle (*Silybum marianum*): A concise overview on its chemistry, pharmacological, and nutraceutical uses in liver diseases. *Phytother. Res.* **2018**, *32*, 2202–2213.
114. Andrew, R.; A Izzo, A. Principles of pharmacological research of nutraceuticals. *J. Cereb. Blood Flow Metab.* **2017**, *174*, 1177–1194. <https://doi.org/10.1111/bph.13779>.
115. Ruggeri, M.; Coretti, S.; Prete, S.; Romano, F.; Orlando, V.; Codella, P.; Di Brino, E. Economic evaluation of screening programs for hepatitis C virus infection: Evidence from literature. *Risk Manag. Health Policy* **2015**, *8*, 45–54. <https://doi.org/10.2147/rmhp.s56911>.
116. Daliu, P.; Santini, A.; Novellino, E. A decade of nutraceutical patents: Where are we now in 2018? *Expert Opin. Ther. Pat.* **2018**, *28*, 875–882.
117. Daliu, P.; Santini, A.; Novellino, E. From pharmaceuticals to nutraceuticals: Bridging disease prevention and management. *Expert Rev. Clin. Pharmacol.* **2018**, *12*, 1–7. <https://doi.org/10.1080/17512433.2019.1552135>.
118. Durazzo, A.; D'Addezio, L.; Camilli, E.; Piccinelli, R.; Turrini, A.; Marletta, L.; Marconi, S.; Lucarini, M.; Lisciani, S.; Gabrielli, P.; et al. From Plant Compounds to Botanicals and Back: A Current Snapshot. *Molecules* **2018**, *23*, 1844. <https://doi.org/10.3390/molecules23081844>.
119. Durazzo, A.; Lucarini, M.; Souto, E.B.; Cicala, C.; Caiazzo, E.; Izzo, A.A.; Novellino, E.; Santini, A. Polyphenols: A concise overview on the chemistry, occurrence, and human health. *Phytother. Res.* **2019**, *33*, 2221–2243. <https://doi.org/10.1002/ptr.6419>.
120. Kumari, A.; Singh, R.K. Medicinal chemistry of indole derivatives: Current to future therapeutic perspectives. *Bioorg. Chem.* **2019**, *89*, 103021. <https://doi.org/10.1016/j.bioorg.2019.103021>.
121. Menditto, E.; Cahir, C.; Aza-Pascual-Salcedo, M.; Bruzzese, D.; Poblador-Plou, B.; Malo, S.; Costa, E.; Gonzalez-Rubio, F.; Gimeno-Miguel, A.; Orlando, V.; et al. Adherence to chronic medication in older populations: Application of a common protocol among three European cohorts. *Patient Prefer. Adherence* **2018**, *12*, 1975–1987.
122. Salehi, B.; Armstrong, L.; Rescigno, A.; Yeskaliyeva, B.; Seitimova, G.; Beyatli, A.; Sharmeen, J.; Mahomoodally, M.F.; Sharopov, F.; Durazzo, A.; et al. Lamium Plants—A Comprehensive Review on Health Benefits and Biological Activities. *Molecules* **2019**, *24*, 1913. <https://doi.org/10.3390/molecules24101913>.
123. Salehi, B.; Venditti, A.; Sharifi-Rad, M.; Kregiel, D.; Sharifi-Rad, J.; Durazzo, A.; Lucarini, M.; Santini, A.; Souto, E.B.; Novellino, E.; et al. The Therapeutic Potential of Apigenin. *Int. J. Mol. Sci.* **2019**, *20*, 1305. <https://doi.org/10.3390/ijms20061305>.
124. Santini, A.; Novellino, E. Nutraceuticals: Beyond the diet before the drugs. *Curr. Bioact. Compd.* **2014**, *10*, 1–12.
125. Santini, A.; Tenore, G.C.; Novellino, E. Nutraceuticals: A paradigm of proactive medicine. *Eur. J. Pharm. Sci.* **2017**, *96*, 53–61. <https://doi.org/10.1016/j.ejps.2016.09.003>.

126. Scala, D.; Menditto, E.; Armellino, M.F.; Manguso, F.; Monetti, V.M.; Orlando, V.; Antonino, A.; Makoul, G.; De Palma, M. Italian translation and cultural adaptation of the communication assessment tool in an outpatient surgical clinic. *BMC Health Serv. Res.* **2016**, *16*, 1–7. <https://doi.org/10.1186/s12913-016-1411-9>.
127. Duan, Y.-D.; Jiang, Y.-Y.; Guo, F.-X.; Chen, L.-X.; Xu, L.-L.; Zhang, W.; Liu, B. The antitumor activity of naturally occurring chromones: A review. *Fitoterapia* **2019**, *135*, 114–129. <https://doi.org/10.1016/j.fitote.2019.04.012>.
128. Vanguru, M.; Merugu, R.; Garimella, S.; E, L. A Review on the Synthetic Methodologies of Chromones. *Asian J. Pharm. Clin. Res.* **2018**, *11*, 9–16. <https://doi.org/10.22159/ajpcr.2018.v11i12.27960>.
129. Meydani, A.; Yousefi, S.; Gharibi, R.; Kazemi, S.; Teimouri, M.B. Synthesis of a new series of furopyranone- and furocoumarin-chromone conjugates followed by in-vitro cytotoxicity activity evaluation, and molecular docking study. *Chem. Sel.* **2019**, *4*, 3315–3324.
130. Elsayed, S.A.; Butler, I.S.; Claude, B.J.; Mostafa, S.I. Synthesis, characterization and anticancer activity of 3-formylchromone benzoylhydrazone metal complexes. *Transit. Met. Chem.* **2014**, *40*, 179–187. <https://doi.org/10.1007/s11243-014-9904-z>.
131. Gautam, R.; Jachak, S.M.; Kumar, V.; Mohan, C.G. Synthesis, biological evaluation and molecular docking studies of stellatin derivatives as cyclooxygenase (COX-1, COX-2) inhibitors and anti-inflammatory agents. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 1612–1616. <https://doi.org/10.1016/j.bmcl.2011.01.116>.
132. Jachak, S.M.; Gautam, R.; Selvam, C.; Madhan, H.; Srivastava, A.; Khan, T. Anti-inflammatory, cyclooxygenase inhibitory and antioxidant activities of standardized extracts of *Tridax procumbens* L. *Fitoterapia* **2011**, *82*, 173–177. <https://doi.org/10.1016/j.fitote.2010.08.016>.
133. Shaveta; Singh, A.; Kaur, M.; Sharma, S.; Bhatti, R.; Singh, P. Rational design, synthesis and evaluation of chromone-indole and chromone-pyrazole based conjugates: Identification of a lead for anti-inflammatory drug. *Eur. J. Med. Chem.* **2014**, *77*, 185–192. <https://doi.org/10.1016/j.ejmech.2014.03.003>.
134. Joo, C.; Venkateswararao, E.; Lee, K.-C.; Sharma, V.K.; Kyung, M.-S.; Kim, Y.; Jung, S.-H. Novel interleukin-5 inhibitors based on hydroxyethylaminomethyl-4H-chromen-4-one scaffold. *Bioorg. Med. Chem.* **2012**, *20*, 5757–5762. <https://doi.org/10.1016/j.bmc.2012.08.006>.
135. Thanigaimalai, P.; Le Hoang, T.A.; Lee, K.-C.; Sharma, V.K.; Bang, S.-C.; Yun, J.H.; Roh, E.; Kim, Y.; Jung, S.-H. Synthesis and evaluation of novel chromone analogs for their inhibitory activity against interleukin-5. *Eur. J. Med. Chem.* **2010**, *45*, 2531–2536. <https://doi.org/10.1016/j.ejmech.2010.02.041>.
136. Venkateswararao, E.; Kim, M.-S.; Sharma, V.K.; Lee, K.-C.; Subramanian, S.; Roh, E.; Kim, Y.; Jung, S.-H. Identification of novel chromenone derivatives as interleukin-5 inhibitors. *Eur. J. Med. Chem.* **2013**, *59*, 31–38. <https://doi.org/10.1016/j.ejmech.2012.11.007>.
137. Venkateswararao, E.; Manickam, M.; Boggu, P.; Kim, Y.; Jung, S.-H. Exploration of benzamidochromenone derivatives with conformational restrictor as interleukin-5 inhibitors. *Bioorg. Med. Chem.* **2015**, *23*, 2498–2504. <https://doi.org/10.1016/j.bmc.2015.03.045>.
138. Altavilla, D.; Squadrito, F.; Bitto, A.; Polito, F.; Burnett, B.; Di Stefano, V.; Minutoli, L. Flavocoxid, a dual inhibitor of cyclooxygenase and 5-lipoxygenase, blunts pro-inflammatory phenotype activation in endotoxin-stimulated macrophages. *J. Cereb. Blood Flow Metab.* **2009**, *157*, 1410–1418. <https://doi.org/10.1111/j.1476-5381.2009.00322.x>.
139. Ribeiro, D.; Freitas, M.; Tomé, S.M.; Silva, A.M.; Porto, G.; Cabrita, E.J.; Marques, M.M.B.; Fernandes, E. Inhibition of LOX by flavonoids: A structure–activity relationship study. *Eur. J. Med. Chem.* **2013**, *72*, 137–145. <https://doi.org/10.1016/j.ejmech.2013.11.030>.
140. An, H.-J.; Nugroho, A.; Song, B.-M.; Park, H.-J. Isoeugenin, a Novel Nitric Oxide Synthase Inhibitor Isolated from the Rhizomes of *Imperata cylindrica*. *Molecules* **2015**, *20*, 21336–21345. <https://doi.org/10.3390/molecules201219767>.
141. Förstermann, U.; Sessa, W.C. Nitric oxide synthases: Regulation and function. *Eur. Heart J.* **2012**, *33*, 829–837. <https://doi.org/10.1093/eurheartj/ehr304>.
142. Gao, H.-Y.; Wang, H.-Y.; Li, G.-Y.; Du, X.-W.; Zhang, X.-T.; Han, Y.; Huang, J.; Li, X.-X.; Wang, J.-H. Constituents from Zhuyeqing Liquor and their inhibitory effects on nitric oxide production. *Phytochem. Lett.* **2013**, *7*, 150–155. <https://doi.org/10.1016/j.phytol.2013.11.011>.
143. Jia, B.-X.; Zeng, X.-L.; Ren, F.-X.; Jia, L.; Chen, X.-Q.; Yang, J.; Liu, H.-M.; Wang, Q. Baeckins F–I, four novel C-methylated biflavonoids from the roots of *Baeckea frutescens* and their anti-inflammatory activities. *Food Chem.* **2014**, *155*, 31–37. <https://doi.org/10.1016/j.foodchem.2014.01.022>.
144. Kim, Y.A.; Kong, C.-S.; Park, H.H.; Lee, E.; Jang, M.-S.; Nam, K.-H.; Seo, Y. Anti-Inflammatory Activity of Heterocarpin from the Salt Marsh Plant *Corydalis heterocarpa* in LPS-Induced RAW 264.7 Macrophage Cells. *Molecules* **2015**, *20*, 14474–14486. <https://doi.org/10.3390/molecules200814474>.
145. Liu, G.-B.; Xu, J.-L.; Geng, M.; Xu, R.; Hui, R.-R.; Zhao, J.-W.; Xu, Q.; Xu, H.-X.; Li, J.-X. Synthesis of a novel series of diphenolic chromone derivatives as inhibitors of NO production in LPS-activated RAW264.7 macrophages. *Bioorg. Med. Chem.* **2010**, *18*, 2864–2871. <https://doi.org/10.1016/j.bmc.2010.03.020>.
146. Pham, T.-A.; Che, H.; Phan, P.-T.; Lee, J.-W.; Kim, S.-S.; Park, H. Oroxylin A analogs exhibited strong inhibitory activities against iNOS-mediated nitric oxide (NO) production. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 2534–2535.
147. Cano, P.A.; Islas-Jacome, A.; Rangel-Serrano, A.; Anaya-Velazquez, F.; Padilla-Vaca, F.; Trujillo-Esquivel, E.; Ponce-Noyola, P.; Martinez-Richa, A.; Gamez-Montano, R. In vitro studies of chromonetetrazoles against pathogenic protozoa, bacteria, and fungi. *Molecules* **2015**, *20*, 12436–12449.

148. He, J.; Li, Z.-H.; Ai, H.-L.; Feng, T.; Liu, J.-K. Anti-bacterial chromones from cultures of the endophytic fungus *Bipolaris eleusines*. *Nat. Prod. Res.* **2018**, *33*, 3515–3520. <https://doi.org/10.1080/14786419.2018.1486313>.
149. Hiruy, M.; Bisrat, D.; Mazumder, A.; Asres, K. Two chromones with antimicrobial activity from the leaf latex of *Aloe monticola* Reynolds. *Nat. Prod. Res.* **2019**, *35*, 1052–1056. <https://doi.org/10.1080/14786419.2019.1614581>.
150. Li, G.-Y.; Li, B.-G.; Yang, T.; Liu, G.-Y.; Zhang, G.-L. Secondary Metabolites from the Fungus *Chaetomium brasiliense*. *Helvetica Chim. Acta* **2008**, *91*, 124–129. <https://doi.org/10.1002/hlca.200890002>.
151. Huang, J.-Q.; Liao, Y.-C.; Chen, H.-J.; Zhang, Z. Chemical solution is an efficient method to induce the formation of 2-(2-phenylethyl) chromone derivatives in *Aquilaria sinensis*. *Phytochem. Lett.* **2017**, *19*, 64–70. <https://doi.org/10.1016/j.phytol.2016.12.004>.
152. Durazzo, A. Study Approach of Antioxidant Properties in Foods: Update and Considerations. *Foods* **2017**, *6*, 17. <https://doi.org/10.3390/foods6030017>.
153. Durazzo, A. Extractable and Non-extractable polyphenols: An overview. In *Non-extractable Polyphenols and Carotenoids: Importance in Human Nutrition and Health (Food Chemistry, Function and Analysis, Volume 5)*; Saura-Calixto, F., Perez-Jimenez, J., Eds.; The Royal Society of Chemistry: London, UK, 2018.
154. Durazzo, A.; Lucarini, M. Extractable and Non-Extractable Antioxidants. *Molecules* **2019**, *24*, 1933. <https://doi.org/10.3390/molecules24101933>.
155. Grazul, M.; Kufelnicki, A.; Wozniczka, M.; Lorenz, I.-P.; Mayer, P.; Jozwiak, A.; Czyz, M.; Budzisz, E. Synthesis, structure, electrochemical properties, cytotoxic effects and antioxidant activity of 5- amino-8-methyl-4H-benzopyran-4-one and its copper (II) complexes. *Polyhedron* **2012**, *31*, 150–158.
156. Kang, J.; Sun, J.-H.; Zhou, L.; Ye, M.; Han, J.; Wang, B.-R.; Guo, D.-A. Characterization of compounds from the roots of *Saposhnikovia divaricata* by high-performance liquid chromatography coupled with electrospray ionization tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* **2008**, *22*, 1899–1911. <https://doi.org/10.1002/rcm.3559>.
157. Kuroda, M.; Uchida, S.; Watanabe, K.; Mimaki, Y. Chromones from the tubers of *Eranthis cilicica* and their antioxidant activity. *Phytochemistry* **2009**, *70*, 288–293. <https://doi.org/10.1016/j.phytochem.2008.12.002>.
158. Park, Y.J.; Kim, H.J.; Lee, S.J.; Choi, H.-Y.; Jin, C.; Lee, Y.S. A New Chromone, 11-Hydroxy-sec-O-glucosylhamaudol from *Ostericum koreanum*. *Chem. Pharm. Bull.* **2007**, *55*, 1065–1066. <https://doi.org/10.1248/cpb.55.1065>.
159. Phosrithong, N.; Samee, W.; Nunthanavanit, P.; Ungwitayatorn, J. In Vitro Antioxidant Activity Study of Novel Chromone Derivatives. *Chem. Biol. Drug Des.* **2012**, *79*, 981–989. <https://doi.org/10.1111/j.1747-0285.2012.01368.x>.
160. Proença, C.; Albuquerque, H.M.T.; Ribeiro, D.; Freitas, M.; Santos, C.M.M.; Silva, A.M.S.; Fernandes, E. Novel chromone and xanthone derivatives: Synthesis and ROS/RNS scavenging activities. *Eur. J. Med. Chem.* **2016**, *115*, 381–392. <https://doi.org/10.1016/j.ejmech.2016.03.043>.
161. Yimam, M.; Brownell, L.; Jia, Q. Aloesin as a medical food ingredient for systemic oxidative stress of diabetes. *World J. Diabetes* **2015**, *6*, 1097–1107. <https://doi.org/10.4239/wjd.v6.i9.1097>.
162. Csepanyi, E.; Szabados-Furjesi, P.; Kiss-Szikszai, A.; Frensemeier, L.M.; Karst, U.; Lekli, I.; Haines, D.D.; Tosaki, A.; Bak, I. Antioxidant Properties and Oxidative Transformation of Different Chromone Derivatives. *Molecules* **2017**, *22*, 588. <https://doi.org/10.3390/molecules22040588>.
163. Demetgül, C.; Beyazit, N. Synthesis, characterization and antioxidant activity of chitosan-chromone derivatives. *Carbohydr. Polym.* **2018**, *181*, 812–817. <https://doi.org/10.1016/j.carbpol.2017.11.074>.
164. Valentina, P.; Ilango, K.; Chander, S.; Murugesan, S. Design, synthesis and α -amylase inhibitory activity of novel chromone derivatives. *Bioorg. Chem.* **2017**, *74*, 158–165. <https://doi.org/10.1016/j.bioorg.2017.07.018>.
165. Dai, H.F.; Liu, J.; Han, Z.; Zeng, Y.B.; Wang, H.; Mei, W.L. Two new 2-(2-phenylethyl) chromones from Chinese aglewood. *J. Asian Nat. Prod. Res.* **2010**, *12*(2), 134–137.
166. Yu, M.; Liu, Y.; Feng, J.; Chen, D.; Yang, Y.; Liu, P.; Yu, Z.; Wei, J. Remarkable Phytochemical Characteristics of Chi-Nan Agarwood Induced from New-Found Chi-Nan Germplasm of *Aquilaria sinensis* Compared with Ordinary Agarwood. *Int. J. Analytical Chemistry* **2021**, *2021*, 1–10. <https://doi.org/10.1155/2021/5593730>.
167. Kuang, T.D.; Chen, H.Q.; Kong, F.D.; Cai, C.H.; Yang, L.; Mei, W.L.; Dai, H.F. Three new 2-(2-phenylethyl)chromone derivatives from artificial holing agarwood of *Aquilaria sinensis*. *Phytochem Lett.* **2018**, *26*, 96–100.
168. Tian, H.; Wang, H.; Yang, L.; Gai, C.J.; Dong, W.H.; Li, W.; Mei, W.L.; Dai, H.F. Two new sesquiterpenoids from agarwood originated from *Aquilaria* sp. *J. Asian Nat. Prod. Res.* **2020**, *22*, 626–631.
169. Ueda, J.Y.; Imamura, L.; Tezuka, Y.; Tran, Q.L.; Tsudab, M.; Kadotaa, S. New sesquiterpene from Vietnamese agarwood and its induction effect on brain-derived neurotrophic factor mRNA expression in vitro. *Bioorg. Med. Chem.* **2006**, *14*, 3571–3574.
170. Shibata, S.; Sugiyama, T.; Uekusa, Y.; Masui, R.; Narukawa, Y.; Kiuchi, F. Five new 2-(2-phenylethyl)chromone derivatives from agarwood. *J Nat Med.* **2020**, *74*, 561–570. doi: 10.1007/s11418-020-01410-z.
171. Yang, D.L.; Li, W.; Dong, W.H.; Wang, J.; Mei, W.L.; Dai, H.F. Five new 5,11-epoxyguaiane sesquiterpenes in agarwood “qi-nan” from *Aquilaria sinensis*. *Fitoterapia* **2016**, *112*, 191–196.
172. Li, W.; Mei, W.L.; Dong, W.H.; Cai, C.H.; Gai, C.J.; Dai, H.F. Study on chemical constituents of Chinese agarwood “Qi-Nan”. *J. Trop. Subtrop. Bot.* **2019**, *27*, 196–202.
173. Nakanishi, T.; Yamagata, E.; Yoneda, K.J. Jinkoh-eremol and jinkohol II, two new sesquiterpene alcohols from agarwood. *Chem. Soc. Perkin Trans.* **1983**, *1*, 601–604.

174. Nakanishi, T.; Yamagata, E.; Yoneda, K.; Miura, I. Jinkohol, a prezizane sesquiterpene alcohol from agarwood. *Phytochemistry* **1981**, *20*, 1597–1599.
175. Ma, C.T.; Ly, T.L.; Van Le, T.H.; Anh Tran, T.V.; Kwon, S.W.; Park, J.H. Sesquiterpene derivatives from the agarwood of *Aquilaria malaccensis* and their anti-inflammatory effects on NO production of macrophage RAW 264.7 cells. *Phytochemistry*, 2021, *183*, 112630. <https://doi.org/10.1016/j.phytochem.2020.112630>.
176. Chen, D.; Bi, D.; Song, Y.L.; Tu, P.F. Flavonoids from the stems of *Aquilaria sinensis*. *Chin. J. Nat. Med.* **2012**, *10*, 287–291.
177. Ma, C.T.; Cho, E.; Nguyen, H.T.; Wu, B.; Le, T.H.V.; Oh, K.B.; Kwon, S.W.; Nguyen, M.D.; Park, J.H. Malacinones A and B, two novel sesquiterpenoids with 6/6/5 tricyclic ring system from the agarwood of *Aquilaria malaccensis*. *Tetrahedron Lett.* **2020**, *61*, 151355.
178. Yang, L.; Qiao, L.-R.; Zhang, J.-J.; Dai, J.-G.; Guo, S.-X. Two new sesquiterpene derivatives from Chinese eaglewood. *J. Asian Nat. Prod. Res.* **2012**, *14*, 1054–1058. <https://doi.org/10.1080/10286020.2012.704910>.
179. Dahham, S.S.; Tabana, Y.M.; Iqbal, M.A.; Ahamed, M.B.K.; Ezzat, M.O.; Majid, A.S.A.; Majid, A.M.S.A. The Anticancer, Antioxidant and Antimicrobial Properties of the Sesquiterpene β -Caryophyllene from the Essential Oil of *Aquilaria crassna*. *Molecules* **2015**, *20*, 11808–11829.
180. Yang, L.; Yang, Y.L.; Dong, W.H.; Li, W.; Wang, P.; Cao, X.; Yuan, J.Z.; Chen, H.Q.; Mei, W.L.; Dai, H.F. New tricyclic prezizaane sesquiterpenoids from agarwood. *J. Enzym. Inhib. Med. Chem.* **2019**, *34*, 853–862.
181. Mei, W.L.; Lin, F.; Zuo, W.J.; Wang, H.; Dai, H.F. Cucurbitacins from fruits of *Aquilaria sinensis*. *Chin. J. Nat. Med.* **2012**, *10*, 234–237.
182. Adhikari, S.R.; Pokhrel, K.; Baral, S.D. Economic Value of Agarwood and Its Prospects of Cultivation. *Int. J. Appl. Sci. Biotechnol.* **2021**, *9*, 23–31. <https://doi.org/10.3126/ijasbt.v9i1.35984>.
183. Swee, L.L.C. *Agarwood (Aquilaria malaccensis) in Malaysia, International Expert Workshop on CITES Non-Detriment Findings*; Publisher: Cancun, Mexico, 2008.
184. Compton, J.; Ishihara, A. *The Use and Trade of Agarwood in Japan*; TRAFFIC International: Cambridge, UK, 2004.
185. Antonopoulou, M.; Compton, J.; Perry, L.S.; Al-Mubarak, R. *The Trade and Use of Agarwood (Oudh) in the United Arab Emirates*; TRAFFIC Southeast Asia: Petaling Jaya, Selangor, Malaysia, 2010.
186. Kiet, L.K. History and ecology of agarwood in Vietnam. In Proceedings of the 1st International Agarwood Conference, Ho Chi Minh City, Vietnam, 10–15 November 2003.
187. Lim, T.W.; Anak, N.A. *Wood for the Trees: A Review of the Agarwood (Gaharu) Trade in Malaysia*; TRAFFIC Southeast Asia: Petaling Jaya, Selangor, Malaysia, 2010.
188. Persoon, G.A. Growing ‘the Wood of The Gods’: Agarwood production in Southeast Asia. In *Smallholder Tree Growing for Rural Development and Environmental Services*; Snelder, D.J., Lasco, R.D., Eds.; Springer Science Business Media: Berlin, Germany, 2008; pp. 245–262. https://doi.org/10.1007/978-1-4020-8261-0_12.
189. Sitepu, I.R.; Santoso, E.; Siran, S.A.; Turjaman, M. *Fragrant Wood Gaharu: Whenthe Wild Can No Longer Provide. Forestry Research and Development Agency (FORDA)*; International Tropical Timber Organization: Yokohama, Japan, 2011.
190. CITES. *The Trade and Use of Agarwood in Taiwan, Province of China*; TRAFFIC East Asia: Taipei, Taiwan, 2005. Available online: <http://cites.org/sites/default/files/common/com/pc/15/X-PC15-07-Inf.pdf> (accessed on 18 May 2022).
191. LaFrankie, J.V. Population dynamics of some tropical trees that yield nontimber forest products. *Econ. Bot.* **1994**, *48*, 301–309.
192. Babatunde, O.J. Oud: Arabia’s Traditional Scent. 2015. Available online: <http://www.masterpiece-ng.com/2015/09/01/oud-arabias-traditional-scent/> (18 May 2022).
193. Abdin, M.J. The agar wood industry: Yet to utilize in Bangladesh. *Int. J. Econ. Manag. Sci.* **2014**, *3*, 163–166. <https://doi.org/10.2139/ssrn.2430055>.