

Article

An In Situ Formation of Ionic Liquid for Enrichment of Triazole Fungicides in Food Applications Followed by HPLC Determination

Rawikan Kachangoon ¹, Jitlada Vichapong ^{1,2,*}, Yanawath Santaladchaiyakit ³ and Supalax Srijaranai ⁴

¹ Creative Chemistry and Innovation Research Unit, Department of Chemistry and Center of Excellent for Innovation in Chemistry, Faculty of Science, Mahasarakham University, Maha Sarakham 44150, Thailand; r.kachangoon@hotmail.com

² Multidisciplinary Research Unit of Pure and Applied Chemistry (MRUPAC), Department of Chemistry and Center of Excellent for Innovation in Chemistry, Faculty of Science, Mahasarakham University, Maha Sarakham 44150, Thailand

³ Department of Chemistry, Faculty of Engineering, Rajamangala University of Technology Isan, Khon Kaen Campus, Khon Kaen 40000, Thailand; sanyanawa@gmail.com

⁴ Materials Chemistry Research Center, Department of Chemistry and Center of Excellent for Innovation in Chemistry, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand; supalax@kku.ac.th

* Correspondence: jitlada.v@msu.ac.th

Abstract: An in situ formation of ionic liquid was used for preconcentration of four triazole fungicides in food samples. The microextraction method was used for the first time in the literature for preconcentration of triazole fungicides. In the developed method, tributylhexadecylphosphonium bromide ([P₄₄₄₁₂]Br) and potassium hexafluorophosphate (KPF₆) were used for the formation of hydrophobic ionic liquid. After centrifugation, the fine microdroplets were produced in one step, providing the extraction step in a quick and environmentally friendly manner. The functional group of the hydrophobic ionic liquid was investigated using FT-IR. Various extraction parameters were studied and optimized. In the extraction method, 0.01 g of [P₄₄₄₁₂]Br and 0.01 g of KPF₆, centrifugation at 4500 rpm for 10 min were used. The optimized technique provided a good linear range (90–1000 µg L⁻¹) and high extraction recovery, with a low limit of detection (30–50 µg L⁻¹). Methods for the proposed in situ formation of ionic liquid were successfully applied to honey, fruit juice, and egg matrices. The recoveries were obtained in a satisfactory range of 62–112%. The results confirmed the suitability of the proposed microextraction method for selective extraction and quantification of triazole fungicides.

Keywords: ionic liquid; in situ extraction; triazole fungicides; extraction; HPLC

Citation: Kachangoon, R.; Vichapong, J.; Santaladchaiyakit, Y.; Srijaranai, S. An In Situ Formation of Ionic Liquid for Enrichment of Triazole Fungicides in Food Applications Followed by HPLC Determination. *Molecules* **2022**, *27*, 3416. <https://doi.org/10.3390/molecules27113416>

Academic Editor: Branimir Pavlic

Received: 2 May 2022

Accepted: 23 May 2022

Published: 25 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Triazole fungicides (TFs) are a class of extremely efficient systemic fungicides that contain a hydroxyl group (ketone group), a substituted phenyl group, and a 1,2,4-triazole group in the main chain [1]. The TFs are promoted with broad-spectrum, internal absorption characteristics, providing good stability for a long time [2]. They are commonly applied in the cultivation and postharvest storage of crops to protect them from fungal infections [3,4]. However, they may disturb endocrine activities, and have induced developmental toxicity in animals and human health [5]. For this reason, a reliable and effective method is still needed for the determination of triazole fungicides in various matrices.

Chromatographic methods such as liquid chromatography [6–10] and gas chromatography [11–15] have been used for the simultaneous analysis of triazole fungicides. Due to the trace levels of triazole fungicides in complex matrices, some preconcentration/extraction procedures are necessary prior to instrumental analysis. The advancement of

sample-preparation methods has mainly promoted their simplification, miniaturization (reduction in the amounts of sample and extraction solvent), and environmental friendliness (use of an alternative extraction solvent) [4]. Recently, miniaturized techniques have been advanced to enrich triazoles in various matrices, such as ultrasonic-assisted dispersive solid-phase extraction (UA-DSPE) [16], magnetic solid-phase extraction (MSPE) [6], homogeneous liquid–liquid extraction based on salting out (SHLLE) [14], vortex-assisted ionic liquid dispersive liquid–liquid microextraction (VA-IL-DLLME) [9], aqueous two-phase extraction (ATPE) [17], and microsolid-phase extraction [18].

Room temperature ionic liquids (RTILs), which are defined as “alternative” solvents, are promising and suitable solvents that have been recently used as replacement extraction solvents for sample preparation and preconcentration methods [19]. They have some unique attributes, such as a negligible vapor pressure, high thermal stability, good extractability for various organic compounds and metal ions as neutral or charged complexes, and tunable viscosity and miscibility with water and organic solvents [20]. The RTIL types are alkylammonium, tetraalkylammonium, tetraalkylphosphonium, 1,3-dialkylimidazolium, and N-alkylpyridinium salts formed with weak nucleophilic anions such as hexafluorophosphate, tetrafluoroborate, and perfluoroalkylsulfonate [21]. There are some reports on sample preconcentration using RTILs [19–22]. Some of extraction methods were based on liquid–liquid microextraction and dispersive liquid–liquid microextraction (DLLME). Generally in the DLLME process, a disperser solvent is required to induce the mass transfer of the extraction solvents into an aqueous solution [23]. In situ DLLME was developed to avoid the use of these dispersive solvents [24,25]. In this method, a hydrophilic RTIL was used as a starting extraction solvent capable of completely dissolving in the aqueous analyte solution. Then, an in situ metathesis reaction was used to form an RTIL immiscible with water by adding an ion-exchange reagent into the aqueous solution to complete the extraction process. This exchange process formed a cloudy solution with fine microdroplets, which exceedingly improved the surface area of the RTIL, resulting in the high extraction efficiency of the target analytes. Due to the high price of RTILs, the metathesis reaction and extraction were investigated in a single step, making the transfer of analyte into the extracting phase very quick and efficient. The benefits of the in situ metathesis reaction-assisted IL were the rejection of the requirement for using a disperser solvent and a remarkable increase in the surface area of the IL extraction solvent.

In this study, an in situ metathesis reaction that generated an ionic liquid was combined with a liquid–liquid microextraction for the preconcentration of triazole fungicides prior to their analysis using high-performance liquid chromatography. A tributyldecylphosphonium bromide ($[P_{44412}]\text{Br}$) as the starting extraction solvent and an ion-exchange reagent (KPF_6) were used to produce a hydrophobic ionic liquid (IL) ($[P_{44412}][\text{PF}_6]$). The starting extraction solvent could be completely dissolved in the aqueous solution, which promoted analyte extraction in the absence of a dispersion solvent. In this work, an in situ metathesis reaction was performed to yield a cloudy solution with fine microdroplets, which significantly increased the extraction efficiency. After centrifugation, the extraction solvent was easily collected with a syringe after solidification. Using in situ formation, the fine microdroplets were produced in one step, providing a short extraction time and environmental friendliness. Various experimental parameters that affected the extraction efficiency of the method were investigated and optimized. Finally, the selected extraction conditions were used to determine triazole fungicide residues in water, honey, fruit juice, and egg matrices.

2. Results and Discussion

2.1. Characterization of In Situ Ionic Liquid

The halide anion of tributylhexadecylphosphonium bromide ($[P_{44412}]Br$) and potassium hexafluorophosphate (KPF_6) was the main force for the formation of the hydrophobic ionic liquid. FT-IR spectra were used to confirm the formation of hydrogen bonding, as shown in Figure 1. In the FT-IR spectra, the characteristic peaks presented at 2850, 2920, and 2960 cm^{-1} were assigned to the C-H stretching or CH_3 stretching, while those at 1410 and 1465 cm^{-1} were attributed to the C-H bending and CH_3 bending vibrations of pure $[P_{44412}]Br$ [26]. Moreover, the FT-IR spectra of the characteristic peaks showed the halogen compound vibrations at 577 and 795 cm^{-1} . This may have been due to a transfer of the bromide ion cloud electron to hydrogen bonding, and consequently, a decrease in the force constant [27]. Thus, the shift of the hexafluorophosphate counter-ion vibrations suggested the existence of hydrogen bonding between $[P_{44412}]Br$ and KPF_6 when the hydrophobic ionic liquid was formed.




Figure 1. FT-IR spectra of (a) KPF_6 , (b) $[P_{44412}]Br$, and (c) the in situ ionic liquid when it was formed.

2.2. Optimization of In Situ Metathesis-Reaction-Generated Ionic Liquid Combined with Liquid-Liquid Microextraction

Due to their low concentrations and matrix interferences in real samples, it is difficult to directly analyze triazole fungicides. Therefore, a sample-preparation method is necessary before analysis. In this work, an in situ metathesis reaction that generated the ionic liquid was combined with a liquid-liquid microextraction in the triazole fungicide analysis. In order to obtain a high extraction efficiency, various parameters had to be investigated. The optimization was carried out using the aqueous solution (10 mL) containing 100 $\mu g L^{-1}$ of each triazole fungicides. All experiments were performed at least in triplicate.

In order to form the ionic liquid ($[P_{44412}][PF_6]$), $[P_{44412}]Br$ and KPF_6 were selected because the starting extraction solvent could be completely dissolved in the aqueous solution, which promoted the analyte extraction in the absence of a dispersion solvent; the melting of this IL ($[P_{44412}][PF_6]$) was between 10 and 30 °C, and it could solidify at low temperatures [24]. Moreover, $[P_{44412}][PF_6]$ is denser than water, therefore it was easy to collect as the bottom layer after centrifugation. The amount of $[P_{44412}]Br$ was studied in the range of 0.01–0.10 g. The results are shown in Figure 2. The results showed that the maximum peak area was obtained at 0.01 g of $[P_{44412}]Br$. Therefore, 0.01 g of $[P_{44412}]Br$ was chosen.

The amount of KPF_6 varied in a range of 0.01–0.10 g. The results are shown in Figure 3. The results showed that the maximum peak area was obtained at 0.01 g of KPF_6 . Then, the signal decreased at a higher amount of KPF_6 . Thus, 0.01 g of KPF_6 was selected. The molar ratio of $[P_{44412}]Br$ to KPF_6 was selected as 1:2.5 because the excess KPF_6 ensured 100% completion of the in situ metathesis reaction [24].




Figure 2. Effects of the amount of $[P_{44412}]Br$ on the extraction efficiency.




Figure 3. Effects of the amount of KPF₆ on the extraction efficiency.

Because the process of mass transfer and completely phase separation in an extraction procedure should be time-dependent, the effects of extraction speed and time on the peak area was studied [22]. The effect of centrifugation speed was evaluated in the range of 1500–5000 rpm (see Figure 4). It was found that the peak areas of all the analytes increased up to a 4500 rpm centrifugation speed, above which the peak areas slightly decreased due to disintegration of the phase of target analytes. Therefore, a centrifugation speed of 4500 rpm was selected. The extraction times studied ranged from 3 to 15 min, while other experimental conditions were kept constant (data not shown). The peak areas of most of the triazoles increased with increases in the extraction time, and reached the highest at 10 min. Therefore, 10 min was chosen to ensure an efficient extraction.




Figure 4. Effects of centrifugation speed (rpm) on the extraction efficiency.

Before HPLC analysis, acetonitrile (ACN) was added to dissolve the RTIL-rich phase due to its solubility property in the $[P_{44412}]PF_6$ phase and the compatibility with the mobile phase being used. The ACN volume was investigated in the range of 30–250 μL (as shown in Figure 5). It was found that ACN 200 μL provided the highest peak areas of all analytes. After that, the peak area decreased due to the dilution effect. When using ACN at less than 30 μL , the phase could not be completely dissolved. Thus, 200 μL of ACN was chosen.





Figure 5. Effects of dissolving solvent (acetonitrile, μL) on the extraction efficiency.

2.3. Analytical Performance of the Proposed Method

Under the chosen extraction condition, a series of experiment were conducted to study the enrichment factors (EFs), extraction recoveries (ERs), linear ranges, limits of detection (LODs), limits of quantitation (LOQs), and precisions (intraday and interday). The results are shown in Table 1. The calibration graphs were linear over concentration ranges of 90–1000 $\mu\text{g L}^{-1}$ for myclobutanil, triadimefon, and tebuconazole, and 150–1000 $\mu\text{g L}^{-1}$ for hexaconazole, with a coefficient of determination in the range of 0.9988–0.9991. The LODs were evaluated as the concentrations giving a signal-to-noise ratio of 3 ($S/N = 3$), and were in the range of 30–50 $\mu\text{g L}^{-1}$. The LOQs ($S/N = 10$) were in the range of 90–150 $\mu\text{g L}^{-1}$. The relative standard deviation (RSD) was determined using five solutions of 150 $\mu\text{g L}^{-1}$ of each triazole fungicide. The RSD values of the retention times and peak areas were in the ranges of 0.40–0.80% and 4.84–6.23%, respectively. The EFs were in the range of 8.53–12.26. The ERs were in the range of 59.71–85.82. The chromatograms obtained from the proposed preconcentration method and direct analysis (without preconcentration) are shown in Figure 6. It was found that the developed method showed high chromatographic signals when compared to the direct analysis.

Table 1. Analytical performance of the proposed method.

Analyte	Linear Range ($\mu\text{g L}^{-1}$)	Linear Equation	R^2	LOD ($\mu\text{g L}^{-1}$)	LOQ ($\mu\text{g L}^{-1}$)	Intraday Precision (n = 5),		Interday Precision		EF	ER
						RSD (%)		(n = 5), RSD (%)			
						t_R	Peak Area	t_R	Peak Area		
Myclobutanil	90–1000	$y = 1 \times 10^6x + 9995.5$	0.9994	30	90	0.61	2.48	0.77	2.70	12.26	85.82
Triadimefon	90–1000	$y = 907,260x + 11,705$	0.9989	30	90	0.51	2.92	0.80	3.50	8.53	59.71
Tebuconazole	90–1000	$y = 484,103x + 4328.4$	0.9991	30	90	0.40	2.60	0.46	2.72	11.01	77.07
Hexaconazole	150–1000	$y = 689,281x + 3751.6$	0.9988	50	150	0.70	2.55	0.77	2.60	11.34	79.38

**Figure 6.** Chromatograms of standard neonicotinoid insecticides obtained (a) without preconcentration and (b) with preconcentration using the proposed microextraction method; the concentration of all standards was $300 \mu\text{g L}^{-1}$.


2.4. Analysis of Real Samples

To study the applicability of the proposed method, an in situ formation and preconcentration method using the ionic liquid was applied in the determination of triazole fungicides in honey, fruit juice, and egg samples. A standard addition method was applied to study the matrix effect (ME). ME(%) is expressed as the ratio of the slopes obtained from calibration curves of each analyte spiked into the samples to those obtained after extraction using the proposed method, according to Equation (1) [28]:

$$ME(\%) = \frac{\text{slope of spiked real sample}}{\text{slope of standard solution}} \times 100 \quad (1)$$

In the absence of the ME, the slope of the calibration curves from both the standard solution and spiked real samples should be similar ($ME(\%) \approx 100\%$). However, in the presence of ME, the signal intensity for the analytes can decrease or increase. Generally, ME values between 80 and 120% indicate no ME, ME values between 50 and 80% or 120 and 150% indicate minor MEs, and ME values less than 50% or greater than 150% indicate major MEs [29]. It was found that the MEs (%) were in the range of 76.9–113.9%.

In order to confirm the accuracy of the proposed method, the relative recoveries (RRs) were investigated using an analysis of three real samples spiked with five triazole fungicides at a concentration of $150 \mu\text{g}\cdot\text{L}^{-1}$ within one day. As shown in Table 2, acceptable recoveries (62–112%) with relative standard deviations (RSDs) of less than 7.9% were obtained. The obtained results showed that no triazole fungicide residues were detected in all samples. These results confirmed that the proposed microextraction method could successfully be utilized to estimate triazole fungicide residues at trace levels in real samples with high accuracy and validity. The chromatograms of blank and spiked samples are shown in Figure 7.






Figure 7. Chromatograms of sample and spiked sample ($150 \mu\text{g L}^{-1}$ for fruit juice sample and $150 \mu\text{g kg}^{-1}$ for honey and egg samples); (a) honey sample, (b) passion fruit juice sample, (c) pomegranate juice sample and (d) egg yolk sample.

Table 2. Recoveries obtained from the analysis of triazole fungicides in real samples (n = 3).

Sample	MCBT		TDF		TBZ		HCZ	
	%RR *	%RSD **	%RR	%RSD	%RR	%RSD	%RR	%RSD
Honey I	66.17	0.94	92.41	2.82	99.72	1.99	94.12	1.06
Honey II	86.98	2.47	72.03	3.12	60.26	2.60	62.22	2.57
Passion fruit juice	76.14	0.84	77.87	1.80	81.50	1.82	87.63	1.20
Pomegranate juice	72.14	0.71	61.77	1.78	69.19	0.20	62.98	1.53
Grape juice	72.50	1.82	62.65	1.39	66.16	0.87	63.59	2.87
Egg yolk	84.19	0.31	72.72	0.47	69.41	1.69	69.05	1.08

* RR: relative recovery; ** RSD: relative standard deviation.

2.5. Comparison of the Proposed Microextraction Method with Other Sample-Preparation Methods

Table 3 shows a comparison of the developed microextraction in this work with other published methods [1,4,29,30]. The ionic liquid could be prepared under facile conditions, which simplified the experimental operation, and no complicated instrument was required. Compared to the other sample-preparation methods, the proposed method provided a wide linear calibration range (150–1000 $\mu\text{g L}^{-1}$), a high precision (less than 5), and acceptable recoveries (61–112) within shorter extraction times for simultaneous extraction determinations for various samples. In conclusion, the proposed microextraction method was demonstrated as a simple, fast, effective, and environmentally friendly technique.

Table 3. Comparisons of the proposed method with other methods for the quantitation of triazole fungicides.

Extraction Method **	Analytical Method	Samples	Adsorbent/Desorbed ***	Linear Range ($\mu\text{g L}^{-1}$)	Recovery (%)	LOD ($\mu\text{g L}^{-1}$ or $\mu\text{g Kg}^{-1}$)	EF	%RSD	Ref.
SMMH-d-SPME	HPLC-UV	Water	$\text{Fe}_3\text{O}_4/\text{SiO}_2/\text{ACN}$	5–100 and 2.5–50	90–104	1.0–2.5	40–237	Less than 8%	[4]
MSPE	HPLC-UV	River water	BCDP/ACN – 0.1% (v/v) HCOOH	1–1200	82.8–113.2	0.2–0.3	281–283	1.2–4.6	[29]
SBSE	HPLC-DAD	Grape and cabbage Water	HC-POF/ACN-water	0.1–500	80.7–111	0.022–0.071	49–57	6.4–12.4	[1]
MSPE	LC-MS/MS	and fruit juices	GO- PmAP/ACN –water in NaOH	0.001–0.5	80.3–106.3	0.08–2.04 ng g^{-1}	0.45–0.60	2.1–13.4	[30]
Ionic liquid combined with liquid-liquid microextraction	HPLC-DAD	Honey, fruit juices, and egg yolk	IL-ACN	150–1000	61–112	30–50	8–11	≤ 5	This work

** MSPE-HPLC-UV: magnetic solid-phase extraction–high-performance liquid chromatography–ultraviolet detector; SBSE-HPLC-DAD: stir bar sorption extraction–high-performance liquid chromatography–diode array detector; SPME-HPLC-DAD: solid-phase microextraction–high-performance liquid chromatography–diode array detector; MSPE-LC-MS/MS: magnetic solid-phase extraction–liquid chromatography–tandem mass spectrometry; μSPE -HPLC-PDA: micro solid-phase extraction–high-performance liquid chromatography–photodiode array; SMMH-d-SPME: in situ surfactant-mixed metal hydroxide (SMMH); d-SPME: dispersive solid-phase microextraction; ***

Fe₃O₄@SiO₂: magnetic nanoparticle; ACN: acetonitrile; BCDP: bridged *bis*CD-bonded chiral column; HCOOH: formic acid; HC-POF: hydroxyl-containing porous organic framework; GO-PmAP: graphene oxide-poly 3-aminophenol.

3. Materials and Methods

3.1. Chemicals and Reagents

Four triazole herbicides (myclobutanil (C₁₅H₁₇ClN₄), triadimefon (C₁₄H₁₆ClN₃O₂), tebuconazole (C₁₆H₂₂ClN₃O), and hexaconazole (C₁₄H₁₇Cl₂N₃O)) from Dr. Ehrenstorfer GmbH (Augsburg, Germany) were used. The stock solution of each fungicide (1000 mg L⁻¹) was prepared using methanol and used for further dilutions with water. Potassium hexafluorophosphate (KPF₆) (Sigma-Aldrich, Beijing, China) and tributyldecylphosphonium bromide ([P₄₄₄₁₂]⁺Br⁻) (Sigma-Aldrich, Schaffhausen, Switzerland) were used. Methanol and acetonitrile (HPLC-grade) were obtained from Merck (Darmstadt, Germany). Deionized water with a resistivity of 18.2 MΩ.cm was obtained from a Type 1 Simplicity[®] ultrapure water system (Merck, Darmstadt, Germany). All solutions were filtered through a 0.45 µm filter before being injected into the HPLC system. All chemicals and reagents mentioned above were of analytical grade.

3.2. Instrumentations

The HPLC analysis was conducted on a Waters 1525 Binary HPLC pump (Water, Massachusetts, USA) equipped with a diode array detector (DAD). A Rheodyne injector with a 20 µL injection loop was used. Empower 3 software was used to acquire and analyze the chromatographic data. A Purospher[®] STAR RP-18 endcapped (4.6 × 150 mm, 5 µm) column (Merck, Darmstadt, Germany) with an isocratic elution of acetonitrile and water at a ratio of 50:50 (%v/v) was used for the separations. The mobile phase flow rate was 1 mL min⁻¹. The detection wavelength was set at 220 nm.

The Fourier-transformed infrared spectra (FTIR) of the ionic liquid phase were measured using a Bruker Invenio-S FT-IR (Bruker Corp., Massachusetts, USA). Diamond-lens-attenuated total reflectance (ATR) was used. A centrifuge (Centurion, Aberdeen, England) also was used.

3.3. Sample Preparation

3.3.1. Honey Samples

Honey samples were purchased from a supermarket in Maha Sarakham province. A total of 5 g of the sample was weighed into a 50 mL volumetric flask and diluted to the marker. The solution was filtered through a Whatman (no. 1) filter paper. After that, the filtrate was passed through a 0.45 µm nylon membrane filter before extraction using the proposed method.

3.3.2. Fruit Juice Samples

Passion fruit juice and pomegranate juice (commercial juice samples) were bought from the supermarket in Maha Sarakham province. An aliquot of fruit juice (30.0 mL) was centrifuged at 4000 rpm for 10 min and then filtered through a Whatman (no. 1) filter paper. The solution was then passed through a 0.45 µm nylon membrane filter before extraction using the proposed method.

3.3.3. Egg Yolk Sample

Chicken eggs were purchased from local markets in Maha Sarakham province. The yolk was separated from the white to reduce interference, since in the analysis of egg collected from animals treated with anthelmintics, it is known that the interferences are greater in the yolk [31,32]. Fortification of the sample was performed directly in the yolk, and a period of about 12 h was allowed to elapse before continuing with any of the extraction processes in order to improve the interaction between the analytes and the matrix

compounds [33]. A total of 10.00 g of egg yolk was mixed well with 0.2 g of anhydrous Na_2SO_4 . After that, 1% (v/v) acetic acid in acetonitrile (2.00 mL) was added and shaken vigorously by hand for 1 min, and the homogenized eggs were centrifuged at 3500 rpm for 5 min for complete fat and protein precipitation. The supernatants were collected using a microsyringe. The solutions were diluted with deionized water to 10.00 mL, 100 μL of acetic acid was added, and the solutions were centrifuged to ensure the complete precipitation of fat and proteins [32]. The samples were spiked with the triazole fungicides at different concentrations before fat and protein precipitation. The obtained clear solutions were then extracted using the proposed microextraction method.

3.4. In Situ Metathesis-Reaction-Generated Ionic Liquid Combined with Liquid–Liquid Microextraction

A schematic diagram of the microextraction procedure is shown in Figure 8. The mixed solution included the standard or sample solution (10.00 mL), 0.01 g of $[\text{P}^{44412}]\text{Br}$ and 0.01 g of KPF_6 , which were added into a 15 mL conical centrifuge tube. Then, the tube was shaken to dissolve the $[\text{P}^{44412}]\text{Br}$ and KPF_6 and to complete the in situ metathesis reaction. The solution was then centrifuged at 4500 rpm for 10 min. After that, the centrifuge tube was cooled in an ice bath until the $[\text{P}^{44412}]\text{PF}_6$ was generated. The $[\text{P}^{44412}]\text{PF}_6$ phase (upper phase) was separated and then diluted with acetonitrile (200 μL) to decrease the viscosity before being injected into the HPLC system.




Figure 8. Schematic illustration of the microextraction procedure using in situ formation of the ionic liquid for triazole fungicides and HPLC analysis.

3.5. Evaluation of Enrichment Factor (EF), Extraction Recovery (ER), Relative Recovery (RR), and Matrix Effect (ME)

To study the effects of experimental conditions on the extraction efficiency, the EF was calculated between the analyte concentration in the final phase (C_{final}) and the initial concentration in the analyte in aqueous sample solution (C_0) according to the following equations:

$$\text{EF} = C_{\text{final}}/C_0 \quad (2)$$

The %ER was expressed as the total percentage amount of the target analytes extracted into the sediment phase using the proposed microextraction method:

$$\%ER = EF \times (V_{sed}/V_0) \times 100 \quad (3)$$

where V_{sed} and V_0 are the volume of sediment phase and the sample solution, respectively.

The %RR was defined as the % amount of analyte recovered from the matrix (real samples) with reference to the extracted standard (standard spiked into the same matrix).

$$RR(\%) = \frac{C_{found} - C_{real}}{C_{added}} \times 100 \quad (4)$$

where C_{found} is the concentration of analyte after adding a known amount of working standard to the real sample, C_{real} is the analyte concentration in the real sample, and C_{added} represents the concentration of a known amount of working standard that was spiked into the real samples.

4. Conclusions

In the present study, an in situ extraction and preconcentration method using an ionic liquid for separation and preconcentration of triazole fungicides in honey, fruit juice, and egg samples was performed prior to high-performance liquid chromatographic analysis. The halide anion of tributylhexadecylphosphonium bromide ($[P_{44412}]Br$) and potassium hexafluorophosphate (KPF_6) was used for the formation of the hydrophobic ionic liquid. In the proposed microextraction method, forming the immiscible IL extraction phase and the transfer of analytes occurred simultaneously. The metathesis reaction and extraction were accomplished in single step, making the transfer of the analytes into the extracting phase very quick and efficient. The proposed method provided good repeatability, a wide linearity range, a high enrichment factor, and an acceptable extraction recovery for each compound, and matrix effects did not interfere with the quantification process. Therefore, the proposed method is recommended as a fast, simple, sensitive, and environmentally friendly sample-preparation technique.

Author Contributions: R.K., J.V., and Y.S. participated in designing the study. R.K. and J.V. conducted the study. Data were collected and analyzed by J.V. The manuscript was written by R.K., J.V., Y.S., and S.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research is supported by Mahasarakham University and Thailand Science Research and Innovation (TSRI) 2021 through grant number 6406019.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Acknowledgments: The authors gratefully acknowledge financial support for this research from Thailand Science Research and Innovation (TSRI) 2021, Mahasarakham University, and the Center of Excellence for Innovation in Chemistry (PERCH-CIC). We would like to thank the National Research Council of Thailand (NRCT; N41A640218). We also thank everyone for their assistance in the laboratory.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Wang, Y.; He, M.; Chen, B.; Hu, B. Hydroxyl-containing porous organic framework coated stir bar sorption extraction combined with high performance liquid chromatography-diode array detector for analysis of triazole fungicides in grape and cabbage samples. *J. Chromatogr. A* **2020**, *1633*, 461628.
2. Buerge, I.J.; Poiger, T.; Müller, M.D.; Buser, H.R. Influence of pH on the stereoselective degradation of the fungicides epoxiconazole and cyproconazole in soils. *Environ. Sci. Technol.* **2006**, *40*, 5443–5450.

3. Dong, F.S.; Li, J.; Chankvetadze, B.; Cheng, Y.P.; Xu, J.; Liu, X.G.; Li, Y.B.; Chen, X.; Bertucci, C.; Tedesco, D.; et al. Chiral triazole fungicide difenoconazole: Absolute stereochemistry, stereoselective bioactivity, aquatic toxicity, and environmental behavior in vegetables and soil. *Environ. Sci. Technol.* **2013**, *47*, 3386–3394.
4. Seebunrueng, K.; Tamuang, S.; Ruangchai, S.; Sansuk, S.; Srijaranai, S. In situ self-assembled coating of surfactant-mixed metal hydroxide on Fe₃O₄@SiO₂ magnetic composite for dispersive solid phase microextraction prior to HPLC analysis of triazole fungicides. *Microchem. J.* **2021**, *168*, 106396.
5. Han, X.; Chen, J.; Li, Z.; Quan, K.; Qiu, H. Magnetic solid-phase extraction of triazole fungicides based on magnetic porous carbon prepared by combustion combined with solvothermal method. *Anal. Chim. Acta* **2020**, *1129*, 85–97.
6. Liu, G.; Li, L.; Gao, Y.; Gao, M.; Huang, X.; Lv, J.; Xu, D. A beta-cyclodextrin-functionalized magnetic metal organic framework for efficient extraction and determination of prochloraz and triazole fungicides in vegetables samples. *Ecotoxicol. Environ. Saf.* **2019**, *183*, 109546.
7. Jing, X.; Yang, L.; Zhao, W.; Wang, F.; Chen, Z.; Ma, L.; Jia, L.; Wang, X. Evaporation-assisted dispersive liquid-liquid microextraction based on the solidification of floating organic droplets for the determination of triazole fungicides in water samples by high-performance liquid chromatography. *J. Chromatogr. A* **2019**, *1597*, 46–53.
8. Pang, J.; Mei, M.; Yuan, D.; Huang, X. Development of on-line monolith-based in-tube solid phase microextraction for the sensitive determination of triazoles in environmental waters. *Talanta* **2018**, *184*, 411–417.
9. Zhang, Y.; Zhang, Y.; Zhao, Q.; Chen, W.; Jiao, B. Vortex-assisted ionic liquid dispersive liquid-liquid microextraction coupled with high-performance liquid chromatography for the determination of triazole fungicides in fruit juices. *Food Anal. Methods* **2016**, *9*, 596–604.
10. Bordagaray, A.; Garcia-Arrona, R.; Millán, E. Determination of triazole fungicides in liquid samples using ultrasound-assisted emulsification microextraction with solidification of floating organic droplet followed by high-performance liquid chromatography. *Food Anal. Methods* **2014**, *7*, 1195–1203.
11. Wang, P.; Zhao, Y.; Wang, X.; Yu, G.W.; Wang, J.; Li, Z.G.; Lee, M.R. Microwave-assisted-demulsification dispersive liquid-liquid microextraction for the determination of triazole fungicides in water by gas chromatography with mass spectrometry. *J. Sep. Sci.* **2018**, *41*, 4498–4505.
12. Wei, Q.; Song, Z.; Nie, J.; Xia, H.; Chen, F.; Li, Z.; Lee, M. Tablet-effervescence-assisted dissolved carbon flotation for the extraction of four triazole fungicides in water by gas chromatography with mass spectrometry. *J. Sep. Sci.* **2016**, *39*, 4603–4609.
13. Bagheri, H.; Es'haghi, A.; Es'haghi, A.; Basiripour, F. Microwave-assisted extraction and high-throughput monolithic-polymer-based micro-solid-phase extraction of organophosphorus triazole, and organochlorine residues in apple. *J. Sep. Sci.* **2016**, *39*, 576–583.
14. Farajzadeh, M.A.; Sheykhzadeh, S.; Khorram, P. Extraction and preconcentration of some triazole pesticides in grape juice by salting out homogeneous liquid-liquid extraction in a narrow-bore tube prior to their determination by gas chromatography-flame ionization detection. *Food Anal. Methods* **2014**, *7*, 1229–1237.
15. Farajzadeh, M.A.; Feriduni, B.; Mogaddam, M.R.A. Extraction and enrichment of triazole and triazine pesticides from honey using air-assisted liquid-liquid microextraction. *J. Food Sci.* **2014**, *79*, H2140–H2148.
16. Aladaghlo, Z.; Fakhari, A.R.; Alavioon, S.I.; Dabiri, M. Ultrasound assisted dispersive solid phase extraction of triazole fungicides by using an N-heterocyclic carbene copper complex supported on ionic liquid-modified graphene oxide as a sorbent. *Microchim. Acta* **2019**, *6*, 209.
17. Zeng, H.; Xie, X.; Huang, Y.; Chen, J.; Liu, Y.; Zhang, Y.; Mai, X.; Deng, J.; Fan, H.; Zhang, W. Enantioseparation and determination of triazole fungicides in vegetables and fruits by aqueous two-phase extraction coupled with online heart-cutting two-dimensional liquid chromatography. *Food Chem.* **2019**, *301*, 125265.
18. Kachangoon, R.; Vichapong, J.; Santaladchaiyakit, Y.; Srijaranai, S. Green fabrication of Moringa oleifera seed as efficient bio-sorbent for selective enrichment of triazole fungicides in environmental water, honey and fruit juice samples. *Microchem. J.* **2022**, *175*, 107194.
19. Vichapong, J.; Burakham, R.; Santaladchaiyakit, Y.; Srijaranai, S. A preconcentration method for analysis of neonicotinoids in honey samples by ionic liquid-based cold-induced aggregation microextraction. *Talanta* **2016**, *155*, 216–221.
20. Gharehbaghi, M.; Shemirani, F.; Farahani, M.D. Cold-induced aggregation microextraction based on ionic liquids and fiber optic-linear array detection spectrophotometry of cobalt in water samples. *J. Hazard. Mater.* **2009**, *165*, 1049–1055.
21. Vichapong, J.; Burakham, R.; Srijaranai, S.; Grudpan, K. Room temperature imidazolium ionic liquid: A solvent for extraction of carbamates prior to liquid chromatographic analysis. *Talanta* **2011**, *84*, 1253–1258.
22. Liu, J.; Chi, Y.; Jiang, G.; Tai, C.; Peng, J.; Hu, J.-T. Ionic liquid-based liquid-phase microextraction, a new sample enrichment procedure for liquid chromatography. *J. Chromatogr. A* **2004**, *1026*, 143–147.
23. Hu, L.; Zhang, P.; Shan, W.; Wang, X.; Li, S.; Zhou, W.; Gao, H. In situ metathesis reaction combined with liquid-phase microextraction based on the solidification of sedimentary ionic liquids for the determination of pyrethroid insecticides in water samples. *Talanta* **2015**, *144*, 98–104.
24. Yao, C.; Anderson, J.L. Dispersive liquid-liquid microextraction using an in situ metathesis reaction to form an ionic liquid extraction phase for the preconcentration of aromatic compounds from water. *Anal. Bioanal. Chem.* **2009**, *395*, 1491–1502.
25. Baghdadi, M.; Shemirani, F. In situ solvent formation microextraction based on ionic liquids: A novel sample preparation technique for determination of inorganic species in saline solutions. *Anal. Chim. Acta* **2009**, *634*, 186–191.

26. Ullah, Z.; Azmi Bustam, M.; Man, Z.; Khan, A.S. Phosphonium-based ionic liquids and their application in separation of dye from aqueous solution. *ARPN J. Eng. Appl. Sci.* **2016**, *11*, 1653–1659.
27. Guo, W.; Hou, Y.; Wu, W.; Ren, S.; Tian, S.; Marsh, K.N. Separation of phenol from model oils with quaternary ammonium salts via forming deep eutectic solvents. *Green Chem.* **2013**, *15*, 226–229.
28. Khezeli, T.; Daneshfar, A.; Sahraei, R. A green ultrasonic-assisted liquid-liquid microextraction based on deep eutectic solvent for the HPLC-UV determination of ferulic, caffeic and cinnamic acid from olive, almond, sesame and cinnamon oil. *Talanta* **2016**, *150*, 577–585.
29. Shuang, Y.; Zhang, T.; Zhong, H.; Li, L. Simultaneous enantiomeric determination of multiple triazole fungicides in fruits and vegetables by chiral liquid chromatography/tandem mass spectrometry on a bridged bis(β -cyclodextrin)-bonded chiral stationary phase. *Food Chem.* **2021**, *345*, 128842.
30. Kadivar, M.; Aliakbar, A. A new composite based on graphene oxide-poly 3-aminophenol for solid-phase microextraction of four triazole fungicides in water and fruit juices prior to high-performance liquid chromatography analysis. *Food Chem.* **2019**, *299*, 125127.
31. Hu, X.Z.; Wang, J.X.; Feng, Y.Q. Determination of benzimidazole residues in edible animal food by polymer monolith microextraction combined with liquid chromatography-mass spectrometry. *J. Agric. Food Chem.* **2010**, *58*, 112–119.
32. Vichapong, J.; Santaladchaiyakit, Y.; Burakham, R.; Srijaranai, S. Determination of Benzimidazole Anthelmintics in Eggs by Advanced Microextraction with High-Performance Liquid Chromatography. *Anal. Lett.* **2015**, *48*, 617–631.
33. Domínguez-álvarez, J.; Mateos-Vivas, M.; García-Gómez, D.; Rodríguez-Gonzalo, E.; Carabias-Martínez, R. Capillary electrophoresis coupled to mass spectrometry for the determination of anthelmintic benzimidazoles in eggs using a QuEChERS with preconcentration as sample treatment. *J. Chromatogr. A* **2013**, *1278*, 166–174.