

Activation of Thioglycosides with Copper(II) Bromide

Faranak Pooladian,^a Samira Escopy,^{a,b} and Alexei V. Demchenko^{a,b*}

^[a] Department of Chemistry, Saint Louis University, 3501 Laclede Ave, St. Louis, Missouri 63103, USA;

^[b] Department of Chemistry and Biochemistry, University of Missouri – St. Louis, One University Boulevard, St. Louis, Missouri 63121, USA

Contents:

S1. NMR Monitoring to understand mechanistic aspects of the approach	S2S2. ¹ H
NMR spectra of known compounds	S4




Figure S1: NMR Monitoring to Understand Mechanistic Aspects of the Reaction.




Figure S2: ^1H NMR Spectra of Known Compounds.
S3




Figure S3: ¹H NMR (CDCl₃ 400 MHz) of compound 3.




Figure S4: ¹H NMR (CDCl₃ 400 MHz) of compound 7.




Figure S5: ¹H NMR (CDCl₃ 400 MHz) of compound 9.






Figure S6: ¹H NMR (CDCl₃ 400 MHz) of compound 13.






Figure S9: ^1H NMR (CDCl_3 400 MHz) of compound 20.





Figure S10: ^1H NMR (CDCl_3 400 MHz) of compound 22.




[illegible]

Figure S16: ^1H NMR (CDCl_3 400 MHz) of compound **34**.