

Article

SNH Amidation of 5-Nitroisoquinoline: Access to Nitro- and Nitroso Derivatives of Amides and Ureas on the Basis of Isoquinoline

Elena K. Avakyan, Anastasia A. Borovleva, Diana Yu. Pobedinskaya, Oleg P. Demidov, Artem P. Ermolenko, Alexander N. Larin and Ivan V. Borovlev *

Department of Chemistry, North Caucasus Federal University, 1a Pushkin St., Stavropol 355017, Russia

* Correspondence: ivborovlev@rambler.ru

Abstract: For the first time, amides and ureas based on both 5-nitroisoquinoline and 5-nitrosoisoquinoline were obtained by direct nucleophilic substitution of hydrogen in the 5-nitroisoquinoline molecule. In the case of urea and monosubstituted ureas, only 5-nitrosoisoquinoline-6-amine is formed under anhydrous conditions.

Keywords: heterocycles; S_NH methodology; metal-free C–N bond formation; 5-nitroisoquinoline; redox process; oxidative S_NH amidation; S_NH dialkylcarbamoylation; regioselectivity

Citation: Avakyan, E.K.; Borovleva, A.A.; Pobedinskaya, D.Yu.; Demidov, O.P.; Ermolenko, A.P.; Larin, A.N.; Borovlev, I.V. SNH Amidation of 5-Nitroisoquinoline: Access to Nitro- and Nitroso Derivatives of Amides and Ureas on the Basis of Isoquinoline. *Molecules* **2022**, *27*, 7862. <https://doi.org/10.3390/molecules27227862>

Academic Editor:
Alexander F. Khlebnikov

Received: 26 October 2022
Accepted: 10 November 2022
Published: 14 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The molecular framework of isoquinoline is the basis of an extensive family of alkaloids exhibiting diverse biological activity [1–3], and their synthetic derivatives are widely used in pharmacology and medicine [4]. Thus, further functionalization of isoquinoline seems to be a very promising direction. Without detracting from the advantages of multistage functionalization methods, where the final stage is cyclization with the formation of an isoquinoline ring [5–13], we note that modern requirements for the synthesis of derivatives of aromatic and heteroaromatic compounds imply their direct C–H functionalization, the economy of all transformation parameters, which is in line with the so-called PASE (Pot-, Atom-, and Step-Economic) concept [14], as well as the principles of green chemistry [15].

In the case of π -deficient azines and nitroarenes, oxidative nucleophilic hydrogen substitution (S_NH) reactions satisfy these requirements, which do not require the preliminary introduction of a good leaving group into the substrate or reagent molecule, as well as the use of expensive catalysts and ligands [16–19]. They include an additional step with the formation of σ^H adduct and its subsequent aromatization due to an external oxidizing agent. Organic and inorganic compounds and atmospheric oxygen are used to oxidize σ^H adducts [20,21], while electrochemical oxidation is used for stable intermediates [22–24]. Even in the absence of an external oxidant, the NO_2 [25] group or the C=N bond of the substrate [26] can also act as hydride anion acceptors. The most probable mechanism for the dehydroaromatization of the σ^H complex is the successive transfer of an electron, a proton, and one more electron (EPE mechanism) to the oxidant molecule [27]. Later, in the arylamination of nitroarenes, in addition to the oxidative one, another way of aromatization of the σ^H adduct was discovered by its dehydration with the formation of the corresponding nitroso compounds [28–34]. On the whole, the S_NH methodology has already found application in industry [35,36] and, in some cases, is a good alternative to cross-coupling reactions with the participation of transition metals [37].


The aim of the first stage of the work was to study the possibility of direct nucleophilic substitution of hydrogen by the N-amide function in the 5-nitroisoquinoline mol-

ecule. It is known that this compound readily enters into reactions of oxidative amination [38], methylation [39], whereas its quinoline analogue even into arylamination reaction [40]. However, unlike other N-nucleophiles, reactions of direct hydrogen substitution by the N-amide function in the series of azines and nitroarenes are still quite rare. The first report on the S_N^H amidation of nitrobenzene appeared only in 1993 [41] during the development of an industrial method for the preparation of *p*-nitroaniline [42].

In continuation of these studies, the oxidative S_N^H amidation of 1,3,7-triazapyrene [43], acridine [44], 3-nitropyridine [45] and 5(6,7,8)-nitroquinolines [46] was successfully performed in our laboratory. In all cases, the process was carried out in anhydrous DMSO by the action on the substrate with the previously obtained anion of the corresponding carboxamide at room temperature, using atmospheric oxygen [43,46] or $K_3Fe(CN)_6$ [44,45] as an oxidizing agent.

2. Results and Discussion

We implemented two approaches to the S_N^H amidation of 5-nitroisoquinoline (**1**), which differed mainly in the presence or absence of a small amount of water in the reaction mass, and which led to significantly different results (Scheme 1). The optimization of the first of them (method A) was carried out using the example of the reaction of substrate **1** with *p*-methylbenzamide in anhydrous DMSO at room temperature, preliminarily generating the amide anion by the action of NaH in the same solvent. The best result was shown by using 2 equiv. amide anion per 1 equiv. substrate (Table 1, entry 1). After adding 5-nitroisoquinoline (**1**), the process has completed within 1.5 h with the formation of a mixture of 4-methyl-*N*-(5-nitroisoquinolin-8-yl)benzamide (**2a**) and 4-methyl-*N*-(5-nitrosoisoquinoline-6-yl)benzamide (**3a**) with a total yield of 45%, the separation of which was carried out by chromatography on silica gel. All spectral data are available in the Supplementary Materials file submitted with this article.



Scheme 1. S_N^H Amidation of 5-nitroisoquinoline.

Table 1. Optimization of the Reaction Conditions of S_N^H Amidation of 5-Nitroisoquinoline (**1**) with the *p*-Methylbenzamide N-anion in anhydrous DMSO (Method A).


Entry	Reaction Time, h	Excess of NaH, Equiv	Excess of Amide, Equiv	Yield, % ^a	
				2a	3a
1 ^b	1.5	2	2	19	26
2 ^b	1.5	4	4	30	traces
3 ^b	1.5	6	6	24	7.5
4 ^b	1.5	1	3	45	traces
5 ^b	1.5	4	2	28	15
6 ^{b,c}	1.5	2	2	28	traces
7 ^d	0.5	2	2	40	traces
8 ^{b,e}	1.5	2	2	15	23

^a Isolated yields after column chromatography. ^b The experiment was performed at room temperature. ^c The experiment was performed in presence of K₃Fe(CN)₆. ^d The experiment was performed at 60 °C. ^e The reaction was performed under argon.

Increasing the excess of the amide anion (entries 2,3), changing the ratio of *p*-methylbenzamide and NaH (entries 4,5), using K₃Fe(CN)₆ as an external oxidizer (entry 6), and increasing the temperature (entry 7) turned out to be ineffective. The reaction was carried out without isolation from air oxygen; however, its performance in an argon atmosphere only slightly reduced the yield of nitro product **2a** (entry 8). These data suggest that, as in the case of 3-nitropyridine [45] and 5-nitroquinoline [46], 5-nitroisoquinoline exhibits a dual reactivity, being not only a substrate, but also the main oxidizer of σ^H adducts in the formation of nitro-amides **2**. Naturally, this reduces the yield of target compounds and leads to the appearance of by-products.

Anions of benzamide- and *p*-methoxybenzamide react similarly with the formation of the corresponding nitroamides **2b,c** and nitrosoamides **3b,c**. Note, however, that the reaction of the initial substrate with *p*-, *m*-, and *o*-nitrobenzamides gives only nitroamides **2d–f** (Scheme 1).

The mechanism of amidation of 5-nitroisoquinoline (**1**) includes the addition of a nucleophile in the *ortho*- and *para*-positions towards the NO₂-group, and the *para* σ^H adduct **5** further undergoes oxidative aromatization to form nitroamides **2a–f** (Scheme 2, route a), while its *ortho* analog **6** is aromatized by proton transfer and elimination of a water molecule, giving nitrosoamides **3a–c** (Scheme 2, route b).



Scheme 2. Proposed mechanism for the synthesis of nitro- **2a–f** and nitroso compounds **3a–c**.

The starting point for the development of another approach to the S_N^H amidation of 5-nitroisoquinoline (**1**) was the fact that the authors of the first work on amidation by direct hydrogen substitution [41] carried out the process in a non-absolute medium, using dihydrate of tetramethylammonium hydroxide as the base, which led to the formation of only products substitution at the *p*-position of the nitrobenzene molecule. In this approach, we used commercial DMSO containing ~0.5% water as well as KOH instead of NaH as the base (method B). As it turned out, when using a 4-molar excess of the corresponding aromatic amides, only the corresponding

N-(5-nitroisoquinolin-8-yl)benzamides **2a–f** were formed in 34–53% yields (Scheme 1). Apparently, in the presence of water, the hydrated amide anions, which have a large volume, experience steric difficulties in entering the *o*-position with respect to the NO₂ group, i.e., in position 6 of the 5-nitroisoquinoline molecule. Despite relatively low yields, nitro- **2** and nitroso compounds **3** are of considerable interest for further functionalization of isoquinoline, and it is very problematic to obtain nitrosoamides **3a–c** by other methods.

Increasing the water content in the mixture with DMSO to 5% in method B, or using amides of aliphatic acids (acetic, propionic and isobutyric) under the conditions of both methods, led to a complex mixture of substances.


A feature of the ¹H NMR spectra of nitrosoamides **3a–c** in CDCl₃ is a strong down-field shift of NH proton signals (δ ~13.5–13.6 ppm), which indicates a strong intramolecular hydrogen bond NH...O=N. The structures of the compounds **2a** (CCDC 2159575) and **3a** (CCDC 2159573) (Figure 1) were confirmed by X-ray determination [47].



Figure 1. (a) ORTEP diagram of nitro compound **2a**; (b) ORTEP diagram of nitroso compound **3a** (The dashed line shows the intramolecular hydrogen bond). The crystallographic data could be found in the Supplementary Materials.


The aim of the second stage of the work was to study the possibility of S_N^H amidation of 5-nitroisoquinoline with urea and its derivatives. We have previously shown that unsubstituted urea can act as an aminating agent in nucleophilic substitution reactions [47–50]. For example, the reaction of acridine with the urea anion in anhydrous DMSO led to 9-aminoacridine in 78% yield [50]. However, alkylureas under the same conditions entered into the S_N^H alkyl(dialkyl)carbamoylamine reaction, allowing to introduce the residues of the corresponding ureas into acridine [50] and 3-nitropyridine [51] molecules. Under anhydrous conditions, urea anions easily form stable 6-adducts at position 9 with 10-alkylacridinium cations [52].

We have found that urea and its monosubstituted derivatives such as phenyl-, tert-butyl- and (1,1-dimethylpentyl)urea react with 5-nitroisoquinoline under anhydrous conditions (method A) to form the same compound—5-nitrosoisoquinoline-6-amine (**7**; Scheme 3). In the ¹H NMR spectrum of compound **7**, even in such a polar solvent as DMSO-d₆, the NH₂ group gives two broadened singlets at δ 11.53 and 8.90 ppm, the first of which corresponds to a proton bound by a strong intramolecular hydrogen bond NH...O=N. In the ¹³C NMR spectrum, the signals at δ 136.6 and 139.2 ppm appear only with increasing accumulation time and are strongly broadened. In our opinion, this is the result of the well-known prototropic tautomerism of the nitrosamine-azaquinone oxime type [53–55] (Scheme 3), and the tautomerization rate for this compound is relatively slow in the NMR time scale, and these signals refer to the C₅ and C₆ atoms of the isoquinoline cycle. All spectral data are available in the Supplementary Materials file submitted with this article.



Scheme 3. Synthesis and tautomerism of compound 7.


Unlike amides, S_N^H reactions of 5-nitroisoquinoline with ureas under anhydrous conditions (method A) proceed exclusively at position 6 in accordance with the mechanism of formation of nitrosoamides **3a–c** (Scheme 2, route **b**). However, the nucleophilic substitution product **8** is unstable, and under the reaction conditions, the urea radical is converted into an amino group according to the route we proposed earlier [43,48] (Scheme 4). The key to it is the elimination of the isocyanic acid molecule or its ester (RNCO) and the formation of the anion **9**.



Scheme 4. Proposed pathway for the formation of the compound 7.

Application of the conditions of method B to the reaction of 5-nitroisoquinoline with anions of urea and its monosubstituted derivatives was unsuccessful, since a complex mixture of substances is formed in the presence of water.

However, 1,1-dialkylurea anions react with 5-nitroisoquinoline (**1**) under the conditions of both methods, but form different substitution products, albeit in relatively low yields (Scheme 5). So, under anhydrous conditions (method A), the 1,1-dimethylurea anion leads to the S_N^H product at position 6—1,1-dimethyl-3-(5-nitrosoisoquinolin-6-yl)urea (**10a**) in 41% yield. Amides of pyrrolidine-1-carboxylic, piperidine-1-carboxylic, and morpholine-4-carboxylic acids also react similarly, forming ureas based on 5-nitrosoisoquinoline **10b–d**. Undoubtedly, the mechanism of formation of compounds **10a–d** corresponds to the general mechanism for obtaining nitroso compounds (Scheme 2, route **b**).



Scheme 5. S_NH Reactions of 5-nitrosoquinoline (**1**) with an ureas.

In the 1H NMR spectra of these compounds, the NH proton signal is strongly shifted downfield in both $CDCl_3$ and $DMSO-d_6$ (δ ~13.5–13.6 ppm), which also indicates a strong intramolecular hydrogen bond. In addition, if under normal conditions of recording the spectrum (25 °C) in $CDCl_3$, the protons of two methyl groups of compound **10a** give one broadened singlet, then when cooled to only 13.7 °C, it splits into two signals, which indicates the nonequivalence of methyl groups under these temperature conditions. In our opinion, this is due to the well-known difficulty in rotation relative to the C(O)-N amide bond. At 25 °C the rotation accelerates and the signals from these groups coalesce. However, in the case of N-(5-nitrosoquinolin-6-yl)pyrrolidine-1-carboxamide (**10b**), the protons of both the α - and β -methylene groups of the pyrrolidine ring are not equivalent even at 25 °C and give separate signals (as do the signals of carbon atoms of these groups in ^{13}C NMR). The difficulty of rotation relative to the amide bond of compounds **10c** and **10d** is noticeable only in the broadening of the signals of the corresponding protons, which appear equivalent. Apparently, this is due to the lower conformational rigidity of the piperidine and morpholine rings of compounds **10c** and **10d** compared to the pyrrolidine one.

In the presence of water (method B), the reactions of 5-nitrosoquinoline with 1,1-dialkylurea anions proceed by the mechanism of oxidative nucleophilic substitution (Scheme 2, route a) to position 8 with the formation of exclusively S_NH dialkylcarbamoylamination products **11a–d** (Scheme 5). The structure of the compound **10a** (CCDC 2159579, Figure 2) was confirmed by X-ray determination [47].




Figure 2. ORTEP diagram of nitroso compound **10a**. The crystallographic data could be found in the Supplementary Materials.

3. Materials and Methods

^1H and ^{13}C NMR spectra were recorded on a Bruker Avance HD 400 spectrometer in the solvent indicated relative to residual DMSO signals [56] or TMS as internal standard when CDCl_3 was used as a solvent. and—85% H_3PO_4 . The NMR-spectra of the newly synthesized compounds could be found in the Supplementary Materials. HRMS were registered on a Bruker UHR-TOF Maxis™ Impact instrument using the ESI technique. All melting points were determined in glass capillaries using REACH Devices RD-MP and Electrothermal IA 9200 instruments and are uncorrected. The reaction progress and the purity of the obtained compounds were controlled by TLC on Silufol UV-254 plates. All experiments were carried out in a reactor protected from atmospheric moisture, but without isolation from atmospheric oxygen. Sodium hydride (60% paraffin oil suspension, Merck, Darmstadt, Germany) and 5-nitrosoquinoline (abcr GmbH & Co. KG, Heilbronn, Germany) were used without further purification.

N-(5-nitrosoquinolin-8-yl)benzamides **2a–f** and *N*-(5-nitrosoisoquinoline-6-yl)benzamides **3a–c**; (General Procedures):

Method A: To a solution of 1 mmol of the corresponding benzamide in 4 mL of anhydrous DMSO 40 mg of a suspension of sodium hydride in paraffin oil (1 mmol of NaH) and after 10 min 87 mg (0.5 mmol) 5-nitrosoquinoline were added sequentially at room temperature. The mixture was intensively stirred at room temperature for 1.5 h. Then, the reaction mass was poured into 50 g of ice and, upon reaching room temperature, it was acidified with dilute HCl to pH~7. The precipitate that formed was filtered off, washed with water, and dried. The dry product was separated into the appropriate fractions by dry flash chromatography [57] on silica gel. In the synthesis of compounds **2a–c** and **3a–c**, the mixture was eluted with toluene—ethyl acetate (15:1) and the second and third fractions were collected (the first one, slightly colored fraction was discarded; it contains the starting benzamides and non-polar impurities). Nitrosoamides **3a–c** were obtained from the second green fraction, and nitroamides **2a–c** were obtained from the third yellow fraction. In the synthesis of compounds **2d–f**, the mixture was eluted with toluene—ethyl acetate (5:1) and the second yellow fraction was collected.

Method B: To a solution of 2 mmol of the corresponding benzamide in 8 mL of DMSO containing 0.5% water, 112 mg of KOH (2 mmol), 87 mg (0.5 mmol) of 5-nitrosoquinoline, and 658 mg (2 mmol) of $\text{K}_3\text{Fe}(\text{CN})_6$ were added sequentially at room temperature. The mixture was vigorously stirred at room temperature for 2.5 h, then the reaction mixture was poured into 50 g of ice and, upon reaching room temperature, acidified with dilute HCl to pH~7. The precipitate was filtered off, washed with water, and dried. The dry product was purified by dry flash chromatography on silica gel eluting with benzene-ethyl acetate (5:1) and collecting a second yellow fraction.

4-Methyl-*N*-(5-nitrosoquinolin-8-yl)benzamide (**2a**). Yellow solid; yield: 29.2 mg (19%, Method A); 81.4 mg (53%, Method B); mp 205–206 °C (EtOAc). ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ = 11.01 (br s, 1H, NH), 9.68 (d, J = 0.5 Hz, 1H, H-1), 8.79 (d, J = 6.2 Hz, 1H, H-3), 8.74 (d, J = 8.6 Hz, 1H, H-6), 8.45 (d, J = 6.2 Hz, 1H, H-4), 8.15 (d, J = 8.6 Hz, 1H, H-7), 8.04 (d, J = 8.1 Hz, 2H, H-2,6 Ar), 7.42 (d, J = 8.1 Hz, 2H, H-3,5 Ar), 2.43 (s, 3H, CH_3). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ = 166.6, 149.4, 146.3, 142.6, 141.9, 140.4, 130.9, 129.9, 129.1, 128.5, 128.3, 122.2, 121.2, 115.1, 21.1. HRMS (ESI): m/z $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{17}\text{H}_{13}\text{N}_3\text{NaO}_3$: 330.0849; found: 330.0831.

4-Methyl-*N*-(5-nitrosoisoquinolin-6-yl)benzamide (**3a**). Green solid; yield: 37.8 mg (26%, Method A); mp 168–169 °C (dec., EtOAc). ^1H NMR (400 MHz, CDCl_3): δ = 13.57 (br s, 1H, NH·O), 9.76 (d, J = 6.6 Hz, 1H, H-4), 9.73 (s, 1H, H-1), 9.56 (d, J = 9.5 Hz, 1H, H-8), 8.84 (d, J = 6.6 Hz, 1H, H-3), 8.64 (d, J = 9.5 Hz, 1H, H-7), 8.11 (d, J = 8.1 Hz, 2H, H-2,6 Ar), 7.50 (d, J = 8.56 Hz, 2H, H-3,5 Ar), 2.53 (s, 3H, CH_3). ^{13}C NMR (100 MHz, CDCl_3): δ = 168.2, 146.1, 145.7, 145.4, 142.3, 141.1, 136.4, 130.4, 129.5, 128.6, 127.9, 124.3, 123.1, 119.3, 22.0. HRMS (ESI): m/z $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{14}\text{N}_3\text{O}_2$: 292.1081; found: 292.1072.

N-(5-Nitrosoquinolin-8-yl)benzamide (**2b**). Yellow solid; yield: 32.2 mg (22%, Method A); 64.5 mg (44%, Method B); mp 221–222 °C (EtOAc). ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ =

11.09 (br s, 1H, NH), 9.70 (s, 1H, H-1), 8.80 (d, $J = 6.1$ Hz, 1H, H-3), 8.75 (d, $J = 8.6$ Hz, 1H, H-6), 8.45 (d, $J = 6.1$ Hz, 1H, H-4), 8.17 (d, $J = 8.6$ Hz, 1H, H-7), 8.13 (d, $J = 8.1$ Hz, 2H, H-2,6 Ph), 7.69 (t, $J = 7.7$ Hz, 1H, H-4 Ph), 7.64–7.59 (m, 2H, H-3,5 Ph). ^{13}C NMR (100 MHz, DMSO- d_6): $\delta = 166.8, 149.4, 146.3, 141.8, 140.5, 133.8, 132.4, 129.8, 128.6, 128.5, 128.3, 122.2, 121.3, 115.1$. HRMS (ESI): m/z $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{16}\text{H}_{11}\text{N}_3\text{NaO}_3$: 316.0693; found: 316.0692.

N-(5-Nitrosoisoquinolin-6-yl)benzamide (**3b**). Green solid; yield: 31.9 mg (23%, Method A); mp 173–174 °C (dec., EtOAc). ^1H NMR (400 MHz, CDCl_3): $\delta = 13.53$ (br s, 1H, NH \cdots O), 9.41 (br s, 1H, H-1), 9.36–9.33 (m, 2H, H-4,8), 8.88 (d, $J = 6.1$ Hz, 1H, H-3), 8.43 (d, $J = 9.4$ Hz, 1H, H-7), 8.20 (d, $J = 7.0$ Hz, 2H, H-2,6 Ph), 7.74–7.65 (m, 3H, H-3,4,5 Ph). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 168.5, 150.6, 147.6, 145.7, 141.5$ (2C), 133.9 (2C), 133.0, 129.6, 128.4, 124.3, 121.0, 116.4. HRMS (ESI): m/z $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{12}\text{N}_3\text{O}_2$: 278.0924; found: 278.0919.

4-Methoxy-*N*-(5-nitrosoisoquinolin-8-yl)benzamide (**2c**). Yellow solid; yield: 35.5 mg (22%, Method A); 66.2 mg (41%, Method B); mp 214–215 °C (EtOAc). ^1H NMR (400 MHz, DMSO- d_6): $\delta = 10.93$ (br s, 1H, NH), 9.67 (s, 1H, H-1), 8.79 (d, $J = 6.3$ Hz, 1H, H-3), 8.73 (d, $J = 8.6$ Hz, 1H, H-6), 8.45 (d, $J = 6.3$ Hz, 1H, H-4), 8.14 (d, $J = 8.6$ Hz, 1H, H-7), 8.12 (d, $J = 8.9$ Hz, 2H, H-2,6 Ar), 7.14 (d, $J = 8.9$ Hz, 2H, H-3,5 Ar), 3.88 (s, 3H, CH_3O). ^{13}C NMR (100 MHz, DMSO- d_6): $\delta = 166.0, 162.6, 149.5, 146.3, 142.1, 140.3, 130.3, 129.9, 128.5, 125.8, 122.1, 121.1, 115.1, 113.8, 55.6$. HRMS (ESI): m/z $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{17}\text{H}_{13}\text{N}_3\text{NaO}_4$: 346.0798; found: 346.0798.

4-Methoxy-*N*-(5-nitrosoisoquinolin-6-yl)benzamide (**3c**). Green solid; yield: 35.3 mg (23%, Method A); mp 186–187 °C (dec., EtOAc). ^1H NMR (400 MHz, CDCl_3): $\delta = 13.64$ (br s, 1H, NH \cdots O), 9.31 (s, 1H, H-1), 9.29–9.24 (m, 2H, H-4,8), 8.88 (d, $J = 6.0$ Hz, 1H, H-3), 8.34 (d, $J = 9.4$ Hz, 1H, H-7), 8.18 (d, $J = 8.8$ Hz, 2H, H-2,6 Ar), 7.14 (d, $J = 8.8$ Hz, 2H, H-3,5 Ar), 3.96 (s, 3H, CH_3). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 168.0, 164.1, 151.9, 148.1, 148.0, 141.5, 130.6$ (2C), 125.3, 124.4, 120.2, 115.5, 114.7 (2C), 55.8. HRMS (ESI): m/z $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{14}\text{N}_3\text{O}_3$: 308.1030; found: 308.1030.

4-Nitro-*N*-(5-nitrosoisoquinolin-8-yl)benzamide (**2d**). Yellow solid; yield: 64.2 mg (38%, Method A); 87.9 mg (52%, Method B); mp 244–245 °C (EtOAc). ^1H NMR (400 MHz, DMSO- d_6): $\delta = 11.37$ (br s, 1H, NH), 9.75 (s, 1H, H-1), 8.81 (d, $J = 6.1$ Hz, 1H, H-3), 8.76 (d, $J = 8.6$ Hz, 1H, H-6), 8.46–8.41 (m, 3H, H-4, H-2,6 Ar), 8.36 (d, $J = 8.9$ Hz, 2H, H-3,5 Ar), 8.18 (d, $J = 8.6$ Hz, 1H, H-7). ^{13}C NMR (100 MHz, DMSO- d_6): $\delta = 165.4, 149.6, 149.5, 146.4, 141.3, 140.8, 139.6, 129.9, 129.8, 128.5, 123.6, 122.3, 121.6, 115.2$. HRMS (ESI): m/z $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{16}\text{H}_{10}\text{N}_4\text{NaO}_5$: 361.0543; found: 361.0552.

3-Nitro-*N*-(5-nitrosoisoquinolin-8-yl)benzamide (**2e**). Yellow solid; yield: 89.6 mg (53%, Method A); 76.1 mg (45%, Method B); mp 215–216 °C (EtOAc). ^1H NMR (400 MHz, DMSO- d_6): $\delta = 11.38$ (br s, 1H, NH), 9.74 (s, 1H, H-1), 8.95 (t, $J = 1.8$ Hz, 1H, H-2 Ar), 8.81 (d, $J = 6.2$ Hz, 1H, H-3), 8.76 (d, $J = 8.6$ Hz, 1H, H-6), 8.57–8.50 (m, 2H, H-4,6 Ar), 8.45 (d, $J = 6.2$ Hz, 1H, H-4), 8.15 (d, $J = 8.6$ Hz, 1H, H-7), 7.91 (t, $J = 8.0$ Hz, 1H, H-5 Ar). ^{13}C NMR (100 MHz, DMSO- d_6): $\delta = 164.8, 149.6, 147.8, 146.3, 141.2, 140.9, 135.3, 134.8, 130.3, 129.8, 128.5, 126.8, 123.1, 122.3, 121.8, 115.1$. HRMS (ESI): m/z $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{16}\text{H}_{10}\text{N}_4\text{NaO}_5$: 361.0543; found: 361.0546.

2-Nitro-*N*-(5-nitrosoisoquinolin-8-yl)benzamide (**2f**). Yellow solid; yield: 64.2 mg (38%, Method A); 57.5 mg (34%, Method B); mp 242–243 °C (EtOAc). ^1H NMR (400 MHz, DMSO- d_6): $\delta = 11.48$ (br s, 1H, NH), 9.71 (s, 1H, H-1), 8.81 (d, $J = 6.2$ Hz, 1H, H-3), 8.79 (d, $J = 8.4$ Hz, 1H, H-6), 8.45 (d, $J = 6.2$ Hz, 1H, H-4), 8.27 (br d, $J = 8.2$ Hz, 2H, H-7, H-3 Ar), 8.03–7.95 (m, 2H, H-5,6 Ar), 7.87–7.81 (m, 1H, H-4 Ar). ^{13}C NMR (100 MHz, DMSO- d_6): $\delta = 165.8, 148.8, 146.5, 146.1, 140.8, 140.5, 134.6, 132.1, 131.4, 130.2, 129.6, 128.6, 124.4, 121.5, 120.1, 115.3$. HRMS (ESI): m/z $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{16}\text{H}_{10}\text{N}_4\text{NaO}_5$: 361.0543; found: 361.0524.

5-Nitrosoisoquinolin-6-amine (**7**)

To a solution of 1 mmol of the corresponding urea (Scheme 3) in 4 mL of anhydrous DMSO 40 mg of a suspension of sodium hydride in paraffin oil (1 mmol of NaH) and

after 10 min 87 mg (0.5 mmol) 5-nitrosoquinoline were added sequentially at room temperature. The mixture was vigorously stirred at room temperature for 1 h. Then, the reaction mass was poured into 50 g of ice and, upon reaching room temperature, acidified with dilute HCl to pH=7. The precipitate formed was filtered off, washed with water, and dried. The dry product was purified by crystallization from ethyl acetate.

Green solid; yield: 32.9 mg (38%, from urea); 50.2 mg (58%, from phenylurea); 65.7 mg (76%, from *tert*-butylurea); 51.9 mg (60%, from 1,1-dimethylpentylurea); mp 256–257 °C (dec., EtOAc). ¹H NMR (400 MHz, DMSO-*d*₆): δ = 11.53 (br s, 1H, NH⋯O), 9.03 (s, 1H, H-1), 8.90 (br s, 1H, NH), 8.82 (d, *J* = 5.7 Hz, 1H, H-4), 8.62 (d, *J* = 5.7 Hz, 1H, H-3), 8.02 (d, *J* = 9.3 Hz, 1H, H-8), 7.13 (d, *J* = 9.3 Hz, 1H, H-7). ¹³C NMR (100 MHz, DMSO-*d*₆): δ = 151.2, 147.8, 147.7, 139.2, 137.9, 136.6, 122.1, 121.3, 112.9. HRMS (ESI): *m/z* [M + H]⁺ calcd for C₉H₈N₃O: 174.0662; found: 174.0662.

1,1-Dialkyl-3-(5-nitrosoisoquinolin-6-yl)ureas (10a–d); (General Procedure):

To a solution of 1 mmol of the corresponding urea (Scheme 5, Method A) in 4 mL of anhydrous DMSO 40 mg of a suspension of sodium hydride in paraffin oil (1 mmol of NaH) and after 10 min 87 mg (0.5 mmol) 5-nitrosoquinoline were added sequentially at room temperature. The mixture was vigorously stirred at room temperature for 2 h. Then, the reaction mass was poured into 50 g of ice and, upon reaching room temperature, acidified with dilute HCl to pH=7. The precipitate formed was filtered off, washed with water, and dried. The dry product was isolated by dry flash chromatography³² on silica gel, collecting in all cases a second yellow or light green fraction, from which compounds **10a–d** were obtained. Eluents: PhMe- EtOAc (10:1) for **10a** and **10d**; PhMe- EtOAc (3:2) for **10b**; PhMe- EtOAc (1:1) for **10c**.

1,1-Dimethyl-3-(5-nitrosoisoquinolin-6-yl)urea (10a). Dark green solid; yield: 50.0 mg (41%); mp 161–162 °C (dec., PhMe-EtOAc). ¹H NMR (400 MHz, DMSO-*d*₆): δ = 13.43 (br s, 1H, NH⋯O), 9.30 (s, 1H, H-1), 8.96 (d, *J* = 5.8 Hz, 1H, H-3), 8.79 (d, *J* = 5.8 Hz, 1H, H-4), 8.77 (d, *J* = 9.4 Hz, 1H, H-8), 8.46 (d, *J* = 9.4 Hz, 1H, H-7), 3.14 (br s, 6H, N(CH₃)₂). ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 13.69 (br s, 1H, NH⋯O), 9.23 (s, 1H, H-1), 9.17 (d, *J* = 5.8 Hz, 1H, H-3), 9.00 (d, *J* = 9.5 Hz, 1H, H-8), 8.81 (d, *J* = 5.8 Hz, 1H, H-4), 8.19 (d, *J* = 9.5 Hz, 1H, H-7), 3.26 (br s, 6H, N(CH₃)₂). ¹H NMR (400 MHz, CDCl₃, 13.7 °C): δ = 13.58 (br s, 1H, NH⋯O), 9.56 (d, *J* = 6.3 Hz, 1H, H-3), 9.46 (s, 1H, H-1), 9.24 (d, *J* = 9.6 Hz, 1H, H-8), 8.79 (d, *J* = 6.3 Hz, 1H, H-4), 8.36 (d, *J* = 9.6 Hz, 1H, H-7), 3.45 (br s, 3H, N(CH₃)₂^b), 3.16 (br s, 3H, N(CH₃)₂^a). ¹³C NMR (100 MHz, CDCl₃, 25 °C): δ = 154.8, 151.0, 148.1, 146.5, 141.0, 139.4, 129.0, 123.4, 120.3, 115.4, 37.0. HRMS (ESI): *m/z* [M + H]⁺ calcd for C₁₂H₁₃N₄O₂: 245.1033; found: 245.1030.

N-(5-Nitrosoisoquinolin-6-yl)pyrrolidine-1-carboxamide (10b). Green solid; yield: 70.2 mg (52%); mp 179–180 °C (dec., PhMe-EtOAc). ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 13.52 (br s, 1H, NH⋯O), 9.28 (br s, 1H, H-1), 9.26 (d, *J* = 6.2 Hz, 1H, H-3), 9.15 (br d, *J* = 9.5 Hz, 1H, H-8), 8.80 (d, *J* = 6.2 Hz, 1H, H-4), 8.22 (d, *J* = 9.5 Hz, 1H, H-7), 3.87 (t, *J* = 5.3 Hz, 2H, NCH₂^b), 3.59 (t, *J* = 5.3 Hz, 2H, NCH₂^a), 2.23–2.20 (m, 2H CH₂CH₂), 2.04–2.01 (m, 2H, CH₂CH₂). ¹³C NMR (100 MHz, CDCl₃): δ = 152.7, 149.8, 147.7, 144.6, 140.7 (2C), 123.1, 121.1, 115.9, 46.8, 46.4, 26.3, 24.9. HRMS (ESI): *m/z* [M + H]⁺ calcd for C₁₄H₁₅N₄O₂: 271.1190; found: 271.1186.

N-(5-Nitrosoisoquinolin-6-yl)piperidine-1-carboxamide (10c). Yellow solid; yield: 35.5 mg (25%); mp 198–199 °C (dec., PhMe-EtOAc). ¹H NMR (400 MHz, CDCl₃): δ = 13.81 (br s, 1H, NH⋯O), 9.26 (br s, 1H, H-1), 9.21 (d, *J* = 5.3 Hz, 1H, H-3), 8.93 (d, *J* = 9.5 Hz, 1H, H-8), 8.80 (br s, 1H, H-4), 8.19 (d, *J* = 9.5 Hz, 1H, H-7), 3.70 (br s, 4H, N(CH₂)₂), 1.75 (br s, 6H, (CH₂)₃). ¹³C NMR (100 MHz, CDCl₃): δ = 153.2, 150.3, 148.0, 145.4, 140.7 (2C), 123.2, 121.1 (2C), 115.7, 29.8, 25.9, 24.3. HRMS (ESI): *m/z* [M + H]⁺ calcd for C₁₅H₁₇N₄O₂: 285.1346; found: 285.1340. HRMS (ESI): *m/z* [M + Na]⁺ calcd for C₁₅H₁₆N₄NaO₂: 307.1163; found: 307.1160.

N-(5-Nitrosoisoquinolin-6-yl)morpholine-4-carboxamide (10d). Brown solid; yield: 28.6 mg (20%); mp 146–147 °C (dec., PhMe-EtOAc). ¹H NMR (400 MHz, CDCl₃): δ = 13.69 (br s, 1H, NH⋯O), 9.32 (s, 1H, H-1), 9.26 (d, 1H, *J* = 6.0 Hz, H-3), 8.97 (d, 1H, *J* = 9.5 Hz, H-8),

8.82 (d, 1H, $J = 6.0$ Hz, H-4), 8.26 (d, 1H, $J = 9.5$ Hz, H-7), 3.83–3.87 (m, 4H, $O(CH_2)_2$), 3.74–3.79 (m, 4H, $N(CH_2)_2$). ^{13}C NMR (100 MHz, $CDCl_3$): $\delta = 153.7, 150.3, 148.0, 145.3, 141.2$ (2C), 129.7, 123.4, 120.8, 115.9, 66.5, 44.6. HRMS (ESI): m/z $[M + H]^+$ calcd for $C_{14}H_{15}N_4O_3$: 287.1139; found: 287.1134.

1,1-Dialkyl-3-(5-nitroisoquinolin-8-yl)ureas (11a–d); (General Procedure):

To a solution of 2 mmol of the corresponding urea (Scheme 5, Method B) in 8 mL of DMSO, containing 0.5% water, 112 mg of KOH (2 mmol), 87 mg (0.5 mmol) of 5-nitroisoquinoline and 658 mg (2 mmol) $K_3Fe(CN)_6$ were added sequentially at room temperature. The mixture was intensively stirred at room temperature for 4 h, then the reaction mass was poured into 50 g of ice and, upon reaching room temperature, it was acidified with dilute HCl to pH~7. The aqueous solution was extracted with ethyl acetate (4×10 mL), the extract was dried with Na_2SO_4 . After evaporation of the solvent, the dry residue was purified by dry flash chromatography on silica gel,³² collecting in all cases the second yellow fraction, from which compounds **11a–d** were obtained. Eluents: PhMe-EtOAc (10:1) for **11a**; PhMe-EtOAc (3:2) for **11b**; PhMe-EtOAc (1:1) for **11c**; ethyl EtOAc for **11d**.

1,1-Dimethyl-3-(5-nitroisoquinolin-8-yl)urea (11a). Yellow solid; yield: 46.8 mg (36%); mp 218–219 °C (dec., PhMe-EtOAc). 1H NMR (400 MHz, $DMSO-d_6$): $\delta = 9.58$ (s, 1H, H-1), 9.25 (br s, 1H, NH), 8.73 (d, $J = 6.1$ Hz, 1H, H-3), 8.66 (d, $J = 8.8$ Hz, 1H, H-6), 8.46 (d, $J = 6.1$ Hz, 1H, H-4), 7.90 (d, $J = 8.8$ Hz, 1H, H-7), 3.07 (s, 6H, $N(CH_3)_2$). ^{13}C NMR (100 MHz, $DMSO-d_6$): $\delta = 155.4, 149.4, 146.2, 144.8, 138.0, 130.4, 128.8, 121.0, 117.9, 115.1, 36.6$. HRMS (ESI): m/z $[M + H]^+$ calcd for $C_{12}H_{13}N_4O_3$: 261.0982; found: 261.0979. HRMS (ESI): m/z $[M + Na]^+$ calcd for $C_{12}H_{12}N_4NaO_3$: 283.0785; found: 283.0792.

N-(5-Nitroisoquinolin-8-yl)pyrrolidine-1-carboxamide (11b). Yellow solid; yield: 32.9 mg (23%); mp 197–198 °C (dec., PhMe-EtOAc). 1H NMR (400 MHz, $DMSO-d_6$): $\delta = 9.61$ (br s, 1H, H-1), 9.10 (br s, 1H, NH), 8.74 (dd, $J = 6.2, 0.8$ Hz, 1H, H-3), 8.66 (dd, $J = 8.8, 0.7$ Hz, 1H, H-6), 8.47 (d, $J = 6.2$ Hz, 1H, H-4), 8.03 (br d, $J = 8.8$ Hz, 1H, H-7), 3.54 (br s, 4H, $N(CH_2)_2$), 1.92 (br s, 4H, $(CH_2)_2$). ^{13}C NMR (100 MHz, $DMSO-d_6$): $\delta = 153.3, 149.4, 146.3, 144.4, 137.9, 130.5, 128.9, 120.8, 117.7, 115.1, 46.1, 25.1$. HRMS (ESI): m/z $[M + H]^+$ calcd for $C_{14}H_{14}N_4O_3$: 287.1139; found: 287.1126.

N-(5-Nitroisoquinolin-8-yl)piperidine-1-carboxamide (11c). Yellow solid; yield: 52.5 mg (35%); mp 205–206 °C (dec., PhMe-EtOAc). 1H NMR (400 MHz, $DMSO-d_6$): $\delta = 9.54$ (br s, 1H, H-1), 9.44 (br s, 1H, NH), 8.73 (d, $J = 6.1$ Hz, 1H, H-3), 8.63 (d, $J = 8.8$ Hz, 1H, H-6), 8.47 (d, $J = 6.1$ Hz, 1H, H-4), 7.83 (d, $J = 8.8$ Hz, 1H, H-7), 3.55 (t, $J = 5.2$ Hz, 4H, $N(CH_2)_2$), 1.63–1.56 (m, 6H, $(CH_2)_3$). ^{13}C NMR (100 MHz, $DMSO-d_6$): $\delta = 154.6, 149.4, 146.3, 145.2, 137.9, 130.5, 128.9, 121.0, 117.8, 115.1, 45.2, 25.6, 24.0$. HRMS (ESI): m/z $[M + Na]^+$ calcd for $C_{15}H_{16}N_4NaO_3$: 323.1115; found: 323.1105.

N-(5-Nitroisoquinolin-8-yl)morpholine-4-carboxamide (11d). Yellow solid; yield: 58.9 mg (39%); mp 230–231 °C (dec., PhMe-EtOAc). 1H NMR (400 MHz, $DMSO-d_6$): $\delta = 9.59$ (s, 1H, H-1), 9.47 (br s, 1H, NH), 8.74 (d, 1H, $J = 6.1$ Hz, H-3), 8.66 (d, 1H, $J = 8.8$ Hz, H-6), 8.46 (d, 1H, $J = 6.1$ Hz, H-4), 7.90 (d, 1H, $J = 8.8$ Hz, H-7), 3.68 (t, 4H, $J = 4.5$ Hz, $O(CH_2)_2$), 3.56 (t, 4H, $J = 4.5$ Hz, $N(CH_2)_2$). ^{13}C NMR (100 MHz, $DMSO-d_6$): $\delta = 154.9, 149.4, 146.3, 144.6, 138.2, 130.4, 128.8, 121.1, 118.1, 115.1, 66.0, 44.6$. HRMS (ESI): m/z $[M + H]^+$ calcd for $C_{14}H_{15}N_4O_4$: 303.1088; found: 303.1089.

4. Conclusions

Thus, different regioselectivity was found in the reactions of S_NH substitution of 5-nitroisoquinoline with N-anions of aromatic amides and ureas, depending on the absence or presence of small amounts of water in the reaction mass. So, interaction with amides in anhydrous DMSO usually results in the formation of a mixture of hitherto unknown 8-arylamino-5-nitroisoquinolines and 6-arylamino-5-nitroisoquinolines in a small or moderate yield. In the presence of water, only nitro derivatives were formed. Anions of 1,1-dialkylureas in anhydrous DMSO give rise to 6-dialkylcarbamoylamino-5-nitroisoquinolines, whereas in the presence of water,

8-dialkylcarbamoylamino-5-nitrosoquinolines form. Urea itself and its monosubstituted derivatives under anhydrous conditions form exclusively 5-nitrosoisoquinoline-6-amine.

Supplementary Materials: Crystallographic data for the structures in this paper have been deposited in the Cambridge Crystallographic Data Center as a supplementary publication (**2a**, CCDC 2159575; **3a**, CCDC 2159573; **10a**, CCDC 2159579, www.ccdc.cam.ac.uk/getstructures). Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB12 1EZ, U.K [Fax: +44 1223 336033 or e-mail: deposit@ccdc.cam.ac.uk. The Supplemental Material file of CCDC includes the CIF file of **2a**, **3a**, **10a**. The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/molecules27227862/s1>, NMR spectroscopy data and X-ray analysis.

Author Contributions: Conceptualization, E.K.A. and I.V.B.; methodology, A.A.B.; validation, O.P.D., investigation, A.A.B., A.P.E., A.N.L. (synthetic chemistry), D.Y.P. (mass analysis) and O.P.D. (X-ray analysis); writing—original draft preparation, A.A.B. and I.V.B.; writing—review and editing, A.A.B. and I.V.B.; visualization, E.K.A.; supervision, O.P.D.; project administration, O.P.D.; funding acquisition O.P.D. All authors have read and agreed to the published version of the manuscript

Funding: The authors express their gratitude to the North Caucasian Federal University for financial support in the framework of the project to support the projects of scientific groups and individual scientists.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Supporting Information data include NMR spectral charts.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the all compounds are available from the authors.

References

1. Bala, M.; Kumar, S.; Pratap, K.; Verma, P.K.; Padwad, Y.; Singh, B. Bioactive isoquinoline alkaloids from *Cissampelos pareira*. *Nat. Prod. Res.* **2019**, *33*, 622–627. <http://doi.org/10.1080/14786419.2017.1402319>.
2. Huang, Q.-Q.; Bi, J.-L.; Sun, Q.-Y.; Yang, F.-M.; Wang, Y.-H.; Tang, G.-H.; Zhao, F.-W.; Wang, H.; Xu, J.-J.; Kennelly, E. J.; et al. Bioactive isoquinoline alkaloids from *Corydalis saxicola*. *Planta Med.* **2012**, *78*, 65–70. <http://doi.org/10.1055/s-0031-1280126>.
3. Giri, P.; Kumar, G.S. Isoquinoline alkaloids and their binding with polyadenylic acid: Potential basis of therapeutic action. *Mini Rev. Med. Chem.* **2010**, *10*, 568–577. <http://doi.org/10.2174/138955710791384009>.
4. Luo, C.; Wireko, M.A.; Wang, H.; Wu, C.; Wang, Q.; Zhang, H.; Cao, Y. Isoquinolines: Important Cores in Many Marketed and Clinical Drugs. *Anti-Cancer Agents Med. Chem.* **2021**, *21*, 811–824. <http://doi.org/10.2174/1871520620666200424132248>.
5. Jaryaram, V.; Sridhar, T.; Sharma, G.V.M.; Berrée, F.; Carboni, B. Synthesis of polysubstituted isoquinolines and related fused pyridines from alkenyl boronic esters via a copper-catalyzed azidation/aza-Wittig Condensation Sequence. *J. Org. Chem.* **2018**, *83*, 843–853. <http://doi.org/10.1021/acs.joc.7b02831>.
6. Zhou, W.; Zhang, Y.-X.; Nie, X.-D.; Si, C.-M.; Sun, X.; Wei, G.-G. Approach to Chiral 1-Substituted Isoquinolone and 3-Substituted Isoindolin-1-one by Addition–Cyclization Process. *J. Org. Chem.* **2018**, *83*, 9879–9889. <http://doi.org/10.1021/acs.joc.8b01282>.
7. Zhu, Z.; Tang, X.; Li, X.; Wu, W.; Deng, G.; Jiang, H. Palladium-Catalyzed C–H Functionalization of Aromatic Oximes: A Strategy for the Synthesis of Isoquinolines. *J. Org. Chem.* **2016**, *81*, 1401–1409. <http://doi.org/10.1021/acs.joc.5b02376>.
8. Zhou, S.; Wang, M.; Wang, L.; Chen, K.; Wang, J.; Song, C.; Zhu, J. Bidentate directing-enabled, traceless heterocycle synthesis: Cobalt-catalyzed access to isoquinolines. *Org. Lett.* **2016**, *18*, 5632–5635. <http://doi.org/10.1021/acs.orglett.6b02870>.
9. Chu, H.; Xue, P.; Yu, J.-T.; Cheng, J. Rhodium-Catalyzed Annulation of Primary Benzylamine with α -Diazo Ketone toward Isoquinoline. *J. Org. Chem.* **2016**, *81*, 8009–8013. <http://doi.org/10.1021/acs.joc.6b01378>.
10. Jacob, J.; Varghese, N.; Rasheed, S.P.; Agnihotri, S.; Sharma, V.; Wakode, S. Recent advances in the synthesis of isoquinoline and its analogue: A review. *WJPPS* **2016**, *5*, 1821–1837. <http://doi.org/10.20959/wjpps20167-7251>.
11. Yang, D.; Burugupalli, S.; Daniel, D.; Chen, Y. Microwave-assisted one-pot synthesis of isoquinolines, furopyridines, and thienopyridines by palladium-catalyzed sequential Coupling–Imination–Annulation of 2-bromoarylaldehydes with terminal acetylenes and ammonium acetate. *J. Org. Chem.* **2012**, *77*, 4466–4472. <http://doi.org/10.1021/jo300494a>.
12. Zheng, L.; Ju, J.; Bin, Y.; Hua, R. Synthesis of isoquinolines and heterocycle-fused pyridines via three-component cascade reaction of aryl ketones, hydroxylamine, and alkynes. *J. Org. Chem.* **2012**, *77*, 5794–5800. <http://doi.org/10.1021/jo3010414>.

13. Chinnagolla, R.K.; Pimparkar, S.; Jeganmohan, M. Ruthenium-Catalyzed Highly Regioselective Cyclization of Ketoximes with Alkynes by C–H Bond Activation: A Practical Route to Synthesize Substituted Isoquinolines. *Org. Lett.* **2012**, *14*, 3032–3035. <http://doi.org/10.1021/ol301091z>.
14. Clarke, P.A.; Santos, S.; Martin, W.H.C. Combining pot, atom and step economy (PASE) in organic synthesis. Synthesis of tetrahydropyran-4-ones. *Green Chem.* **2007**, *9*, 438–440. <http://doi.org/10.1039/B700923B>.
15. Arends, I.; Sheldon, R.; Hanefeld, U. *Green Chemistry and Catalysis*; Wiley-VCH: Weinheim, Germany, 2007.
16. Charushin, V.N.; Chupakhin, O.N. Nucleophilic C–H functionalization of arenes: A contribution to green chemistry. *Russ. Chem. Bull.* **2019**, *68*, 453–471. <http://doi.org/10.1007/s11172-019-2441-3>.
17. Chupakhin, O.N.; Charushin, V.N. Recent advances in the field of nucleophilic aromatic substitution of hydrogen. *Tetrahedron Lett.* **2016**, *57*, 2665–2672. <http://doi.org/10.1016/j.tetlet.2016.04.084>.
18. Charushin, V.N.; Chupakhin, O.N. Metal-free C–H functionalization of aromatic compounds through the action of nucleophilic reagents. *Top. Heterocycl. Chem.* **2014**, *37*, 1–50. http://doi.org/10.1007/7081_2013_119.
19. Gulevskaya, A.V.; Pozharskii, A.F. The S_N^H -Amination of Heteroaromatic Compounds. *Top. Heterocycl. Chem.* **2013**, *37*, 179–239. http://doi.org/10.1007/7081_2013_114.
20. Budyka, M.F.; Terent'ev, P.B.; Kost, A.N. Amination of 5-azacinnoline with aromatic amines. *Chem. Heterocycl. Compd.* **1978**, *14*, 663–666. <http://doi.org/10.1007/BF00481143>.
21. Borovlev, I.V.; Demidov, O.P.; Saigakova, N.A.; Amangasieva, G.A. S_N^H - and S_N^{ipso} -Arylation of 1, 3, 7-Triazapyrenes. *Eur. J. Org. Chem.* **2014**, *2014*, 7675–7683. <http://doi.org/10.1002/ejoc.201402891>.
22. Shchepochkin, A.V.; Chupakhin, O.N.; Charushin, V.N.; Steglenko, D.V.; Minkin, V.I.; Rusinov, G.L.; Matern, A.I. C–H functionalization of azines. Anodic dehydroaromatization of 9-(hetero) aryl-9, 10-dihydroacridines. *RSC Adv.* **2016**, *6*, 77834–77840. <http://doi.org/10.1039/C6RA17783B>.
23. Makhava, G.F.; Lushchekina, S.V.; Boltneva, N.P.; Serebryakova, O.G.; Rudakova, E.V.; Ustyugov, A.A.; Bachurin, S.O.; Shchepochkin, A.V.; Chupakhin, O.N.; Charushin, V.N.; et al. 9-Substituted acridine derivatives as acetylcholinesterase and butyrylcholinesterase inhibitors possessing antioxidant activity for Alzheimer's disease treatment. *Bioorg. Med. Chem.* **2017**, *25*, 5981–5994. <http://doi.org/10.1016/j.bmc.2017.09.028>.
24. Shchepochkin, A.V.; Chupakhin, O.N.; Charushin, V.N.; Rusinov, G.L.; Subbotina, Y.O.; Slepukhin, P.A.; Budnikova, Y.G. Stable σ^H -adducts in the reactions of the acridinium cation with heterocyclic N-nucleophiles. *Russ. Chem. Bull.* **2013**, *62*, 773–779. <http://doi.org/10.1007/s11172-013-0105-2>.
25. Gulevskaya, A.V.; Tyaglivaya, I.N.; Verbeeck, S.; Maes, B.U.W.; Tkachuk, A.V. Oxidative arylation of 1, 3-dinitrobenzene and 3-nitropyridine under anaerobic conditions: The dual role of the nitroarenes. *Arkivoc* **2011**, *2011*, 238–251. <http://doi.org/10.3998/ark.5550190.0012.917>.
26. Garnier, E.; Audoux, J.; Pasquinet, E.; Suzenet, F.; Poullain, D.; Lebre, B.; Guillaumet, G. Easy access to 3-or 5-heteroarylamino-1, 2, 4-triazines by S_N^{Ar} , S_N^H , and palladium-catalyzed N-heteroarylations. *J. Org. Chem.* **2004**, *69*, 7809–7815. <http://doi.org/10.1021/jo0490898>.
27. Matern, A.I.; Charushin, V.N.; Chupakhin, O.N. Progress in the studies of oxidation of dihydropyridines and their analogues. *Russ. Chem. Rev.* **2007**, *76*, 23–40. <http://doi.org/10.1070/RC2007v076n01ABEH003647>.
28. Makosza, M.; Wojciechowski, K. Nucleophilic Substitution of Hydrogen in Arenes and Heteroarenes. *Top. Heterocycl. Chem.* **2013**, *37*, 51–105. http://doi.org/10.1007/7081_2013_115.
29. Makosza, M. How does nucleophilic aromatic substitution in nitroarenes really proceed: General mechanism. *Synthesis* **2017**, *49*, 3247–3254. <http://doi.org/10.1055/s-0036-1588444>.
30. Wróbel, Z.; Kwast, A. 2-Nitroso-N-arylanilines: Products of acid-promoted transformation of σ^H adducts of arylamines and nitroarenes. *Synlett* **2007**, *10*, 1525–1528. <http://doi.org/10.1055/s-2007-982534>.
31. Wróbel, Z.; Kwast, A. Simple synthesis of N-aryl-2-nitrosoanilines in the reaction of nitroarenes with aniline anion derivatives. *Synthesis* **2010**, *2010*, 3865–3872. <http://doi.org/10.1055/s-0030-1258230>.
32. Kwast, A.; Stachowska, K.; Trawczyński, A.; Wróbel, Z. N-Aryl-2-nitrosoanilines as intermediates in the synthesis of substituted phenazines from nitroarenes. *Tetrahedron Lett.* **2011**, *52*, 6484–6488. <http://doi.org/10.1016/j.tetlet.2011.09.113>.
33. Wróbel, Z.; Stachowska, K.; Grudzień, K.; Kwast, A. N-Aryl-2-nitrosoanilines as Intermediates in the Two-Step Synthesis of Substituted 1, 2-Diarylbenzimidazoles from Simple Nitroarenes. *Synlett* **2011**, *10*, 1439–1443. <http://doi.org/10.1055/s-0030-1260764>.
34. Wróbel, Z.; Wieclaw, M.; Bujok, R.; Wojciechowski, K. Synthesis of pyrrolo [3, 2-a] phenazines from 5-nitroindoles and anilines. *Monatsh. Chem.* **2013**, *144*, 1847–1853. <http://doi.org/10.1007/s00706-013-1087-3>.
35. Bashkin, J.K.; Rains, R.; Stern, M. Taking green chemistry from the laboratory to chemical plant. *Green Chem.* **1999**, *1*, G41–G43. <http://doi.org/10.1039/GC990G41>.
36. Triplett, R.D.; Rains, R.K. Process for Preparing 4-aminodiphenylamine Intermediates. US Patent 7504539 B2, 17 March 2009.
37. Patriciu, O.-I.; Finaru, A.-L.; Sandulescu, I.; Guillaumet, G. Synthesis of Nitro N, N'-Dipyridinylamines via Oxidative Nucleophilic Substitution of Hydrogen. *Synthesis* **2007**, *2007*, 3868–3876. <http://doi.org/10.1055/s-2007-990896>.
38. Wozniak, M.; Baranski, A.; Nowak, K.; Poradowska, H. Regioselectivity of the amination of some nitroisoquinolines by liquid ammonia/potassium permanganate. *Liebigs Ann. Chem.* **1990**, *1990*, 653–657. <http://doi.org/10.1002/jlac.1990199001124>.
39. Woźniak, M.; Nowak, K. Amination of some nitroisoquinolines with liquid methylamine/potassium permanganate. *Liebigs Ann. Chem.* **1994**, *1994*, 355–360. <http://doi.org/10.1002/jlac.199419940407>.

40. Demidov, O.P.; Pobedinskaya, D.Y.; Avakyan, E.K.; Amangasieva, G.A.; Borovlev, I.V. S_N^H Arylamination of Nitroquinolines: Access to Nitro and Nitroso Derivatives of Arylaminoquinolines. *Chem. Heterocycl. Compd.* **2018**, *54*, 875–886. <http://doi.org/10.1007/s10593-018-2368-x>.
41. Stern, M.K.; Cheng, B.K. Amination of nitrobenzene via nucleophilic aromatic substitution for hydrogen: Direct formation of aromatic amide bonds. *J. Org. Chem.* **1993**, *58*, 6883–6888. <http://doi.org/10.1021/jo00076a059>.
42. Stern, M.K.; Bashkin, K.J. Method of preparing 4-aminodiphenylamine. US Patent 5 117 063, 26.05.1992.
43. Borovlev, I.V.; Demidov, O.P.; Kurnosova, N.A.; Amangasieva, G.A.; Avakyan, E.K. Direct oxidative S_N^H amidation of 1,3,7-triazapyrene. *Chem. Heterocycl. Compd.* **2015**, *51*, 170–175. <http://doi.org/10.1007/s10593-015-1677-6>.
44. Demidov, O.P.; Borovlev, I.V.; Amangasieva, G.A.; Avakyan, E.K. Oxidative S_N^H amidation of acridine and tautomerism of N-(acridin-9-yl) benzamides. *Chem. Heterocycl. Compd.* **2016**, *52*, 104–109. <http://doi.org/10.1007/s10593-016-1841-7>.
45. Amangasieva, G.A.; Borovlev, I.V.; Demidov, O.P.; Avakyan, E.K.; Borovleva, A.A. Synthesis of Amides by Nucleophilic Substitution of Hydrogen in 3-Nitropyridine. *Russ. J. Org. Chem.* **2018**, *54*, 867–872. <http://doi.org/10.1134/S1070428018060076>.
46. Amangasieva, G.A.; Avakyan, E.K.; Demidov, O.P.; Borovleva, A.A.; Pobedinskaya, D.Y.; Borovlev, I.V. S_N^H Amidation of nitroquinolines: Synthesis of amides on the basis of nitro- and nitrosoquinolines. *Chem. Heterocycl. Compd.* **2019**, *55*, 623–631. <http://doi.org/10.1007/s10593-019-02508-3>.
47. Amangasieva, G.A.; Borovlev, I.V.; Demidov, O.P.; Kurnosova, N.A.; Avakyan, E.K. Urea in an aminodemethoxylation reaction of 6-methoxy-1,3,7-triazapyrenes. *Chem. Heterocycl. Compd.* **2015**, *51*, 586–588. <http://doi.org/10.1007/s10593-015-1743-0>.
48. Borovlev, I.V.; Demidov, O.P.; Amangasieva, G.A.; Avakyan, E.K.; Kurnosova, N.A. Ureas as new nucleophilic reagents for S_N^H amination and carbamoyl amination reactions in the 1,3,7-triazapyrene series. *ARKIVOC*, **2016**, *2016*, 58–70. <http://doi.org/10.3998/ark.5550190.p009.412.48>.
49. Borovlev, I.V.; Demidov, O.P.; Amangasieva, G.A.; Avakyan, E.K.; Kurnosova, N.A. Ureas as a New Nucleophilic Reagents for S_N^Ar Amination and Carbamoyl Amination Reactions in 1,3,7-Triazapyrene Series. *J. Heterocycl. Chem.* **2017**, *54*, 406–412. <http://doi.org/10.1002/jhet.2597>.
50. Borovlev, I.V.; Demidov, O.P.; Amangasieva, G.A.; Avakyan, E.K. Direct and facile synthesis of 9-aminoacridine and acridin-9-yl-ureas. *Tetrahedron Lett.* **2016**, *57*, 3608–3611. <http://doi.org/10.1016/j.tetlet.2016.06.103>.
51. Avakyan, E.K.; Borovlev, I.V.; Demidov, O.P.; Amangasieva, G.A.; Pobedinskaya, D.Y. S_N^H Alkyl carbamoyl amination of 3-nitropyridine: Competitive synthesis of nitro- and nitrosopyridine derivatives. *Chem. Heterocycl. Compd.* **2017**, *53*, 1207–1213. <http://doi.org/10.1007/s10593-018-2201-6>.
52. Demidov, O.P.; Amangasieva, G.A.; Avakyan, E.K.; Borovlev, I.V. Nucleophilic Addition of Amides to 10-Alkylacridinium Cations: A Case of Double N-Nucleophilicity of Some Monoamides. *Synthesis* **2017**, *49*, 3710–3719. <http://doi.org/10.1055/s-0036-1588786>.
53. Iriepa, I.; Bellanato, J. Synthesis, spectroscopic, structural and conformational study of some tri-substituted ureas derived from N-methylpiperazine containing phenyl and N-heterocyclic substituents. *J. Mol. Struct.* **2013**, *1044*, 215–220. <http://doi.org/10.1016/j.molstruc.2013.01.001>.
54. Annese, M.; Corradi, A.B.; Forlani, L.; Rizzoli, C.; Sgarabotto, P. Tautomerism in some acetamido derivatives of nitrogen-containing heterocycles: X-ray structural analysis of 2-amino and 2-imino forms of benzothiazole derivatives. *J. Chem. Soc. Perkin Trans. 2* **1994**, 615–621. <http://doi.org/10.1039/P29940000615>.
55. Forlani, L.; Mezzina, E.; Boga, C.; Forconi, M. Tautomerism and Dimerization of Acetamidothiazole Derivatives—UV/Vis and NMR Spectroscopic Investigation. *Eur. J. Org. Chem.* **2001**, *2001*, 2779–2785. [http://doi.org/10.1002/1099-0690\(200107\)2001:14<2779::AID-EJOC2779>3.0.CO;2-N](http://doi.org/10.1002/1099-0690(200107)2001:14<2779::AID-EJOC2779>3.0.CO;2-N).
56. Gottlieb, H.E.; Kotlyar, V.; Nudelman, A. NMR chemical shifts of common laboratory solvents as trace impurities. *J. Org. Chem.* **1997**, *62*, 7512–7515. <http://doi.org/10.1021/jo971176v>.
57. Sharp, J.T.; Gosney, I.; Rowley, A.G. *Practical Organic Chemistry*; Springer: London, UK; New York, NY, USA, 1989; p. 57.