

Supporting Information

For

Effect of Controlling Thiophene Rings on D-A Polymer Photocatalysts Accessed via Direct Arylation for Hydrogen Production

Dongnai Ye ^{1,2}, Lei Liu ¹, Qimin Peng ¹, Jiabin Qiu ^{1,*}, Hao Gong ¹, Aiguo Zhong ³ and Shiyong Liu ^{1,*}

¹ Jiangxi Provincial Key Laboratory of Functional Molecular Materials Chemistry, College of Materials, Metallurgical and Chemistry, Jiangxi University of Science and Technology, Ganzhou 341000, China; yedongnai2014@gnnu.edu.cn (D.Y.); 13125257083@139.com (L.L.); qm2474572831@163.com (Q.P.); 6720210821@mail.jxust.edu.cn (H.G.)

² School of Chemistry and Chemical Engineering, Gannan Normal University, Ganzhou 341000, China

³ Department of Pharmacy & Chemistry, Taizhou University, Taizhou 318000, China; zhongaiguo@tzc.edu.cn

* Correspondence: jiabinqiu@jxust.edu.cn (J.Q.); chelsy@jxust.edu.cn or chelsy@zju.edu.cn (S.L.)




Figure. S1. ¹H NMR spectra of DBCS in CDCl₃.




Figure. S2. ¹³C NMR spectra of DBCS in CDCl₃.




Figure. S3. CV curves of the as-prepared CPs.




Figure. S4. Tauc plots of the transformed Kubelka-Munk function vs energy of CPs.