



Article Identification and Pharmacological Characterization of a Low-Liability Antinociceptive Bifunctional MOR/DOR Cyclic Peptide

Yangmei Li^{1,*}, Shainnel O. Eans², Michelle Ganno-Sherwood³, Abbe Eliasof¹, Richard A. Houghten³ and Jay P. McLaughlin^{2,*}

- ¹ College of Pharmacy, University of South Carolina, Columbia, SC 29208, USA; aeliasof@email.sc.edu
- ² Department of Pharmacodynamics, College of Pharmacy, University of Florida, Gainesville, FL 32610, USA; shaieans@cop.ufl.edu
- ³ Torrey Pines Institute for Molecular Studies, Port St. Lucie, FL 34987, USA; condor1082@aol.com (M.G.-S.); houghten@tpims.org (R.A.H.)
- * Correspondence: liyangme@cop.sc.edu (Y.L.); jmclaughlin@cop.ufl.edu (J.P.M.); Tel.: +1-803-777-0849 (Y.L.); Tel.: +1-352-273-7207 (J.P.M.)

Abstract: Peptide-based opioid ligands are important candidates for the development of novel, safer, and more effective analgesics to treat pain. To develop peptide-based safer analgesics, we synthesized a mixture-based cyclic pentapeptide library containing a total of 24,624 pentapeptides and screened the mixture-based library samples using a 55 °C warm water tail-withdrawal assay. Using this phenotypic screening approach, we deconvoluted the mixture-based samples to identify a novel cyclic peptide Tyr-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt), which produced dose- and timedependent antinociception with an ED_{50} (and 95% confidence interval) of 0.70 (0.52–0.97) mg/kg *i.p.* mediated by the mu-opioid receptor (MOR). Additionally, higher doses (\geq 3 mg/kg, *i.p.*) of CycloAnt antagonized delta-opioid receptors (DOR) for at least 3 h. Pharmacological characterization of CycloAnt showed the cyclic peptide did not reduce breathing rate in mice at doses up to 15 times the analgesic ED₅₀ value, and produced dramatically less hyperlocomotion than the MOR agonist, morphine. While chronic administration of CycloAnt resulted in antinociceptive tolerance, it was without opioid-induced hyperalgesia and with significantly reduced signs of naloxone-precipitated withdrawal, which suggested reduced physical dependence compared to morphine. Collectively, the results suggest this dual MOR/DOR multifunctional ligand is an excellent lead for the development of peptide-based safer analgesics.

Keywords: mixed-based combinatorial library; structure–activity relationships; cyclic peptide; mu-opioid receptor; delta-opioid receptor; bifunctional ligand; antinociception; opioid antagonist; opioid liabilities

1. Introduction

The perception of pain arises from a variety of nociceptive insults, each transduced and transmitted through distinct systems to which specific analgesics may provide varying degrees of relief [1]. Opioid analgesics have a long history of use to relieve moderate to extreme pain [1]. However, repeated opioid use results in clinical liabilities such as the development of tolerance and dependence and a potentially lethal respiratory depression [2]. The misuse of opioid analgesics leads to the increasing risk of opioid abuse and overdose, which have contributed to the escalating economic burden imposed on society. Thus, there is a critical public health need for novel, safer, and more efficacious analgesics to treat pain.

One strategy to address this need focuses on the development of opioid peptides into safer analgesics [3–5]. The opioid system utilizes opioid receptors and the endogenous opioid peptides to modulate pain, yet the endogenous opioid system is not observed



Citation: Li, Y.; Eans, S.O.; Ganno-Sherwood, M.; Eliasof, A.; Houghten, R.A.; McLaughlin, J.P. Identification and Pharmacological Characterization of a Low-Liability Antinociceptive Bifunctional MOR/DOR Cyclic Peptide. *Molecules* 2023, 28, 7548. https://doi.org/ 10.3390/molecules28227548

Academic Editor: Nuno Vale

Received: 13 October 2023 Revised: 4 November 2023 Accepted: 9 November 2023 Published: 11 November 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). to produce respiratory depression and addictive side effects. A recent study using a genetically encoded biosensor revealed the mu-opioid peptide receptor (MOR) and the delta-opioid peptide receptor (DOR) not only function at the cell membrane, but also inside the cell at different locations, depending on the chemistry of ligand types [6]. After peptides activated the MOR in the plasma membrane, MOR quickly internalized and was detected in an active state in endosome for 20 min, while opiate alkaloids penetrated the membrane to produce Golgi-localized receptor activation after MOR was activated in the plasma membrane [6]. The spatiotemporal specificity is thought to affect signal duration and pathway selection downstream, contributing to distinct downstream physiological effects [7,8] As the endogenous opioid peptides, but not opiate alkaloids, can precisely modulate pain, the different spatiotemporal patterns of opioid receptor activation may link to the severity of adverse effects limiting opiate use in pain management [9].

The endogenous opioid peptides such as enkephalins, endorphins, and dynorphins exert their analgesic effects through the MOR, DOR, and kappa (KOR) opioid receptors. These peptides have varying affinities at the opioid receptors and none of them bind exclusively to one opioid receptor type [10]. For example, enkephalins bind and activate the DOR and MOR. Endorphins exert their effect primarily through the MOR, but they also have affinity for the DOR. While dynorphins have high affinity for the KOR, they have significant affinity for the MOR and DOR as well. Most importantly, as a whole, the endogenous opioid peptides activate all opioid receptors.

All three opioid receptors mediate opioid-dependent analgesia. While MOR has been the primary target for clinically used opioid analgesics, activation of MOR may lead to respiratory depression and physical dependence. Unlike MOR, activation of DOR does not cause respiratory depression and physical dependence [11]. Since DOR is mainly located in the cytoplasm, with a very low level of receptors being associated with the plasma membrane [12], activation of DOR alone has been reported to produce insufficient analgesia in a model of acute pain [11]. However, mounting evidence indicates that the activation of DOR produces analgesia in various models of chronic inflammatory and neuropathic pain [13,14].

Importantly, *multifunctional* ligands, which simultaneously modulate MOR and DOR, have been demonstrated to produce enhanced antinociception with fewer adverse effects [15–20]. For example, co-administration of the DOR agonist SNC80 with the MOR agonist heroin enhanced the antinociceptive effect of heroin while decreasing the sedative and reinforcing effects [21]. Dual agonists effective at both MOR and DOR, such as MMP-2200 [22,23] and SRI-22141 [24], produce high antinociceptive potency as well as decreasing the propensity to produce locomotor stimulation, tolerance, and physical dependence in mouse models of chronic pain. Simultaneous activation of MOR and DOR using a dual agonist LP2 not only significantly ameliorated mechanical allodynia, but also produced neuroprotective effects in a rodent model of neuropathic pain [25]. Interestingly, dual-function ligands acting as a MOR agonist/DOR antagonist have also been reported to produce enhanced antinociceptive efficacy with reduced tolerance and dependence [15,26,27]. DIPP-NH₂[Ψ] [28], SoRi20411 [29], and 14-alkoxy pyridomorphinans [30] are potent DOR antagonists and MOR agonists which produce analgesia with reduced tolerance when administered repeatedly at the ED_{80} antinociceptive dose supraspinally twice daily for 5–7 days. Ligands simultaneously modulating multiple opioid receptors, such as MOR and DOR, therefore hold great promise for the development of safe analgesics.

We have previously reported the synthesis of a mixture-based cyclic peptide library consisting of 24,624 anthraniloyl(Ant)-tagged pentapeptides in a positional scanning format (Figure 1) [31]. The fully defined cyclic peptide Tyr-[D-Lys-Dap(Ant)-Tyr-Gly] identified from this library demonstrated high affinity for the MOR in a competition receptor-binding assay [31]. As cyclic peptides have improved stability against proteolysis, we hypothesized a cyclic peptide with a desired antinociceptive, safety, and pharmacokinetic profile would be found within the library. To identify such a cyclic peptide analgesic from this library, we employed a direct phenotypic *in vivo* screening using the 55 °C warm water

tail-withdrawal (WWTW) assay to facilitate the discovery of the analgesic lead [32,33]. As the WWTW assay measures the overall antinociception of a sample in mice, the antinociceptive lead compounds discovered should have a favored pharmacokinetic profile without a preconceived bias for the mechanism of action [34]. Using this direct *in vivo* screening, we identified a fully defined, novel, potent, low-liability cyclic peptide analgesic from the mixture library. Further target identification characterization using opioid receptor knock-out mice elucidated the analgesic lead produced agonist activity at MOR, while pharmacological testing demonstrated DOR antagonism, indicating that it acted as a MOR agonist/DOR antagonist in mice.



-																														-						
R ² sample	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
O at R ²	А	D	Е	F	G	н	Т	К	L	М	Ν	Ρ	Q	R	S	Т	V	w	Υ	а	d	е	f	h	i	k	Т	m	n	р	q	r	s	t	v	у
R ⁴ sample	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72
O at R ⁴	А	D	Е	F	G	Н	Т	К	L	М	Ν	Ρ	Q	R	S	Т	V	W	Y	а	d	е	f	h	i	k	T	m	n	р	q	r	s	t	v	y
R ⁵ sample	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91																	
O at R ⁵	А	D	Е	F	G	н	1	к	L	М	Ν	Ρ	Q	R	s	Т	V	w	Υ																	

Figure 1. The mixture-based cyclic peptide library and library composition. (**Top**). General peptide sequence in the library. (**Bottom**). The defined single amino acid (*O*) of each sample in the library.

2. Results

2.1. Mixture-Based Cyclic Peptide Library

As we reported previously, the cyclic peptides in the library [31] were synthesized from peptide thioesters [35] using an imidazole-promoted cyclization strategy [36]. All the pentapeptides in this library have a Gly fixed at R¹ position and a Dap(Ant) at fixed R³ position. The rest of the R², R⁴, and R⁵ positions are made either with a single amino acid (*O*) or a mixture of amino acids (*X*) [31]. The library was composed of the R²-, R⁴-, and R⁵-sub-libraries (Figure 1). R²-position and R⁴-position were constituted by a total of 36 amino acids (AAs) including 19 *L*- and 17 *D*-amino acids, while R⁵-position was generated with 19 natural *L*-amino acids. Thus, there are 36 R² samples in the R²-sublibrary, 36 R⁴ samples in the R⁴-sublibrary, and 19 R⁵ samples in the R⁵-sublibrary; each sample is a mixture of pentapeptides. Each sample contains a total of 684 (36 × 19) pentapeptides at the R²- and R⁴-sublibrary has a total of 91 (36 + 36 + 19) samples and 24,624 (36 × 36 × 19) pentapeptides in the linear sequences.

2.2. Library Deconvolution Using Direct In Vivo Screening

Utilizing a positioning scanning format, the library was screened *in vivo* via antinociception assessed with the 55 °C WWTW assay, as we previously described [32,33]. Each partially defined, mixture-based sample was administered intraperitonially (*i.p.*) at a dose of 5 mg/kg. Additional mice received vehicle alone (10%DMSO/90% saline, *i.p.*) as a negative control, or morphine (10 mg/kg, *i.p.*) as a positive control. The combined time mice demonstrated for withdrawment of their tails for each sample tested was summed over six time points, and was reported by substitution position (Figure 2A–C). Analysis of the responses with one-way ANOVA detected significant global responses in the R²-(F_(37,266) = 12.1, *p* < 0.0001; Figure 2A), R⁴-(F_(37,266) = 13.1, *p* < 0.0001; Figure 2B) and R⁵-sublibraries (F_(25,182) = 9.50, *p* < 0.001; Figure 2C). Post hoc analysis (Dunnett's) identified samples from the R²-, R⁴- and R⁵-sublibraries producing a combined average response time to tail-withdrawal that was both

significantly different from the vehicle response (p < 0.05 vs. dotted line, Figure 2A–C), but not different from the morphine response (p > 0.05 vs. cyan bar, Figure 2A–C). Meeting these conditions for the R²- position were samples 5 (Gly), 6 (His), 11 (Asn), 12 (Pro), 13 (Gln), 16 (Thr) and 25 (D-Ile), whereas for the R⁴ position, only Sample 62 (D-Lys) showed this response, while the R⁵ position demonstrated three samples meeting the criteria: 75 (Glu), 81 (Leu) and 91 (Tyr).





at each position from in vivo screening												
R ²	Gly	His	Asn	Pro	Gln	Thr	D-lle					
\mathbf{R}^4	D-Lys											
R ⁵	Glu	Leu	Tyr									

Figure 2. Positional scan screening of the cyclic peptide library *in vivo* using the 55 °C warm water tail-withdrawal assay in young adult male C57BL/6J mice across an 8 h period. (**A**): library defined at position 2, (**B**): library defined at position 4, (**C**): library defined at position 5. Each sample was given to the mice (n = 8) *i.p.* at a dose of 5 mg/kg. The tail-withdrawal latencies were measured at six time points post-administration, with combined time to withdraw tails (s; y-axis) calculated by taking the sum of the average tail-withdrawal latencies from each time point. Dotted line = vehicle alone (*i.p.*); cyan line = morphine control (10 mg/kg, *i.p.*). (**D**): Summary of key functionality residues identified in the screening. Data represented the combined time to tail withdrawal \pm SEM for each sample. Dark blue bars in (**A**–**C**) represent significantly different from vehicle (p < 0.05), but not morphine (p > 0.05); One-way ANOVA with Dunnett's post hoc tests.

To validate the antinociceptive effects of the five strongest partially defined samples and to determine a potential dose–response relationship, additional doses of 10 mg/kg or 25 mg/kg, *i.p.* of the selected samples were administered to mice and tested *in vivo* using the 55 °C warm water tail-withdrawal assay (Figure 3). Samples 5, 6 and 16 defined at the R²-position produced dose-dependent antinociception that peaked at 60 min post administration (Figure 3A). Estimated from the limited doses tested, antinociceptive ED₅₀ (and 95% confidence intervals) values were near that of morphine (4.12 (2.73–5.82) mg/kg, *i.p.*), at 13.0 (10.2–17.8) mg/kg, *i.p.* (Sample 5), 17.5 (12.5–33.2) mg/kg, *i.p.* (Sample 6) and 7.15 (3.91–9.96) mg/kg, *i.p.* (Sample 16). Similarly, mixture-based samples 81 and 91 defined at the R⁵-position produced antinociceptive ED₅₀ (and 95% confidence intervals) values of 3.62 (0.00–8.58) mg/kg, *i.p.* (Sample 81) and 7.15 (0.52–12.7) mg/kg, *i.p.* (Sample 91) at their peak effect of 120 min post administration (Figure 3B). While limited to estimates, this data collectively confirms the antinociceptive effects of the partially defined, mixture-based samples, validating further deconvolution of the library. Taken together, Thr (16) at R², D-Lys (62) at R⁴, and Tyr (91) and Leu (81) at R⁵ were selected as the most active residues at each position to deconvolute the mixture-based library for the identification of the cyclic peptide that produced the most significant antinociceptive activity in mice with the 55 °C WWTW assay.



Figure 3. Antinociception produced by the top five mixture-based samples in the mouse 55 °C warm water tail-withdrawal assay is dose-dependent. (**A**) Dose–response of mixture-based samples 5, 6 and 16 defined at the \mathbb{R}^2 position, 60 min after administration; (**B**) Dose–response of mixture-based samples 81 and 91 defined at the \mathbb{R}^5 position, 120 min after administration. All samples were administered by the *i.p.* route. Morphine is included as a positive control, with antinociception measured at the 30 min peak effect. All points = 8 mice (7–11 mice for morphine).

2.3. Synthesis of Cyclic Peptides Tyr-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt) and Leu-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt-Leu) for the Deconvolution of the Library

Two cyclic peptides, Tyr-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt) and Leu-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt-Leu), were then synthesized to complete the deconvolution of the library. Using mercaptopropyl isobutyl polyhedral oligomeric silsesquioxane (POSS-SH) as a soluble support [37], Boc-protected amino acids including Gly (R¹), Thr(tBu) (R²), Dap[Ant(Boc)] (R³), D-Lys(Alloc) (R⁴), and Tyr(tBu)/Leu (R⁵) were subsequentially coupled to form the peptide POSS-thioesters, Boc-Tyr(tBu)-D-Lys(Alloc)-Dap(Ant)-Thr-Gly-S-POSS and Boc-Leu-D-Lys(Alloc)-Dap(Ant)-Thr-Gly-S-POSS, respectively (Scheme 1). The peptide POSS-thioesters were then treated with Pd(PPh₃)₄ in the presence of PhSiH₃ in DCM to remove the Alloc protective group, followed by adding imidazole to carry out a one-pot, in situ cyclization to form the final side-to-tail cyclic peptides, CycloAnt and CycloAnt-Leu, respectively [38].



Scheme 1. Solution phase synthesis of the cyclic peptides using POSS-SH as a soluble support.

The cyclic peptides were then treated with 55% TFA to remove the protective groups and the resulting compounds were purified by RP-HPLC and structurally characterized by NMR experiments before characterization of their pharmacological profile.

2.4. Screening of the Cyclic Peptide CycloAnt for Protein Interaction

The potential interaction of CycloAnt with a panel of 47 protein targets, including GPCRs, kinases, nuclear hormone receptors, ion channels, and enzymes, was assessed by assays included in the SAFETYscan service from Eurofins DiscoverX. CycloAnt was tested in both agonist and antagonist modes for each representative GPCRs (see Supplemental Material S9–S11). Tested in a cAMP assay utilizing cells overexpressing each receptor, CycloAnt demonstrated MOR and DOR agonism, with EC₅₀ values of 14.0 and 3.36 nM, respectively (Supplemental Material S9–S11). CycloAnt did not produce agonist or antagonist activity at any other target tested in this panel.

2.5. Evaluation of the Cyclic Peptides for Their Antinociceptive Effects in Wild-Type Mice

We tested the antinociceptive effects of the cyclic peptides using the 55 °C WWTW assay with C57BL/6J male mice (Figure 4). Both peptides produced dose- (Figure 4A) and time-dependent antinociception ($F_{(16,152)} = 38.5$; p < 0.0001; two-way ANOVA; Figure 4B). Peak antinociception was produced by both peptides within 20 min of administration, with a significant duration of 70 min for CycloAnt (3 mg/kg, *i.p.*) and 50 min for the CycloAnt-Leu analog (100 mg/kg, *i.p.*) compared to vehicle (p < 0.05, Dunnett's post hoc test; Figure 4B). The two cyclic peptides produced full antinociception, but with significant differences in potency. CycloAnt-Leu produced antinociception with an ED₅₀ (and 95% C.I.) value of 38.3 (18.7–127.1) mg/kg, *i.p.*, 9.3-fold less potent than the control opioid morphine (ED₅₀ (and 95% C.I.) value = 4.12 (2.73–5.82) mg/kg, *i.p.*; Figure 4A). In contrast, CycloAnt displayed an ED₅₀ (and 95% C.I.) value of 0.70 (0.52–0.97) mg/kg, *i.p.*, 5.9-fold more potent than morphine. Collectively, this data demonstrates CycloAnt is a potent antinociceptive "hit" identified through the direct phenotypic *in vivo* screening and deconvolution of this mixture-based cyclic peptide library.



Figure 4. Acute antinociceptive activity in the 55 °C WWTW assay following *i.p.* administration in C57BL/6J mice. Shown as (**A**) dose–response and (**B**) time-course of antinociception following a maximally efficacious dose of CycloAnt or CycloAnt-Leu. Morphine and vehicle are included as controls where appropriate. Points represent average % Antinociception \pm SEM of 6–11 mice for each compound presented.

2.6. Evaluation of Opioid Receptor Involvement in CycloAnt Antinociception

To assess the contributions of the individual opioid receptors to CycloAnt-mediated antinociception, the cyclic peptide (3 mg/kg, *i.p.*) was administered to mice lacking the mu- (MOR KO), kappa- (KOR KO) or delta-opioid receptor (DOR KO). Compared to the response in wild-type C57BL/6J mice, CycloAnt-mediated antinociception was significantly

reduced 80% only in MOR KO mice ($F_{(3,28)} = 216.3$, p < 0.0001, one-way ANOVA with Dunnett's post hoc test), and not KOR KO or DOR KO mice (Figure 5).



Figure 5. Evaluation of opioid receptor involvement in the antinociceptive activity of CycloAnt (3 mg/kg, *i.p.*) 20 min after administration to wild-type (C57BL/6 J) or mice where individual opioid receptors have been knocked out. * Significantly different from vehicle-control response (p < 0.05); + Significantly different from wild-type mice response (p < 0.0001); one-way ANOVA with Dunnett's post hoc test. Bars each represent average % Antinociception \pm SEM of 8 mice.

2.7. Evaluation of Opioid-Receptor-Selective Antagonism Mediated by CycloAnt

Given the MOR and DOR activity identified in the *in vitro* screening, and the singular MOR agonist activity displayed in mice, CycloAnt was evaluated for antagonist activity against the KOR-selective agonist U50,488 (10 mg/kg, *i.p.*), and the DOR-selective agonist SNC-80 (100 nmol, *i.c.v.*; Figure 6A). MOR KO mice were used to preclude complications of the mu-opioid receptor-mediated antinociception. U50,488 antinociception was not significantly antagonized by pretreatment with CycloAnt (3 or 10 mg/kg, *i.p.*; $F_{(2,21)} = 0.9$, p = 0.42, one-way ANOVA; Figure 6A left bars). However, surprisingly, CycloAnt pretreatment significantly antagonized SNC-80 antinociception in a dose-dependent manner ($F_{(2,21)} = 22.3$, p < 0.0001, one-way ANOVA with Dunnett's post hoc testing; Figure 6A, right bars). Examined at the dose of maximal DOR antagonism (10 mg/kg, *i.p.*), CycloAntmediated DOR antagonism was time dependent ($F_{(4,43)} = 16.2$, p < 0.0001, one-way ANOVA with Dunnett's post hoc testing; Figure 6B), with significant reduction of SNC-80 antinociception for 1 or 3 h (p < 0.0001), but not 6 h (p = 0.07) or 8 h (p = 0.96). This data suggests CycloAnt acts *in vivo* as a multifunctional MOR agonist and DOR antagonist.

2.8. Evaluation of CycloAnt for Respiratory and Hyperlocomotor Effects

Respiratory depression is a major life-threatening effect of clinically used opioid analgesics. To evaluate whether the cyclic peptide-based MOR/DOR multifunctional ligand can dissociate this side effect from antinociception, we examined CycloAnt for its potential to depress breathing rate over a 120 min period using the automated, computer-controlled Comprehensive Lab Animal Monitoring System (CLAMS) apparatus in male C57BL/6J mice [34]. As expected, the positive control morphine (10 or 30 mg/kg, *i.p.*) produced significant dose and time-dependent respiratory depression compared to vehicle $F_{(20,360)} = 6.67$, p < 0.001, two-way RM ANOVA with Tukey's multiple comparison post hoc test; Figure 7A). In contrast, CycloAnt did not significantly impair breathing when administrated *i.p.* at either the peak antinociceptive dose (3.0 mg/kg) or a higher dose 15 times the antinociceptive ED₅₀ value (10.6 mg/kg), and in fact statistically increased it in a time and dose-dependent manner up to 100 min compared to vehicle (p < 0.05, Tukey's test; Figure 7A).



Figure 6. Evaluation of opioid antagonist activity of CycloAnt in the 55 °C warm water tailwithdrawal assay. MOR KO mice were tested to avoid confounding antinociception mediated by this receptor. (**A**) A 60 min pretreatment with CycloAnt (3 or 10 mg/kg, *i.p.*) dose-dependently prevented SNC-80-induced antinociception (100 nmol, *i.c.v.*), but not that of U50,488 (10 mg/kg, *i.p.*). (**B**) Time-dependent DOR antagonism by CycloAnt. The antinociceptive effect of SNC-80 (100 nmol, *i.c.v.*) was determined in mice pretreated for 1, 3, 6 or 8 h with CycloAnt (10 mg/kg, *i.p.*). Points represent average % antinociception \pm SEM from 6 to 12 mice for each bar. *significantly different from response of agonist alone (*p* < 0.05); one-way ANOVA with Dunnett's post hoc test.



Figure 7. Effects of the hit on (**A**) respiration and (**B**) ambulation in C57BL/6J mice tested in the CLAMS/Oxymax system. Respiratory and ambulation were monitored after *i.p.* administration of vehicle (saline, i.p.), morphine (10 or 30 mg/kg, *i.p.*), or CycloAnt (3 or 10.6 mg/kg, *i.p.*). Data from 12-20 mice presented as % vehicle \pm SEM; breaths per minute, BPM (**A**) or ambulation, XAMB (**B**). Dashed lines represent normalized vehicle response. * Significantly different from vehicle control response (p < 0.05); \pm Significantly different from response of morphine at either dose (p < 0.05); two-way RM ANOVA with Tukey's multiple comparison post hoc test.

As a MOR agonist, morphine induces psychostimulation demonstrated by hyperlocomotion in mice. Consistent with this, treatment significantly increased ambulation (factor: time × treatment, $F_{(20,360)} = 36.7$, p < 0.0001, two-way RM ANOVA; Figure 7B), primarily due to morphine (factor: treatment, $F_{(10,250)} = 34.8$, p < 0.001, two-way RM ANOVA). In contrast, CycloAnt induced significant hyperlocomotor activity (factor: treatment, $F_{(10,205)} = 3.19$, p = 0.0008, two-way RM ANOVA with Tukey's test; Figure 7B), but CycloAnt-induced ambulations were significantly less than those induced by morphine after 20 min to the end of the testing period († p < 0.05, Tukey's test; Figure 7B).

2.9. Evaluation of CycloAnt for Potential of Opioid-Induced Hyperalgesia (OIH), Tolerance and Dependence Development

Chronic use of opioid analgesics may cause opioid-induced hyperalgesia, the paradoxical response where a patient becomes more sensitive to painful stimuli, as well as antinociceptive tolerance and physical dependence. To assess this, mice were treated twice daily for four days with the morphine (10 mg/kg, *i.p.*) or CycloAnt (3 mg/kg, *i.p.*), and once more on the 5th day before WWTW testing. Tail-withdrawal latencies were collected both prior to the start of treatment (on day 1), and again on the morning of day 5 prior to drug treatment. Each group of mice showed equivalent initial responses on day 1 to a 48 °C warm water stimulus (morphine, 10 ± 1.14 s vs. CycloAnt, 9.1 ± 0.68 s; p = 0.52, Student's *t*-test; Figure 8A) or 55 °C warm water stimulus (morphine, 1.43 ± 0.05 s vs. CycloAnt, 1.56 ± 0.08 s; p = 0.20, Student's *t*-test), whereas chronic morphine administration induced significant hyperalgesia ($F_{(1,17)}$ = 14.3, p = 0.002; two-way RM ANOVA) against a 48 °C warm water stimulus on day 5 (with a tail withdrawal latency of 4.35 ± 0.47 s; *p* < 0.0001, Sidak's post hoc test; Figure 8A), chronic CycloAnt treatment did not induce significant OIH (day 5 tail withdrawal latency of 9.02 \pm 0.83 s; *p* = 0.997, Sidak's post hoc test; Figure 8A). However, repeated chronic administration of either morphine or CycloAnt over 4 days produced significant reductions in the tail-withdrawal latency of the compound on day 5 ($F_{(1,18)}$ = 5.75, p = 0.03; two-way REML ANOVA with Sidak's post hoc test; Figure 8B), indicating opioid-induced tolerance.



Figure 8. Assessment of chronic treatment on (**A**) induced hyperalgesia in the 48 °C warm water tail-withdrawal test, or (**B**) antinociceptive tolerance in the 55 °C warm water tail-withdrawal test. Mice were first tested on day 1 to establish baseline responses prior to any treatment. Mice were then subjected to a dosing regimen with morphine (10 mg/kg, *i.p.*, twice a day; blue bars) or CycloAnt (3 mg/kg, *i.p.*, twice a day; red bars) for 4 days. On day 5, tail-withdrawal latency from 48 °C water was assessed prior to final dosing (**A**), whereas tail-withdrawal latency from 55 °C water was tested 20 min (for CycloAnt, 3 mg/kg, *i.p.*) or 30 min (for morphine, 10 mg/kg, *i.p.*) after drug administration (**B**). Data are shown as mean \pm SEM of (**A**) pre- and post-dosing testing or (**B**) responses to listed drug in naïve and chronically treated mice. * *p* < 0.01 versus baseline (**A**) or response in naïve mice (**B**), Two-way RM ANOVA with Sidak's multiple comparisons post hoc test. *n* = 9–11 mice/treatment.

We then examined the potential development of physical dependence after chronic administration of morphine or CycloAnt, as above, followed by precipitation of withdrawal with naloxone. Repeated treatment with escalating doses of morphine induced physical dependence, demonstrated by administration of naloxone (Table 1). Morphine produced significant increases in the frequency of forepaw tremor, presence of diarrhea, and frequency of jumping and teeth chattering, along with decreases in rearing and the forepaw licking frequency characteristic of opioid physical dependence (see one-way ANOVA values in Table 1). In contrast, while CycloAnt showed significant increases in diarrhea and reduced rearing frequency compared to saline control mice, most other signs of withdrawal, such as jumping frequency and teeth chattering, significantly differed from those of morphine or were completely absent (Table 1). Collectively, these results indicate chronic CycloAnt treatment induced much less physical dependence than that seen with morphine.

Table 1. Assessment of physical dependence in mice treated chronically with CycloAnt, Saline or Morphine and subjected to naloxone-precipitated withdrawal. \ddagger Saline values reproduced from Wilson et al., 2021 [39]. Average and Standard Error of the Mean (or presence of diarrhea) reported for n = 9-10 mice/treatment. * p < 0.05 versus Saline control, $\ddagger p < 0.05$ versus Morphine control (Tukey's post hoc test).

	Tr	eatment (D	Outcome						
Measure	CycloAn	t (<i>n</i> = 9)	Saline (#	n = 10) ‡	Morphine	(n = 10)	(1-Way ANOVA)		
	Ave	SEM	Ave	SEM	Ave	SEM	$F_{(2,26)} =$		
Forepaw Tremor	23.7 +	6.85	20.2	80.4	66 *	17.0	4.77, <i>p</i> = 0.02		
Wet Dog Shakes	2.22	0.43	0.8	0.51	1.3	0.56	1.95, p = 0.16		
Straightening	1.22	0.47	5.7	2.33	1.1	0.46	3.35, p = 0.05		
Number of soft stools	6.67 *	0.91	1.1	0.41	5.4 *	0.78	16.6, <i>p</i> < 0.001		
Presence of diarrhea	yes		no		yes				
Jumping Frequency	1.89 +	1.31	0	0	41.6 *	2.64	188.1, <i>p</i> < 0.001		
Rearing Frequency	17.8 *	5.94	55.4	6.59	22.2 *	2.46	15.4, <i>p</i> < 0.0001		
Forepaw Licking Frequency	11.2 †	1.80	15.1	3.27	1.7 *	1.00	9.59, <i>p</i> = 0.0008		
Teeth Chattering Frequency	0 +	0	0.4	0.27	9.5 *	1.79	25.0, $p < 0.0001$		

3. Discussion

From the direct *in vivo* phenotypic screening of the mixture-based cyclic peptide library using the 55 °C WWTW assay, we have identified a novel hit CycloAnt, Tyr-[D-Lys-Dap(Ant)-Thr-Gly], exhibiting time- and dose-dependent antinociception with an ED50 value of 0.7 mg/kg (*i.p.*) in mice. Notably, this library was screened earlier using a receptor-binding assay for the MOR, identifying a hit with a binding affinity of 14 nM [31]. However, that cyclic peptide has a sequence of Tyr-[D-Lys-Dap(Ant)-Tyr-Gly], differing from the sequence Tyr-[D-Lys-Dap(Ant)-Thr-Gly] identified in this study. The differing results may stem from the two screening methods utilized, based on different mechanisms and readouts. First, as the *in vivo* direct phenotypic screening tests checked the relative antinociceptive effect of a sample, in terms of the MOR, only agonists can be identified from the assay. While the binding assay at MOR measures the affinity of a sample at the receptor, whether the sample is an agonist or antagonist cannot be differentiated, leaving activity unresolved. Second, the present samples were administered to mice through the intraperitoneal route for the *in vivo* direct phenotypic screening. As the mice are complete organisms with systemic physiology and functions such as metabolism, only these samples with a preferred pharmacokinetic profile will succeed in this screening. It seems likely that the Tyr-[D-Lys-Dap(Ant)-Tyr-Gly] previously identified using the receptor binding assay is unable to adequately activate the MOR under physiological conditions in a complete organism, precluding antinociception in the mouse WWTW assay.

The mixture-based cyclic peptide library is constituted using the imidazole-catalyzed cyclization of linear pentapeptide. While the goal is to cyclize all the linear peptides in a head-to-tail pattern, the composition of the mixture-based positional library is more complicated, as side-to-tail cyclization can take place when Lys or D-Lys is in the linear sequence. Lys and D-Lys are used at positions 2 and 4, and Lys is used at position 5 in the linear samples; therefore, the peptides in the library are not only cyclized in the head-to-tail pattern, but also has the side-to-tail as a side product when Lys or D-Lys is in the sequence.

Since we are tracking the antinociceptive effect and not the purity of the compound in the library, as long as the side reaction is reproducible, the active compounds can be identified through the deconvolution of the mixture library. In this case, CycloAnt, a side-to-tail cyclic peptide, which produced a potent antinociceptive effect, has been successfully identified through the phenotypic screening in mice.

CycloAnt interactions with a broad set of molecular targets were determined with the Safety47 Panel Dose Response, using the SAFETYscan provided by Eurofins DiscoverX. This panel utilizes 78 assays at 47 protein targets, including GPCRs, kinases, nuclear hormone receptors, ion channels, and enzymes. Notably, a number of targets associated with antinociception are included in this panel, including MOR, KOR, and DOR, cyclooxygenase 1 (COX1), cyclooxygenase 2 (COX 2), human serotine receptors 1A (5-HT1A), 1B (5-HT1B), 2A (5-HT2A), and 2B (5-HT2B), human adrenoceptors 3A (5-HT3A), alpha2A (ADRA1), beta1 (ADRB1), and beta2 (ADRB2), voltage gated sodium channel (Nav1.5), and the human glutamate ion channel (NMDAR). CycloAnt only interacted with MOR and DOR in the panel, showing agonist activity at both receptors using the cAMP assay. It did not produce agonist or antagonist activity at any other pain-associated targets in this panel. While the results of MOR testing were confirmed with our *in vivo* testing presently, they diverged with the surprising demonstration in vivo of DOR antagonism. Notably, the DOR agonist activity in the cAMP assay was made with cells expressing a high level of DOR, whereas the *in vivo* results utilizing mice with a physiologically normal level of DOR expression indicated the compound behaved as an antagonist. This discrepancy on the function at DOR may be accounted for receptor reserve [40-42], as a very weak partial agonist/antagonist can appear as a full agonist when receptor reserve is present. While the discrepancy is beyond the scope of this initial identification and characterization, future investigations may prove beneficial in better understanding this interaction. Regardless, consistent with the SAFETYscan results, CycloAnt was not found to produce any significant biological activity at any other targets in this study than DOR and MOR.

The combination of DOR antagonism with MOR agonism has been reported to produce antinociception while possessing safety advantages over typical MOR agonists alone [43]. Although a reduction in antinociceptive tolerance has been shown via a co-administration of MOR agonists with DOR antagonists [44,45] or via a bifunctional MOR agonist/DOR antagonist compounds [46], chronic administration of CycloAnt presently produced a reduction of subsequent antinociception, which was consistent with significant tolerance. However, chronic exposure to CycloAnt did not produce the significant opioid-induced hyperalgesia demonstrated by morphine, and displayed reduced signs of opioid physical dependence, consistent with earlier pharmacological studies [44,45] and reports where *i.c.v.* pretreatment with DOR-selective antisense oligonucleotides decreased morphine dependence in mice [47,48]. It is also consistent with the study in DOR knock out mice in which morphine retained its MOR-mediated analgesic activity without producing tolerance and dependence upon chronic administration [49]. Importantly, CycloAnt did not suppress breathing rate at high doses up to 15 times of the antinociceptive ED₅₀ value, and, in fact, increased breathing rate significantly. CycloAnt treatment also induced significantly less hyperlocomotion than that observed with even lower doses of morphine. All of these results are consistent with the recently reported activity of the bifunctional MOR agonist/DOR antagonist $[Nal(2')^4]CJ-15,208$, which also did not produce hyperlocomotion or respiratory depression; indeed, the bifunctional peptide briefly and significantly increased respiration as well [46]. Given the absence of CycloAnt-mediated activity at off-target proteins, future studies are anticipated to investigate potential mechanisms by which the addition of DOR antagonism may contribute to the improved safety advantages demonstrated by this cyclic peptide.

The potent antinociceptive effect and low liability profile after the *i.p.* administration suggests this cyclic peptide has a preferred pharmacodynamic and pharmacokinetic profile for systemic use as a potentially safer pain modulator. Therefore, this MOR ago-

nist/DOR antagonist is an excellent initial lead for the development of a peptide-based safer opioid analgesic.

4. Materials and Methods

4.1. General Materials

Silica gel, 130–270 mesh, 60 Å, BET surface area $500 \text{ m}^2/\text{g}$, pore volume $0.75 \text{ cm}^3/\text{g}$, was purchased from Sigma-Aldrich Chemical Company Inc., St. Louis, MO, USA. Boc amino acid derivatives, benzotriazole-1-yloxy-trispyrrolidino-phosphonium hexafluorophosphate (PyBOP), N,N-diisopropylethylamine (DIEA), and trifluoroacetic acid (TFA) were purchased from Chem-Impex. HF were purchased from Air Products (San Marcos, CA, USA). Polypropylene mesh (74 micro) was utilized to prepare mesh packets. Phenyltrimethoxysilane and *p*-Chloromethylphenyltrimethoxysilane were purchased from Gelest, Inc. (Tullytown, PA, USA). Mercaptopropyl isobutyl POSS (POSS-SH) was purchased from Hybrid Plastics (Hattiesburg, MS, USA). Pd(PPh_3)_4 and all solvents were purchased from Fisher Scientific, Hampton, NH, USA.

LC-MS (APCI and ESI) was recorded on an Agilent LCMS-1260 (Santa Clara, CA, USA) infinity II at 6120 quadrupole LC/MS mass spectrometer at 214 nm using a Zorbax Eclipse Plus C18 (3.0×100 mm, 3.5μ m) column. Preparative RP-HPLC was performed on a Shimadzu preparative HPLC using a Phenomenex Luna 10 μ C18 100 Å column (21.2×250 mm). NMR spectra were recorded on a Bruker Avance III-HD 400 instrument (Singapore) at 400 MHz for ¹H NMR and 100 MHz for ¹³C NMR. NMR chemical shifts are expressed in ppm relative to internal solvent peak and coupling constant were calculated in hertz.

4.2. Synthesis of the Mixture-Based Cyclic Peptide Library

The cyclic peptide library was synthesized as we have previously reported [31]. Briefly, all linear peptide thioesters were synthesized using the functionalized mercaptomethylphenyl silica gel as 'volatilable' support. Boc-amino acids were coupled on the silica support using PyBOP/DIEA as coupling reagents. X-positions were coupled as mixtures of N^{α} -Boc protected amino acids using concentration ratios to compensate for the relative reaction rates in competitive couplings. Position 3 was coupled with a N^{α} -Boc-N^{β}-Alloc-diaminoproponic acid. After peptide elongation on resin, the Boc-protected peptides were treated with $Pd(PPh_3)_4$ (0.1 equiv.) in the presence of $PhSiH_3$ (20 equiv.) in DCM to remove the Alloc protection group. The resin bound peptides were coupled with 2-nitrobenzoic acid, followed by the treatment of 2 M SnCl₂ in DMF overnight to reduce the nitro group to an amino group to generate the Ant-label. After removal of the Boc group with 55% TFA, the resin-bound Ant-labeled peptides were treated with anhydrous HF in the presence of 5% anisole at 0 °C for 2 h. After removal of the HF with nitrogen stream and lyophilization, the linear peptide thioesters were cyclized in a mixed solution of 1.5 M imidazole (aq.) and acetonitrile (1:7 v/v) at a concentration of 1 mM for 72 h [36], forming the Ant-labeled cyclic peptides.

4.3. Solution-Phase Synthesis of Cyclic Peptides Tyr-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt) and Leu-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt-Leu)

4.3.1. Synthesis of Boc-Dap[Ant(Boc)]-OH

All Boc-amino acids used for the POSS-thioester synthesis were commercially available, except the third amino acid, Boc-Dap[Ant(Boc)], which was synthesized in the lab and directly utilized for the peptide synthesis without further purification. Briefly, Boc-Dap-OH (1 mmol) was dispersed in 2 mL of DMF with the presence of DIEA (1 mmol). In a separate vial, Boc-2-aminobenzoic acid (1 mmol, 0.11 M in DCM) was mixed with PyBOP (1 mmol) and DIEA (1.5 mmol) and activated for 5 min. The solution of Boc-Dap-OH was then added to the activated Boc-2-aminobenzoic acid. The reaction was stirred at r.t. and the completion of the reaction was monitored by LC-MS. After 5 h, the resulting Boc-Dap[Ant(Boc)] was utilized directly for peptide synthesis without purification.

4.3.2. Synthesis of Linear Peptide POSS-Thioesters Boc-Tyr(tBu)-D-Lys(Alloc)-Dap(Ant)-Thr-Gly-S-POSS and Boc-Leu-D-Lys(Alloc)-Dap(Ant)-Thr-Gly-S-POSS

The linear peptide thioester was synthesized using POSS-SH as a soluble support as we previously described [37]. Briefly, Boc-Gly-OH (1.05 mmol) pre-activated with PyBOP (1.05 mmol) and DIEA (2.1 mmol) was added to POSS-SH (0.10 M in 10 mL DCM). The reaction mixture was stirred at room temperature for 3 h. After removal of DCM, Boc-Gly-S-POSS was washed 3 times with 75% acetonitrile containing aqueous HCl (0.05 M). The resulting precipitation was collected by centrifugation and freeze-dried. Boc-Gly-S-POSS was treated with 55% TFA for 30 min to remove the Boc protective group. After removal of TFA/DCM *in vacuo*, the peptide chain was elongated by coupling to the subsequent Boc-amino acid using the method described above. At each of the peptide coupling steps, an extra amount of DIEA was added to the reaction mixture to adjust the pH value to 8. The completion of the amino acid coupling reaction was monitored by a ninhydrin test. The linear peptide POSS-thioesters were obtained at an overall total yield of 81.8%.

4.3.3. Synthesis of Cyclic Peptides Tyr-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt) and Leu-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt-Leu)

The cyclic peptides were synthesized using the imidazole-promoted cyclization methods as we previously reported [38]. Under N2 protection, the Boc-Tyr(tBu)-D-Lys(Alloc)-Dap(Ant)-Thr-Gly-S-POSS (1.0 mmol) or the Boc-Leu-D-Lys(Alloc)-Dap(Ant)-Thr-Gly-S-POSS (1.0 mmol) was dissolved in 100 mL of DCM to a concentration of 10 mM. A portion of PhSiH₃ (24 mmol) and Pd(PPh₃)₄ (0.07 mmol) were then added to facilitate the removal of the Alloc protective group. After reacting overnight, imidazole (1.36 g, 20 mmol) was directly added to the reaction mixture to a concentration of 0.2 M and the reaction was carried out for 24 h. After removal of the solvent *in vacuo*, 75% acetonitrile (aq.) was added to dissolve the yielded cyclic peptide while precipitating the POSS-SH released from peptide cyclization. The supernatant was collected and then lyophilized to dry. The resulting powder was washed with water to remove imidazole. The precipitation was collected and re-dissolved in 65% acetonitrile. After centrifugation at 4000 rpm for 10 min, the supernatant was collected and freeze-dried to yield the protected cyclic peptide Boc-Tyr(tBu)-[D-Lys-Dap(Ant)-Thr-Gly] or Boc-Leu-[D-Lys-Dap(Ant)-Thr-Gly]. Following the treatment with 55% TFA for 30 min, the fully unprotected cyclic peptide Tyr-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt) or Leu-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt) was yielded. The crude compound was purified by reverse-phase HPLC. The cyclic peptide was obtained as a TFA salt after HPLC purification with a purity of 95%.

Tyr-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt). ¹H NMR (400 MHz, DMSO-*d*₆) δ 0.99–1.22 (m, 2H), 1.03 (d, 3H, *J* = 6 Hz), 1.08–1.18 (m, 1H), 1.29–1.39 (m, 3H), 1.49–1.56 (m, 1H), 2.85–2.87 (m, 1H), 2.90 (m, 1H), 2.96–2.98 (m, 1H), 3.08–3.12 (m, 1H), 3.50 (td, 1H, *J* = 14, 6 Hz), 3.88 (d, 1H, *J* = 6.8 Hz), 3.90–3.94 (m, 1H), 3.99–4.02 (m, 2H), 4.08 (t, 1H, *J* = 6 Hz), 4.2–4.24 (m, 1H), 5.07 (d, 1H, *J* = 4.8 Hz), 6.34 (brs, 2H), 6.49–6.52 (dd, 1H, *J* = 7.2, 1 Hz), 6.54 (d, 1H, *J* = 2 Hz), 6.72 (d, 3H, *J* = 8 Hz), 7.03 (d, 2 H, *J* = 8 Hz), 7.16 (td, 1H, *J* = 7.6, 1.6 Hz), 7.33 (t, 1H, *J* = 5.2 Hz), 7.46 (dd, 1H, *J* = 8, 1.2 Hz), 7.64 (d, 1H, *J* = 5.2 Hz), 7.72 (brs, 1H), 8.13 (brs, 2H), 8.29 (t, 1H, *J* = 5.6 Hz), 8.38 (t, 1H, *J* = 5.6 Hz), 8.43 (d, 1H, *J* = 7.2 Hz), 8.47 (d, 1H, *J* = 6.4 Hz), 9.39 (s, 1H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 20.1, 21.2, 28.1, 31.5, 36.8, 38.2, 40.7, 43.8, 53.6, 54.2, 56.4, 60.2, 66.3, 114.9, 115.3, 115.7, 115.8, 116.9, 118.7, 125.2, 128.8, 130.9, 132.5, 149.7, 157.1, 158.4, 158.7, 168.2, 169.2, 170.5(2), 170.6, 172.1. ¹⁹F NMR (376 MHz) δ-37.7 (s). [M + H]⁺ calculated 655.3; found 655.3

Leu-[D-Lys-Dap(Ant)-Thr-Gly] (CycloAnt-Leu). ¹H NMR (400 MHz, DMSO- d_6) δ 0.92 (d, 6H, J = 8 Hz), 1.03 (d, 3H, J = 4 Hz), 1.29–1.32 (m, 2H), 1.38–1.39 (m, 2H), 1.57–1.65 (m, 5H), 2.96–2.99 (m, 1H), 3.12–3.16 (m, 1H), 3.36 (dd, 1H, J = 16, 4 Hz), 3.53 (m, 1H), 3.88 (dd, 1H, J = 16, 8 Hz), 3.93–4.00 (m, 1H), 4.02–4.04 (m, 2H), 4.12 (t, 1H, J = 8 Hz), 4.19 (t, 1H, J = 8 Hz), 6.51 (td, 1H, J = 8, 1 Hz), 6.73 (d, 1H, J = 8 Hz), 7.16 (td, 1 H, J = 8, 1 Hz), 7.41 (t, 1H, J = 8 Hz), 7.48 (dd, 1H, J = 7.6, 1.6 Hz), 7.66 (d, 1H, J = 8 Hz), 8.15 (br.s, 3H), 8.26 (t, 1H, J = 6 Hz), 8.38 (t, 1H, J = 5.6 Hz), 8.57 (d, 1H, J = 8 Hz), 8.83 (d, 1H, J = 6.8 Hz). ¹³C NMR (100 MHz, DMSO- d_6) δ 20.1, 21.7, 22.1, 23.2, 24.1, 28.2, 31.6, 38.2, 40.4, 40.5, 43.8, 51.4, 54.4, 56.4, 60.0, 66.3, 114.7, 115.2,

115.8, 116.9, 118.8, 128.8, 132.5, 149.8, 158.4, 158.8, 169.2, 169.5, 170.5(2), 170.7, 172.4. [M + H]⁺ calculated 605.3; found 605.3.

4.4. Animals

Male C57BL/6J mice (19–27 g each, Jackson Laboratories, Bar Harbor, ME, USA) were used for screening and deconvoluting the library. Male mu-opioid receptor gene knockout mice (MOR KO), kappa-opioid receptor gene-knockout mice (KOR KO), and delta-opioid receptor gene-knockout mice (DOR KO), obtained from a breeding colony established at the University of Florida with progenitors from Jackson Labs, were used for additional testing of the lead. Four mice per cage were housed in a room with controlled temperature and 12 h light/dark cycle controls with the lights on from 07:00 a.m. to 19:00 p.m.; food and water were available ad libitum. All procedures with mice were approved by the local IACUC.

4.5. Assessment of Opioid-Induced Hyperalgesia Using the 48 °C Tail-Withdrawal Assay in Mice

The 48 °C warm water tail-withdrawal assay was performed as previously described [39], with the latency to withdraw the tail taken as the end point. The water temperature of 48 °C was selected for this work to ensure a moderate tail-withdrawal response, with a measurable decrease in withdrawal time possible, but also a significant temperature for hyperalgesic testing. If the mouse failed to display a tail-flick in 30 s, the tail was removed from the water to minimize tissue damage. After determining control latencies on day 1, mice received chronic morphine (10 mg/kg, *i.p.*) or CycloAnt (3 mg/kg, *i.p.*) treatment twice a day for 4 days, as described below. Dosing of these compounds was based on the demonstration of equianalgesic efficacy (see Section 2). Mice were then tested prior to treatment on day 5 for evidence of induced hyperalgesia. Experimentally induced decreases in tail-withdrawal latencies in this assay indicate effects of hyperalgesia [50,51].

4.6. Antinociceptive Test Using the 55 °C Tail-Withdrawal Assay in Mice

The 55 °C warm water tail-withdrawal assay was performed, as previously described, [52] to screen the library [32]. Mixed-based samples were dissolved at a concentration of 5 mg/mL in saline containing 10% DMSO. Each sample was given to mice (n = 8) intraperitoneally (*i.p.*) at a dose of 5 mg/kg. At 0.5, 1, 2, 3.5, 5, 8 h post-administration of the samples, the tail-withdrawal latencies were measured for each sample. The average response time to tail-withdrawal at the six time points were then combined. The combined response time to withdrawal tail was used to determine the antinociceptive effect of each sample. A cutoff of 15 s was used to avoid tissue damage; those mice failing to withdraw their tails within this time were assigned a maximal antinociceptive score of 100%.

To determine antinociceptive activity of individual compounds, the tail-withdrawal latency was determined repeatedly every 10 min, following administration of the compounds for 1 h or until latencies returned to baseline values. At each time point, antinociception was calculated according to the following formula: % antinociception = $100 \times$ (test latency – control latency)/(15 – control latency). Tail-withdrawal data points are the means of 8–16 mice, unless otherwise indicated, with SEM shown by error bars.

The opioid receptor involvement in the agonist activity of the individual compounds was determined by testing in MOR KO, KOR KO and DOR KO mice as utilized previously [53].

To determine antagonist activity, MOR KO mice were pretreated with CycloAnt 20 min prior to the administration of the KOR-selective agonist U50,488 (10 mg/kg, *i.p.*), or DOR-selective agonist SNC-80 (100 nmol, *i.c.v.*); the MOR KO mice do not display antinociceptive activity of CycloAnt. Antinociception produced by the established opioid agonists was then measured 40 min after their administration. Additionally, the duration of DOR antagonism produced by CycloAnt was further determined by evaluating SNC-80-induced antinociception 3, 6 or 8 h after administration of CycloAnt.

4.7. Respiratory Depression and Hyperlocomotion

Male C57Bl/6J mice were tested for respiratory effects using automated, computercontrolled Comprehensive Lab Animal Monitoring System (CLAMS) apparatus. Mice (n = 8) were habituated in closed apparatus cages ($23.5 \times 11.5 \times 13 \text{ cm}^3$) for 30 min before being administrated with the hit i.p. at the experimented high dose of 3.0 mg/kg, and an even higher dose at 15 times of the ED₅₀ value, 10.6 mg/kg, *i.p.* Mice were then returned to chambers, and their respiration rate (breaths/min) was measured for 100 min in 30 s intervals using a pressure transducer built into the enclosed, sealed CLAMS cage. The resultant change in pressure during mouse breathing was recorded to reveal the respiration rate (breaths/min).

Infrared beams located in the floor measured locomotion as the number of beam breaks. Respiration and locomotive data were averaged over 20 min periods for 120 min post-injection of the test compound. Data are presented as % vehicle response \pm SEM, ambulation or breaths per minute.

4.8. Antinociceptive Tolerance and Naloxone-Precipitated Opioid Withdrawal Assay

Mice were randomly assigned to one of three groups and given saline (*i.p.*; n = 10), morphine (10 mg/kg, *i.p.*; n = 10) or CycloAnt (3 mg/kg, *i.p.*; n = 9) twice daily for 4 days, and once more on the morning of the fifth day. Mice were then tested in the 55 °C WWTW assay to assess for the loss of therapeutic efficacy (antinociceptive tolerance; see above). Two hours after this test, naloxone (10 mg/kg, *i.p.*) was administrated to all mice to precipitate opioid withdrawal [39]. Mouse activity was digitally recorded and evaluated later for an established record of withdrawal behaviors for 15 min [39]. Immediately upon injection of naloxone, all mice were scored for number of jumps, forepaw tremor, wet dog shakes, straightening, rearing, forepaw licking, teeth chattering, and the presence (as +/-) and number of soft stools.

4.9. Statistical Analysis

All dose–response lines were analyzed by regression, and ED₅₀ (effective dose producing 50% antinociception) values and 95% C.I. were determined using individual data points from graded dose–response curves with Prism 10.0 software (GraphPad, La Jolla, CA, USA). Percent antinociception was used to determine within-group effects and to allow comparison to baseline latency in tail-withdrawal experiments. The statistical significance of differences between ED₅₀ values was determined by evaluation of the ED₅₀ value shift via nonlinear regression modeling with Prism 10.0 software. Significant differences in behavioral data were analyzed by ANOVA (one-way or two-way with repeated measures (RM), as appropriate). Significant results were further analyzed with Sidak's, Tukey's, or Dunnett's multiple comparison post hoc tests, as appropriate. All data are presented as mean \pm SEM, with significance set at p < 0.05.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/molecules28227548/s1, NMR profiles of CycloAnt and CycloAnt-Leu and Safety47 Panel response.

Author Contributions: Conceptualization, R.A.H.; methodology, Y.L. and J.P.M.; formal analysis, J.P.M. and Y.L.; investigation, Y.L., M.G.-S., S.O.E., A.E. and J.P.M.; writing—original draft preparation, Y.L.; writing—review and editing, J.P.M. and Y.L.; visualization, J.P.M. and Y.L.; supervision, J.P.M. and Y.L.; project administration, J.P.M. and Y.L.; funding acquisition, J.P.M. and Y.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research was partially supported by grants from the National Institute for Health 5P20 GM109091 and by an ASPIRE grant from the Office of the Vice President for Research at the University of South Carolina.

Institutional Review Board Statement: The animal study protocols governing this work were approved prior to testing by the Institutional Animal Care and Use Committees at the Torrey Pines Institute for Molecular Studies and the University of Florida (protocol #201808990, approved 7/17/2018).

Informed Consent Statement: Not applicable.

Data Availability Statement: Data is contained within the article and the Supplementary Material.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

Alloc: allyloxycarbonyl; ANOVA: analysis of variance; Ant, anthraniloyl; Boc, tert-butyloxycarbonyl; CI, confidence interval; CycloAnt, Tyr-[D-Lys-Dap(Ant)-Thr-Gly]; CycloAnt-Leu, Leu-[D-Lys-Dap(Ant)-Thr-Gly]; CLAMS; Comprehensive Lab Animal Monitoring System; DIEA, diisopropylethylamine; DCM, dichloromethane; DMSO, dimethyl sulfoxide; DOR, delta opioid receptor; DOR KO, delta opioid receptor gene-disrupted mice; ED50, dose for 50% maximum effect; HOBt, 1-hydroxybenzotriazole; *i.c.v.*, intracerebroventricular; *i.p.*, intraperitoneal; KOR, kappa opioid receptor; KOR KO, kappa opioid receptor gene-knock-out mice; LC-MS, liquid chromatography-mass spectrometry; MOR, mu opioid receptor; MOR KO, mu opioid receptor gene-disrupted mice; MS, morphine sulfate; NMR, nuclear magnetic resonance; POSS, polyhedral oligomeric silsesquioxane; PyBOP, benzotriazole-1-yloxy-trispyrrolidino-phosphonium hexafluorophosphate; RM, repeated measures; RP-HPLC, reverse-phase high-performance liquid chromatography; SNC-80, (+)-4-[(α R)- α ((2S,5R)-4-allyl-2,5-dimethyl-1-piperazinyl)-3-methoxybenzyl]-*N*,*N*-diethylbenzamide; TFA, trifluoroacetic acid; U50,488,(±)-*trans*-3,4-dichloro-*N*-methyl-*N*-[2-(1-pyrrolidinyl) cyclohexyl]benzeneacetamide.

Amino acids are the L-isomer unless otherwise specified; abbreviations for standard amino acids follow IUPAC-IUB Joint Commission of Biochemical Nomenclature [54].

References

- 1. Alorfi, N.M. Pharmacological methods of pain management: Narrative review of medication used. *Int. J. Gen. Med.* **2023**, *16*, 3247–3256.
- 2. Savage, S.R.; Kirsh, K.L.; Passik, S.D. Challenges in using opioids to treat pain in persons with substance use disorders. *Addict. Sci. Clin. Pract.* **2008**, *4*, 4–25. [CrossRef]
- 3. Schiller, P.W. Opioid peptide-derived analgesics. AAPS J. 2005, 7, E560–E565. [PubMed]
- 4. Webster, L.; Schmidt, W.K. Dilemma of addiction and respiratory depression in the treatment of pain: A prototypical endomorphin as a new approach. *Pain Med.* **2020**, *21*, 992–1004.
- DiMaio, J.; Nguyen, T.M.-D.; Lemieux, C.; Schiller, P.W. Synthesis and Pharmacological Characterization in Vitro of Cyclic Enkephalin Analogues: Effect of Conformational Constraints on Opiate Receptor Selectivity. J. Med. Chem. 1982, 25, 1432–1438. [CrossRef]
- Stoeber, M.; Jullié, D.; Lobingier, B.T.; Laeremans, T.; Steyaert, J.; Schiller, P.W.; Manglik, A.; Von Zastrow, M. A genetically encoded biosensor reveals location bias of opioid drug action. *Neuron* 2018, *98*, 963–976. [PubMed]
- Tsvetanova, N.G.; Von Zastrow, M. Spatial encoding of cyclic AMP signaling specificity by GPCR endocytosis. *Nat. Chem. Biol.* 2014, 10, 1061–1065. [PubMed]
- Jong, Y.-J.I.; Sergin, I.; Purgert, C.A.; O'Malley, K.L. Location-dependent signaling of the group 1 metabotropic glutamate receptor mGlu5. *Mol. Pharmacol.* 2014, 86, 774–785.
- Valentino, R.J.; Volkow, N.D. Untangling the complexity of opioid receptor function. *Neuropsychopharmacology* 2018, 43, 2514–2520. [CrossRef] [PubMed]
- 10. Gomes, I.; Sierra, S.; Lueptow, L.; Gupta, A.; Gouty, S.; Margolis, E.B.; Cox, B.M.; Devi, L.A. Biased signaling by endogenous opioid peptides. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 11820–11828. [PubMed]
- Gallantine, E.L.; Meert, T.F. A Comparison of the antinociceptive and adverse effects of the mu-opioid agonist morphine and the delta-opioid agonist SNC80. *Clin. Pharmacol. Toxicol.* 2005, *97*, 39–51. [CrossRef]
- Zhang, X.; Arvidsson, L.B.; Elde, U.R.; Hökfelt, T. Localization and regulation of the delta-opioid receptor in dorsal root ganglia and spinal cord of the rat and monkey: Evidence for association with the membrane of large dense-core vesicles. *Neuroscience* 1998, *82*, 1225–1242. [CrossRef]
- Mas-Orea, X.; Basso, L.; Blanpied, C.; Gaveriaux-Ruff, C.; Cenac, N.; Dietrich, G. Delta opioid receptors on nociceptive sensory neurons mediate peripheral endogenous analgesia in colitis. *J. Neuroinflamm.* 2022, 19, 7. [CrossRef]
- Conibear, A.E.; Asghar, J.; Hill, R.; Henderson, G.; Borbely, E.; Tekus, V.; Helyes, Z.; Palandri, J.; Bailey, C.; Starke, I.; et al. A novel G protein–biased agonist at the δ opioid receptor with analgesic efficacy in models of chronic pain. *J. Pharmacol. Exp. Ther.* 2020, 372, 224–236. [CrossRef] [PubMed]
- Vekariya, R.H.; Lei, W.; Ray, A.; Saini, S.K.; Zhang, S.; Molnar, G.; Barlow, D.; Karlage, K.L.; Bilsky, E.J.; Houseknecht, K.L.; et al. Synthesis and structure–activity relationships of 5'-aryl-14-alkoxypyridomorphinans: Identification of a μ opioid receptor agonist/δ opioid receptor antagonist ligand with systemic antinociceptive activity and diminished opioid side effects. *J. Med. Chem.* 2020, *63*, 7663–7694. [CrossRef]

- Faouzi, A.; Uprety, R.; Gomes, I.; Massaly, N.; Keresztes, A.I.; Le Rouzic, V.; Gupta, A.; Zhang, T.; Yoon, H.J.; Ansonoff, M.; et al. Synthesis and pharmacology of a novel μ–δ opioid receptor heteromer-selective agonist based on the carfentanyl template. *J. Med. Chem.* 2020, *63*, 13618–13637. [CrossRef]
- Matsumoto, K.; Narita, M.; Muramatsu, N.; Nakayama, T.; Misawa, K.; Kitajima, M.; Tashima, K.; Devi, L.A.; Suzuki, T.; Takayama, H.; et al. Orally active opioid μ/δ dual agonist MGM-16, a derivative of the indole alkaloid mitragynine, exhibits potent antiallodynic effect on neuropathic pain in mice. *J. Pharmacol. Exp. Ther.* **2014**, *348*, 383–392. [CrossRef]
- Podolsky, A.T.; Sandweiss, A.; Hu, J.; Bilsky, E.J.; Cain, J.P.; Kumirov, V.K.; Lee, Y.S.; Hruby, V.J.; Vardanyan, R.S.; Vanderah, T.W. Novel fentanyl-based dual μ/δ-opioid agonists for the treatment of acute and chronic pain. *Life Sci.* 2013, 93, 1010–1016. [CrossRef] [PubMed]
- Lenard, N.R.; Daniels, D.J.; Portoghese, P.S.; Roerig, S.C. Absence of conditioned place preference or reinstatement with bivalent ligands containing mu-opioid receptor agonist and delta-opioid receptor antagonist pharmacophores. *Eur. J. Pharmacol.* 2007, 566, 75–82. [CrossRef] [PubMed]
- Dietis, N.; Niwa, H.; Tose, R.; McDonald, J.; Ruggieri, V.; Filaferro, M.; Vitale, G.; Micheli, L.; Ghelardini, C.; Salvadori, S.; et al. In vitro and in vivo characterization of the bifunctional mu and delta opioid receptor ligand UFP-505. *Br. J. Pharmacol.* 2018, 175, 2881–2896. [CrossRef]
- Stevenson, G.W.; Folk, J.E.; Rice, K.C.; Negus, S.S. Interactions between delta and mu opioid agonists in assays of schedulecontrolled responding, thermal nociception, drug self-administration, and drug versus food choice in rhesus monkeys: Studies with SNC80 [(+)-4-[(alphaR)-alpha-((2S,5R)-4-allyl-2,5-dimethyl-1-piperazinyl)-3-methoxybenzyl]-N,N-diethylbenzamide] and heroin. J. Pharmacol. Exp. Ther. 2005, 314, 221–231.
- Lowery, J.J.; Raymond, T.J.; Giuvelis, D.; Bidlack, J.M.; Polt, R.; Bilsky, E.J. In vivo characterization of MMP-2200, a mixed δ/μ opioid agonist, in mice. J. Pharmacol. Exp. Ther. 2011, 336, 767–778. [CrossRef]
- Stevenson, G.W.; Luginbuhl, A.; Dunbar, C.; Lavigne, J.; Dutra, J.; Atherton, P.; Bell, B.; Cone, K.; Giuvelis, D.; Polt, R.; et al. The mixed-action delta/mu opioid agonist MMP-2200 does not produce conditioned place preference but does maintain drug self-administration in rats, and induces in vitro markers of tolerance and dependence. *Pharmacol. Biochem. Behav.* 2015, 132, 49–55. [CrossRef]
- 24. Lei, W.; Vekariya, R.H.; Ananthan, S.; Streicher, J.M. A novel mu-delta opioid agonist demonstrates enhanced efficacy with reduced tolerance and dependence in mouse neuropathic pain models. *J. Pain* **2020**, *21*, 146–160. [CrossRef]
- Vicario, N.; Pasquinucci, L.; Spitale, F.M.; Chiechio, S.; Turnaturi, R.; Caraci, F.; Tibullo, D.; Avola, R.; Gulino, R.; Parenti, R.; et al. Simultaneous activation of mu and delta opioid receptors reduces allodynia and astrocytic connexin 43 in an animal model of neuropathic pain. *Mol. Neurobiol.* 2019, 56, 7338–7354. [CrossRef]
- Anand, J.P.; Kochan, K.E.; Nastase, A.F.; Montgomery, D.; Griggs, N.W.; Traynor, J.R.; Mosberg, H.I.; Jutkiewicz, E.M. In vivo effects of μ-opioid receptor agonist/δ-opioid receptor antagonist peptidomimetics following acute and repeated administration. Br. J. Pharmacol. 2018, 175, 2013–2027. [CrossRef]
- Nastase, A.F.; Anand, J.P.; Bender, A.M.; Montgomery, D.; Griggs, N.W.; Fernandez, T.J.; Jutkiewicz, E.M.; Traynor, J.R.; Mosberg, H.I. Dual pharmacophores explored via structure-activity relationship (SAR) matrix: Insights into potent, bifunctional opioid ligand design. *J. Med. Chem.* 2019, 62, 4193–4203. [CrossRef]
- Schiller, P.W.; Fundytus, M.E.; Merovitz, L.; Weltrowska, G.; Nguyen, T.M.-D.; Lemieux, C.; Chung, N.N.; Coderre, T.J. The opioid μ agonist/δ antagonist DIPP-NH2[Ψ] produces a potent analgesic effect, no physical dependence, and less tolerance than morphine in rats. *J. Med. Chem.* 1999, 42, 7. [CrossRef]
- Ananthan, S. Opioid ligands with mixed μ/δ opioid receptor interactions: An emerging approach to novel analgesics. *AAPS J.* 2006, *8*, E118–E125. [CrossRef]
- Ananthan, S.; Saini, S.K.; Dersch, C.M.; Xu, H.; McGlinchey, N.; Giuvelis, D.; Bilsky, E.J.; Rothman, R.B. 14-Alkoxy- and 14acyloxypyridomorphinans: Mu agonist/delta antagonist opioid analgesics with diminished tolerance and dependence side effects. J. Med. Chem. 2012, 55, 8350–8363. [CrossRef]
- 31. Li, Y.; Dooley, C.T.; Misler, J.A.; Debevec, G.; Giulianotti, M.A.; Cazares, M.E.; Maida, L.; Houghten, R.A. Fluorescent mu selective opioid ligands from a mixture based cyclic peptide library. *ACS Comb. Sci.* **2012**, *14*, 673–679. [CrossRef] [PubMed]
- Reilley, K.J.; Giulianotti, M.; Dooley, C.T.; Nefzi, A.; McLaughlin, J.P.; Houghten, R.A. Identification of two novel, potent, low-liability antinociceptive compounds from the direct in vivo screening of a large mixture-based combinatorial library. *AAPS J.* 2010, 12, 318–329. [CrossRef]
- Armishaw, C.J.; Banerjee, J.; Ganno, M.L.; Reilley, K.J.; Eans, S.O.; Mizrachi, E.; Gyanda, R.; Hoot, M.R.; Houghten, R.A.; McLaughlin, J.P. Discovery of novel antinociceptive α-conotoxin analogues from the direct in vivo screening of a synthetic mixture-based combinatorial library. ACS Comb. Sci. 2013, 15, 153–161. [CrossRef]
- 34. Houghten, R.A.; Ganno, M.L.; McLaughlin, J.P.; Dooley, C.T.; Eans, S.O.; Santos, R.G.; Lavoi, T.; Nefzi, A.; Welmaker, G.; Giulianotti, M.A.; et al. Direct phenotypic screening in mice: Identification of individual, novel antinociceptive compounds from a library of 734 821 pyrrolidine bis-piperazines. ACS Comb. Sci. 2016, 18, 51–64. [CrossRef] [PubMed]
- Li, Y.; Yu, Y.; Giulianotti, M.; Houghten, R.A. High throughput synthesis of peptide α-thioesters through the use of "volatilizable" support. J. Comb. Chem. 2008, 10, 4. [CrossRef] [PubMed]

- Li, Y.; Yongye, A.; Giulianotti, M.; Martinez-Mayorga, K.; Yu, Y.; Houghten, R.A. Synthesis of cyclic peptides through direct aminolysis of peptide thioesters catalyzed by imidazole in aqueous organic solutions. *J. Comb. Chem.* 2009, 11, 1066–1072. [CrossRef]
- Perry, K.; Sui, B.; Li, Y. Mercapto-functionalized polyhedral oligomeric silsesquioxane as a soluble support for the synthesis of peptide thioesters. *Tetrahedron Lett.* 2021, 85, 153483. [CrossRef]
- Li, Y.; Giulionatti, M.; Houghten, R.A. Macrolactonization of peptide thioesters catalyzed by imidazole and its application in the synthesis of Kahalalide B and analogues. Org. Lett. 2010, 12, 2250–2253. [CrossRef]
- Wilson, L.L.; Eans, S.O.; Cirino, T.J.; Stacy, H.M.; Simons, C.A.; Uprety, R.; Majumdar, S.; McLaughlin, J.P. Kratom alkaloids, natural and semi-synthetic, show less physical dependence and ameliorate opioid withdrawal. *Cell Mol. Neurobiol.* 2021, 41, 1131–1143. [CrossRef] [PubMed]
- 40. Wacker, D.; Stevens, R.C.; Roth, B.L. How Ligands Illuminate GPCR Molecular Pharmacology. Cell 2017, 170, 414–427. [CrossRef]
- Singleton, S.; Bastista-Hon, D.T.; Edelsten, E.; McCaughey, K.S.; Camplisson, E.; Hales, T.G. TRV130 partial agonism and capacity to induce anti-nociceptive tolerance revealed through reducing available μ-opioid receptor number. *Br. J. Pharmacol.* 2021, 178, 1855–1868. [CrossRef]
- 42. Kelly, E. Efficacy and ligand bias at the mu-opioid receptor. Br. J. Pharmacol. 2013, 169, 1430–1446. [CrossRef]
- 43. Schiller, P.W. Bi- or multifunctional opioid peptide drugs. Life Sci. 2010, 86, 598–603. [CrossRef]
- 44. Abdelhamid, E.E.; Sultana, M.; Portoghese, P.S.; Takemori, A.E. Selective blockage of delta-opioid receptors prevents the development of morphine-tolerance and dependence in mice. *J. Pharmacol. Exp. Ther.* **1991**, *258*, 299–303.
- 45. Fundytus, M.E.; Schiller, P.W.; Shapiro, M.; Weltrowska, G.; Coderre, T.J. Attenuation of Morphine Tolerance and Dependence with the Highly Selective Delta-Opioid Receptor Antagonist TIPPY. *Eur. J. Pharmacol.* **1995**, *286*, 105–108. [CrossRef]
- 46. Scherrer, K.H.; Eans, S.O.; Medina, J.M.; Senadheera, S.N.; Khaliq, T.; Murray, T.F.; McLaughlin, J.P.; Aldrich, J.V. Tryptophan substitution in CJ-15,208 (cyclo[Phe-D-Pro-Phe-Trp]) introduces δ-opioid receptor antagonism, preventing antinociceptive tolerance and stress-induced reinstatement of extinguished cocaine-conditioned place preference. *Pharmaceuticals* 2023, *16*, 1218–1238. [CrossRef]
- 47. Sanchez-Blazquez, P.; Garcia-Espana, A.; Garzon, J. Antisense oligodeoxynucleotides to opioid mu and delta receptors reduced morphine dependence in mice: Role of delta-2 opioid receptors. *J. Pharmacol. Exp. Ther.* **1997**, *280*, 1423–1431.
- 48. Kest, B.; Lee, C.E.; McLemore, G.L.; Inturrisi, C.E. An Antisense Oligodeoxynucleotide to the Delta Opioid Receptor (DOR-1) Inhibits Morphine Tolerance and Acute Dependence in Mice. *Brain Res. Bull.* **1996**, *39*, 185–188. [CrossRef]
- Zhu, Y.; King, M.A.; Schuller, A.G.P.; Nitsche, J.F.; Reidl, M.; Elde, R.P.; Unterwald, E.; Pasternak, G.W.; Pintar, J.E. Retention of Supraspinal Delta-like Analgesia and Loss of Morphine Tolerance in δ Opioid Receptor Knockout Mice. *Neuron* 1999, 24, 243–252. [CrossRef]
- 50. Crain, S.M.; Shen, K.F. Acute thermal hyperalgesia elicited by low-dose morphine in normal mice is blocked by ultra-low-dose naltrexone, unmasking potent opioid analgesia. *Brain Res.* 2001, *888*, 75–82. [CrossRef]
- Journigan, V.B.; Mésangeau, C.; Vyas, N.; Eans, S.O.; Cutler, S.J.; McLaughlin, J.P.; Mollereau, C.; McCurdy, C.R. Nonpeptide small molecule agonist and antagonist original leads for neuropeptide FF1 and FF2 receptors. *J. Med. Chem.* 2014, 57, 8903–9827. [CrossRef]
- McLaughlin, J.P.; Hill, K.P.; Jiang, Q.; Sebastian, A.; Archer, S.; Bidlack, J.M. Nitrocinnamoyl and chlorocinnamoyl derivatives of dihydrocodeinone: In vivo and in vitro characterization of μ-selective agonist and antagonist activity. *J. Pharmacol. Exp. Ther.* 1999, 289, 304–311. [PubMed]
- Brice-Tutt, A.C.; Wilson, L.L.; Eans, S.O.; Stacy, H.M.; Simons, C.A.; Simpson, G.G.; Coleman, J.S.; Ferracane, M.J.; Aldrich, J.V.; McLaughlin, J.P. Multifunctional opioid receptor agonism and antagonism by a novel macrocyclic tetrapeptide prevents reinstatement of morphine-seeking behaviour. *Br. J. Pharmacol.* 2020, *177*, 4209–4222. [CrossRef] [PubMed]
- 54. Nomenclature Committee of IUB (NC-IUB) and IUPAC-IUB Joint Commission on Biochemical Nomenclature (JCBN). *Eur. J. Biochem.* **1984**, *138*, 9–37.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.