

Article

Phenolic Profile, Antioxidant Activity, and Chemometric Classification of Carob Pulp and Products

Georgia D. Ioannou ¹, Ioanna K. Savva ¹, Atalanti Christou ¹, Ioannis J. Stavrou ²
and Constantina P. Kapnissi-Christodoulou ^{1,*}

¹ Department of Chemistry, University of Cyprus, 1678 Nicosia, Cyprus

² Department of Life Sciences, European University Cyprus, 2404 Nicosia, Cyprus

* Correspondence: ckapni1@ucy.ac.cy

Abstract: In recent years, carob and its derived products have gained wide attention due to their health-promoting effects, which are mainly attributed to their phenolic compounds. Carob samples (carob pulps, powders, and syrups) were analyzed to investigate their phenolic profile using high-performance liquid chromatography (HPLC), with gallic acid and rutin being the most abundant compounds. Moreover, the antioxidant capacity and total phenolic content of the samples were estimated through DPPH (IC₅₀ 98.83–488.47 mg extract/mL), FRAP (48.58–144.32 µmol TE/g product), and Folin–Ciocalteu (7.20–23.18 mg GAE/g product) spectrophotometric assays. The effect of thermal treatment and geographical origin of carobs and carob-derived products on their phenolic composition was assessed. Both factors significantly affect the concentrations of secondary metabolites and, therefore, samples' antioxidant activity (p -value < 10^{−7}). The obtained results (antioxidant activity and phenolic profile) were evaluated via chemometrics, through a preliminary principal component analysis (PCA) and orthogonal partial least square-discriminant analysis (OPLS-DA). The OPLS-DA model performed satisfactorily, differentiating all samples according to their matrix. Our results indicate that polyphenols and antioxidant capacity can be chemical markers for the classification of carob and its derived products.

Keywords: carobs; carob-derived products; polyphenols; antioxidant activity; HPLC; chemometrics; PCA; OPLS-DA

Citation: Ioannou, G.D.; Savva, I.K.; Christou, A.; Stavrou, I.J.; Kapnissi-Christodoulou, C.P. Phenolic Profile, Antioxidant Activity, and Chemometric Classification of Carob Pulp and Products. *Molecules* **2023**, *28*, 2269. <https://doi.org/10.3390/molecules28052269>

Academic Editors:
Dimitrios Kouretas
and Zoi Skaperda

Received: 2 February 2023
Revised: 16 February 2023
Accepted: 17 February 2023
Published: 28 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The carob tree (*Ceratonia siliqua* L.), which belongs to the Leguminosae family, is a flowering evergreen shrub widely cultivated in the Mediterranean region and the Middle East. The carob or carob pod is the edible fruit produced from the carob tree and has recently attracted great attention due to its unique chemical composition and health-promoting properties. The carob pod consists of two main parts, the pulp and the seeds in a ratio of about 90:10 (w/w) [1,2]. Even though carob seeds represent only a small part of the total weight of the fruit, they are of great industrial interest, as they are used to produce locust bean gum (LBG, E410), a natural food thickening and stabilizing agent [1,3]. The pulp remaining after the industrial exploitation of the seeds, also referred to as carob kibble, is considered an agri-food waste [4,5]. Carob pulp has, so far, been used as livestock feed, while recently the worldwide market has turned its attention to carob pulp as it is considered a promising functional and nutraceutical food component [3,6,7]. Currently, carob pulp is used as a raw material for the preparation of several organic products, such as roasted and unroasted carob powders (a cocoa substitute, free of caffeine and theobromine) and carob syrups or molasses [6].

Carob has recently attracted great attention due to its health-promoting effects and potential use for value-added food production. It is considered an abundant source of

several beneficial substances, including dietary fibers, carbohydrates, minerals, and polyphenolic compounds [8]. The majority of the biological effects and health benefits of carob have been associated with the presence of polyphenols [9]. Phenolic compounds are a widespread class of compounds, produced by the secondary metabolism of plants and present in all species of plant origin. Polyphenols are divided into the following subcategories: phenolic acids, flavonoids, tannins, and stilbenes, which differ in terms of the number of phenol rings they contain and the type of structural unit that connects the aromatic rings [10,11]. In the case of carob pods, the main subcategories of polyphenols are the phenolic acids (gallic acid, sinapic acid, and ferulic acid), the flavonoids (flavanols: catechin, epicatechin; flavonols: myricetin, quercetin; flavones: apigenin), and tannins (digalloyl-glucose, trigalloyl-glucose, and tannic acid). Polyphenols, as natural antioxidants, can protect cellular components from oxidative damage and, as a result, reduce the risk of a variety of degenerative diseases related to oxidative stress [12]. In particular, a reduction in cholesterol along with beneficial effects on the lipid profile of human blood have been reported in human trials after the consumption of insoluble carob dietary fiber rich in polyphenols [13]. Furthermore, the anti-cancer, anti-diabetic, and neuroprotective effects of carob are mainly attributed to its polyphenolic composition [2,14,15].

Several analytical methods have been proposed for the qualitative and quantitative determination of polyphenolic compounds in carob and its derived products. Among them, the spectrophotometric methods represent the easiest way to estimate the total phenolic content as well as the antioxidant activity of the samples. In particular, 2,2-diphenyl-1-picrylhydrazyl (DPPH), Ferric Reducing Antioxidant Power (FRAP), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid (ABTS), and Oxygen Radical Absorbance Capacity (ORAC) assays have been previously reported in the literature to determine the antioxidant activity of carob samples [9,14,16–19]. Since antioxidant activity is directly related to phenolic content, the determination of antioxidant activity is usually followed by the determination of total phenolics, with the Folin–Ciocalteu assay being the most commonly used spectrophotometric method for the estimation of the total phenolic content in carob and its derived products [8,14,20–22]. The choice of two or more spectrophotometric assays is necessary for the correct evaluation of the antioxidant activity, since each test is based on a different mechanism of action and the results for the radical scavenging ability of the samples may significantly vary. The use of chromatographic and electrophoretic methods is essential for the identification and quantification of individual carob phenolic compounds. Gas chromatography (GC), capillary electrophoresis (CE), and high-performance liquid chromatography (HPLC) are the most common separation techniques applied to carob analysis [14,15,23–26]. However, due to the low volatility of phenolic compounds, GC is not preferred [10]. The most widely used technique for the identification and quantification of phenolics in carob is HPLC, coupled with either an Ultraviolet-Visible (UV) detector or mass spectrometry (MS).

Chemometrics is a useful and sophisticated tool for classifying and distinguishing samples based on certain common characteristics. Through powerful statistical modeling tools such as principal component analysis (PCA) and orthogonal partial least square-discriminant analysis (OPLS-DA) can be provide insights into classification of experimental groups based on high-dimensional measurements. Nutritional composition, volatile organic compounds, and IR spectra data have, so far, been used as indicators for the discrimination of carob-derived products as well as carob's geographical origin [7,27–30]. Farag et al. first reported the use of secondary metabolites, as chemical markers, for the classification of carobs [31]. Then, Christou et al. achieved the chemometric classification of carob samples based on their geographical origin, using polyphenols as indicators for the chemometric analysis [6]. To the best of our knowledge, no work has, so far, been performed to classify carobs and their derived products according to their type (matrix) using chromatographic and spectrophotometric data for antioxidant activity (DPPH and FRAP) and phenolic composition as chemical markers.

Under this framework, 30 samples of carob and carob-derived products from Mediterranean countries were examined for their TPC and antioxidant capacity through DPPH and FRAP assays. In addition, the phenolic profile of the samples was investigated via HPLC analysis. The effect of the geographical origin and processing method on carob phenolic composition was also investigated. The results from TPC, DPPH, FRAP, and HPLC were used as variables in chemometrics analysis for the classification of samples based on their matrix.

2. Results and Discussion

2.1. Spectrophotometric Analysis

2.1.1. Total Phenolic Content

Since polyphenolic compounds occur in carob in significant amounts and are responsible for its high antioxidant activity, the total phenolic content of carob and derived products (carob powders and syrups) was investigated. The total phenolic content (TPC) of the samples, determined using the Folin–Ciocalteu assay, is presented in Table S1 (Supplementary Materials). TPC values ranged from 7.20 to 23.18 mg gallic acid equivalents (GAE) per gram of sample. Significant differences in TPC values were observed for each type of product studied. Specifically, the TPC ranged from 7.20–12.28 mg GAE/g, 18.19–23.18 mg GAE/g, and 9.51–14.80 mg GAE/g of sample for carob pulps, powders, and syrups, respectively, with a p -value < 0.001 . The carob powder category presented the highest TPC, with the maximum value recorded from the roasted powder originating in Italy. It is important to highlight that roasted carob pulps and powders demonstrated higher TPC values than unroasted ones, while in the case of thermally treated (>40 °C) syrups, they presented lower TPC than non-thermally treated samples (<40 °C). The increased TPC values in roasted pulps and powders may be attributed to the breakdown of the cellular structure of plant material and the release of previously bound and insoluble phenolics during the roasting process. [4,6,32,33]. In the case of syrups, a decrease in TPC was observed during their thermal treatment, possibly due to the thermal degradation of polyphenols [6].

Data on total phenolics of carobs and their derived products vary significantly in the literature and depend on several factors such as the extraction process, geographical origin, degree of maturation, and processing method [8,20]. The TPC values obtained in the present work for carob pulp are higher than those reported by Goulas and Georgiou (2020) (351.4 ± 4.1 mg GAE/100 g carob fruit) and Gregoriou et al. (2021) (11.57 mg GAE/g extract) [14,22]. Roseiro and Duarte et al. (2013) reported a significantly higher TPC value for carob pulp using a supercritical fluid extraction (SFE) (27.1 ± 0.8 mg GAE/g dry mass) and a slightly lower value using UAE (20.4 ± 1.8 mg GAE/g dry mass) [20]. Additionally, Christou et al. (2021) (14.24 mg GAE/g carob pulp) and Kyriacou et al. (2021) (17.4 mg GAE/g dried weight) reported higher TPC values in mature carobs than those obtained in this work [8,34]. Nonetheless, the present findings on TPC are in very good agreement with several previous investigations [21,26].

Quantitative comparison of results for the TPC of carob-derived products with the literature data is even more difficult since there are not enough studies available. Furthermore, as previously mentioned, the variables affecting the obtained TPC have a decisive role and complicate the comparisons. The present findings for carob powder were found to be greater than those recorded by Petkova et al. (2017) (8.11 mg GAE/g dried weight) and in the same range as those reported by Papagianopoulos et al. (2004) (14.55–24.52 g/kg) for roasted powders [19,26]. In the case of carob syrup, our findings for the TPC appeared to be higher than those reported by Papagianopoulos et al. (2004) (3.94 g/kg) [26].

2.1.2. Antioxidant Activity

It is well known that antioxidant activity cannot be properly assessed with a single assay [35]. Therefore, two commonly used assays with different operating principles and mechanisms were included in the present study. Specifically, the antioxidant activity of carobs and their derived products was evaluated in terms of (i) free radical scavenging activity, with the DPPH assay, and (ii) ferric-reducing antioxidant power, with the FRAP assay. The DPPH assay outcomes were expressed as IC_{50} , which is the extract concentration (mg/mL) required to scavenge 50% of DPPH free radical. In general, a lower IC_{50} value indicates the greater antioxidant capacity of the sample, while a higher IC_{50} value is indicative of the sample's lower antioxidant activity. As far as the FRAP assay results are concerned, they were expressed as μmol Trolox equivalents (TE)/g of the sample.

The results of DPPH and FRAP assays are demonstrated in Figure 1, with IC_{50} values ranging from 98.83 to 488.47 mg extract/mL and FRAP values from 46.36 to 144.32 μmol TE/g sample. For both assays, carob powder is the category with the highest antioxidant capacity. The sample from Italy demonstrated the greatest antioxidant activity, which is in agreement with the TPC results (Figure 1). According to the DPPH assay, the unroasted carob pulp from Paphos (Cyprus) presented the lowest antioxidant potential, while, according to the FRAP test, the thermally treated syrup from Crete (Greece) was the one with the lowest antioxidant capacity. It should be emphasized that the antioxidant activity evaluated using different antioxidant assays may vary, due to different mechanisms and operating principles of the methods [36,37]. As observed, thermal treatment significantly affects the phenolic content and antioxidant activity of the samples. In particular, non-thermal treated syrups (DPPH IC_{50} = 232.21 \pm 19.58, FRAP 82.23 \pm 15.81 μM TE/g product), roasted carob pulps (DPPH IC_{50} = 361.00 \pm 11.92, FRAP 89.22 \pm 4.71 μM TE/g product), and powders (DPPH IC_{50} = 98.83 \pm 3.20, FRAP 144.32 \pm 7.16 μM TE/g product) demonstrated higher antioxidant activity than thermal-treated syrups (DPPH IC_{50} = 388.05 \pm 1.75, FRAP 46.36 \pm 6.48 μM TE/g product) and unroasted products (DPPH IC_{50} = 488.47 \pm 34.67, FRAP 73.94 \pm 3.84 μM TE/g product).




Figure 1. Graphical representation of antioxidant activity and total phenolic content of carob pulps and carob-derived products. (#1: unroasted carob pulp, Cyprus; #2: unroasted carob pulp, Greece; #3: roasted carob pulp, Greece; #4: roasted powder, Italy, #5: roasted powder, Cyprus; #6: roasted powder, Greece; #7: unroasted powder, Greece; #8: thermally treated syrup, Cyprus; #9: thermally treated syrup, Greece; #10: non-thermally treated syrup, Greece). Different lower-case letters indicate significant differences ($p < 0.05$) according to Duncan's multiple range test.

Comparing the DPPH and FRAP results obtained in the present study with the literature data was a difficult task. As previously mentioned, many different parameters can

affect the antioxidant capacity of the samples. For comparison purposes, the antioxidant capacity results should be expressed in terms of the same unit of measurement. DPPH values can also be expressed as % inhibition, ascorbic acid equivalents (AAE), or Trolox equivalents (TE), while FRAP values can also be described as FeSO₄ or ascorbic acid equivalents [14,20–22,38,39]. The samples studied in the present work had lower antioxidant potential, in terms of DPPH radical scavenging activity, than the findings of Goulas and Georgiou (2020) (IC₅₀ 1.4–2.6 mg/mL) and Papagianopoulos et al. (2004) (IC₅₀ kibbles: 33.26 g/L, syrup: 26.33 g/L, roasted flours: 7.04–9.96 g/L) [22,26]. In the case of the results obtained from the FRAP assay, comparison with the other literature studies was considered impossible as the units of measurement of antioxidant capacity were different.

2.2. Chromatographic Analysis

2.2.1. Method Validation

Calibration curve parameters, including linear ranges, regression equations, and coefficients of determination are listed in Table 1. In the concentration ranges examined, all analytes demonstrated good linearity, with the corresponding coefficients of determination being higher than 0.997. The limits of detection (LODs) and quantification (LOQs) varied from 0.1 to 0.9 µg/mL and 0.3 to 2.8 µg/mL, respectively. The precision of the method, in terms of repeatability (intraday) and reproducibility (interday), was evaluated by computing the relative standard deviations (RSDs) of the peak areas. As demonstrated in Table 1, the RSD values for intraday precision ranged from 0.32 to 1.04% and for interday precision from 0.92 to 2.68%.

Table 1. LODs and LOQs, regression equations, coefficients of determination, retention times, repeatability and reproducibility.

Compounds	Rt ± SD	Linear Range	Regression Equation	R ²	LOD (µg/mL)	LOQ (µg/mL)	Precision	
							Intraday	Interday
Gallic Acid	2.52 ± 0.00	0.05–0.0001	y = 99234156x – 35103	0.999	0.7	2.1	1.00	0.92
Catechin	3.34 ± 0.02	0.05–0.0001	y = 13269933x – 2012	0.999	0.3	0.9	0.95	2.53
Epicatechin	3.83 ± 0.03	0.01–0.0001	y = 16600346x + 2404	0.998	0.3	0.8	1.04	1.47
Caffeic Acid	4.54 ± 0.05	0.01–0.0001	y = 55118151x + 8713	0.997	0.3	1.0	0.47	1.96
Rutin	5.54 ± 0.07	0.05–0.0001	y = 23176073x – 12698	0.998	0.9	2.8	0.55	1.80
Sinapic Acid	7.99 ± 0.06	0.01–0.0001	y = 39382578x + 2882	0.997	0.4	1.1	0.33	1.92
Quercitrin	11.49 ± 0.08	0.01–0.0001	y = 12295377x – 1169	0.999	0.1	0.3	0.42	1.84
Myricetin	13.12 ± 0.03	0.01–0.0001	y = 25427092x + 2679	0.998	0.3	0.9	0.90	2.68
Quercetin	14.67 ± 0.04	0.01–0.0001	y = 34063703x + 42715	0.998	0.3	0.9	0.32	1.67

2.2.2. Polyphenolic Profile via HPLC Analysis


HPLC analysis was deemed essential for the identification and quantification of the main phenolic compounds in the carob pulp, powder, and syrup extracts. Furthermore, the chromatographic determination in correlation with the spectrophotometric data will assist to identify the individual compounds, which are responsible for the antioxidant activity of the extracts.

Table 2 demonstrates the qualitative and quantitative results obtained from the analysis of carob and derived products. Rutin, gallic acid, and catechin were the most abundant phenolic compounds in the extracts, as demonstrated in Figure 2. As observed, sample processing significantly affects the amount of extracted polyphenolic compounds. Roasted pulp and carob powders have the highest polyphenolic composition, while thermally treated syrups have the lowest amounts of polyphenols.

Table 2. Phenolic composition of carob pulp and carob-derived products.

Compounds	Concentration (mg/g Product)									
	Carob Pulp				Carob Powder			Carob Syrup		
	Unroasted		Roasted		Roasted		Unroasted	>40 °C		<40 °C
	Cyprus	Greece	Italy	Cyprus	Greece	Cyprus	Greece	Cyprus	Greece	
Gallic Acid	0.31 ± 0.01 _e	0.23 ± 0.01 _e	0.40 ± 0.02 _d	0.56 ± 0.07 _c	0.65 ± 0.09 _c	0.61 ± 0.05 _c	0.60 ± 0.05 _c	0.46 ± 0.01 _d	0.99 ± 0.03 _b	1.34 ± 0.03 _a
Catechin	0.47 ± 0.04 _b	0.33 ± 0.01 _{cd}	0.66 ± 0.07 _a	0.24 ± 0.03 _d	0.49 ± 0.05 _b	0.45 ± 0.02 _{bc}	0.55 ± 0.08 _{ab}	0.43 ± 0.04 _{bc}	0.04 ± 0.00 _e	0.06 ± 0.00 _e
Epicatechin	0.11 ± 0.02 _c	0.05 ± 0.00 _{de}	0.14 ± 0.05 _{bc}	0.02 ± 0.00 _{def}	0.17 ± 0.01 _a	0.16 ± 0.01 _{ab}	0.05 ± 0.01 _d	ND _f	0.03 ± 0.00 _{def}	0.02 ± 0.00 _{ef}
Caffeic Acid	ND _b	NQ _b	ND _b	ND _b	NQ _b	NQ _b	NQ _b	ND _b	0.03 ± 0.01 _a	0.03 ± 0.00 _a
Rutin	0.89 ± 0.06 _a	0.50 ± 0.03 _c	0.80 ± 0.09 _a	0.21 ± 0.03 _d	0.82 ± 0.16 _a	0.81 ± 0.08 _a	0.66 ± 0.08 _b	ND _e	NQ _e	NQ _e
Sinapic Acid	ND _d	ND _d	NQ _d	NQ _d	0.02 ± 0.00 _b	0.02 ± 0.00 _c	0.02 ± 0.00 _a	ND _d	NQ _d	ND _d
Quercitrin	0.03 ± 0.01 _c	0.04 ± 0.00 _c	0.34 ± 0.02 _a	ND _d	0.05 ± 0.01 _c	0.04 ± 0.00 _c	0.12 ± 0.02 _b	ND _d	ND _d	ND _d
Myricetin	0.03 ± 0.00 _{cd}	0.03 ± 0.00 _{cd}	0.05 ± 0.01 _a	ND _e	0.05 ± 0.01 _{ab}	0.04 ± 0.01 _{bc}	0.02 ± 0.00 _d	ND _e	ND _e	ND _e
Quercetin	0.02 ± 0.00 _c	0.02 ± 0.00 _b	NQ _d	0.03 ± 0.00 _a	NQ _d	NQ _d	ND _d	ND _d	ND _d	ND _d
Total	1.86	1.20	2.40	1.06	2.25	2.13	2.03	0.79	1.09	1.44

Different lower-case letters within the same row indicate significant differences ($p < 0.05$) according to Duncan's multiple range test. ND: non-detected; NQ: non-quantified.

**Figure 2.** HPLC chromatogram of carob powder. (1) Gallic acid, (2) catechin, (3) epicatechin, (4) caffeic acid, (5) rutin, (6) sinapic acid, (7) quercitrin, (8) myricetin, and (9) quercetin.

Rutin appeared to occur in greater concentrations in carob pulp, followed by catechin and gallic acid. Minor amounts of epicatechin, quercitrin, myricetin, and quercetin were detected in carob pulp, while caffeic and sinapic acid were absent. The roasting of carob pulp had a positive impact on their polyphenolic composition. This is in agreement with the findings of Christou et al., who also reported a greater amount of phenolics in roasted carob pulps than in raw pods [6]. During the roasting of the carob pulp, the concentrations

of all polyphenols increased, except for quercetin, which decreased to non-measurable levels, possibly due to degradation as a result of thermal treatment [40]. It is worth to mention here that quercitrin demonstrated a significant increase during roasting (about 10 times higher concentration in roasted carob pulp). The increased concentrations of polyphenols detected in roasted carob pulp can be attributed to the breakdown of the cellular structure of the plant matrix during the roasting process, which results in the release of previously bound phenolics. The degradation of hydrolyzed tannins and the release of gallic acid units are other possible explanations for the increased polyphenolic content of roasted carob pulp. In regard to the effect of geographical origin on carob's polyphenolic composition, even though Cyprus and Greece (Crete) have similar climatic conditions, the carob pulp from Cyprus demonstrated higher concentrations of gallic acid, catechin, epicatechin, and rutin. In contrast, quercitrin, myricetin, and quercetin were detected at similar concentration levels in samples from both countries.

The phenolic profile of the carob powder was similar to that of the carob pulp, the initial raw material, except for sinapic acid, which was detected and quantified in both roasted and unroasted carob powders. The higher phenolic composition of the unroasted powder compared to the starting material (carob pulp) may be attributed to the technological processes used to produce the carob flour. During the processing steps, bound phenolic compounds are released due to the rupture of the cellular structure of the initial raw material.

In the case of the carob pulp, the roasted powder appeared to have a higher phenolic content than the unroasted one. According to previous studies, thermal treatment of carob powder can cause hydrolysis of gallotannins and ellagitannins, which, in turn, releases gallic and ellagic acid units [4,26,32,33]. Therefore, the increased levels of polyphenols in roasted carob powders are justified and in agreement with the literature. The geographical origin of the samples also significantly influenced their polyphenolic profile. Although all samples originated from countries in the Mediterranean Basin, they appeared to have significant differences. In particular, the detected phenolic components of the Italian roasted carob powder were in lower concentrations compared to the Cypriot and Greek samples. As far as the phenolic profile of each sample is concerned, sinapic acid, quercitrin, and myricetin were not quantified in the powder from Italy, while the quantification of quercetin was not possible in powders from Cyprus and Greece.

Gallic acid was found to be the most abundant phenolic compound in carob syrups, while minor levels of catechin and epicatechin were also quantified. Most phenolic compounds under study were not detected in either thermally or non-thermally treated syrups. Caffeic acid was quantified only in syrups originating from Greece. Although Cypriot carob pulp and powder had a higher polyphenolic content than the corresponding Greek samples, there was no such trend in the case of syrups, probably due to the different processing methods used in these two countries. As observed, thermal treatment ($>40^{\circ}\text{C}$) had a negative effect on the phenolic content of carob syrups. The increase in temperature during syrup processing causes the thermal degradation of polyphenolic compounds due to decarboxylation reactions. The results of the present work are in agreement with the findings of Christou et al., who also reported a lower amount of polyphenols in thermally treated syrups than in non-thermally treated ones [6].

Overall, the obtained results via HPLC analysis concur with the spectrophotometric data for antioxidant activity and TPC of the samples. The thermal treatment and geographical origin of carob and its derived products have a significant role in their polyphenolic composition.

2.3. Chemometric Analysis

A low-dimensional model plane's systematic variation was estimated in a data matrix using PCA. Thirty samples were divided into three predefined groups according to their matrix (pulp, powder, and syrup). The first three principal components (PCs) of the extracted model explained 33.7% of the total variance in the data, while the first seven

components explained 41.9% of the total variability. As observed from the obtained PCA scatter plot results, the samples were adequately separated based on their matrix (Figure 3). Thermally treated samples were slightly separated from the other members of their groups. This was expected, considering their different chemical composition and antioxidant activity. Moreover, the group of carob powder demonstrated a great variation, due to the quite different phenolic composition of the Italian powder, which indicates the decisive role of the samples' geographical origin.

Furthermore, the data were analyzed statistically in order to find the correlation between the studied variables (phenolic compounds and antioxidant activity). The variable TPC demonstrated a strong correlation with FRAP, while a lower but significant correlation was also observed with DPPH. Rutin was highly correlated with catechin and epicatechin, while quercitrin and catechin appeared to have a significant correlation. Table S3 (Supplementary Materials) presents the correlation matrix of the model's variables.




Figure 3. PCA scatter plot (R^2X (cum) = 0.730, Q^2 (cum) = 0.337).

To discriminate the samples based on their matrix, OPLS-DA was effectively performed. As demonstrated in Figure 4, the samples were well discriminated. The extracted model was described via $R^2X = 0.877$ (the total sum of variation in X that is uncorrelated to Y), $R^2Y = 0.969$ (the total sum of variation in Y explained by the model), and $Q^2 = 0.936$ (the goodness of prediction). Furthermore, the OPLS-DA model was validated using cross-validation analysis (CV-ANOVA), with a p -value equal to 3.29×10^{-13} . To validate the model, the misclassification table was calculated (Table S4, Supplementary Materials). The obtained results demonstrate 100% correct classification of all samples, according to their matrix, with a highly satisfactory Fisher's probability reading equal to 4.8×10^{-13} .




Figure 4. Orthogonal partial least square-discriminant analysis model (R^2X (cum) = 0.946, R^2Y (cum) = 0.985, Q^2 (cum) = 0.975), discriminating the groups of carob pulps, powders, and syrups.

3. Materials and Methods

3.1. Standards and Reagents

For HPLC analysis, acetonitrile and methanol were purchased from Sigma Aldrich (St. Louis, MO, USA), while trifluoroacetic acid (TFA) was provided by Merck (Darmstadt, Germany). The analytical standards of catechin, myricetin, and quercitrin were acquired from HWI ANALYTIK GMBH (Rülzheim, Germany), while quercetin, epicatechin, sinapic acid, caffeic acid, and gallic acid were purchased from Sigma Aldrich (St. Louis, MO, USA). Rutin was provided by PhytoLab GmbH & Co (Vestenbergsgreuth, Germany).

Reagents for spectrophotometric assays, Folin–Ciocalteu reagent, sodium carbonate (Na_2CO_3), 2,4,6-tris(2-pyridyl)-s-triazine (TPTZ), iron(III) chloride hexahydrate ($\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$), and sodium acetate were purchased from Sigma Aldrich (St. Louis, MO, USA). DPPH was obtained from TCI (Tokyo, Japan).

3.2. Plant Material and Commercial Samples

A total of thirty samples were collected and analyzed to compare their antioxidant activity and polyphenolic profile. In particular, carob pods were collected during September (ripe pods) from Cyprus (Paphos, #1–3) and Greece (Crete, #4–6). The carob fruits were washed and crushed, the seeds were separated, and the pulp was then collected and lyophilized to remove moisture (LyoDry Compact Benchtop Freeze Dryer, Mechateck Systems, Bristol, UK). The lyophilized pulp was ground into a powder (Thermomix® TM5, Vorwerk, Wuppertal, Germany) and passed through a 250- μm sieve (Endecotts, London, UK) to obtain particles of uniform size. For comparison purposes, roasted pods collected from Greece (Crete, #7–9) were also studied to evaluate the effect of the roasting treatment, which was processed in the same way as raw carobs. In addition, seven carob-derived commercial products (powders and syrups) were purchased from local supermarkets. Specifically, the following products were studied: roasted powders from Italy (#10–12), Cyprus (Limassol, #13–15), and Greece (Crete, #16–18), raw powder from Greece (Crete, #19–21), thermally treated syrup from Cyprus (Limassol, #22–24) and Greece (Crete, #25–27), and non-thermally treated syrup from Greece (Crete, #28–30).

3.3. Extraction Procedure

Polyphenols were extracted from carob pulp and products according to the procedure described by Christou et al. [8]. Specifically, 2 g of each sample was mixed with 50 mL of 57% (*v/v*) acetone. The obtained mixture was then sonicated using an ultrasonic probe system (CY-500, Optic Ivymen System®, Barcelona, Spain) for 14 min, in a pulsed mode (5 s:5 s). After the ultrasound treatment, the obtained mixture was centrifuged at 4400 rpm for 20 min and filtered through filter paper. A rotary evaporator (RE300, Stuart®, Staffordshire, UK) was used to remove the organic solvent and condense the treated material before it was lyophilized and vacuum-stored at $-20\text{ }^\circ\text{C}$ until further analysis. The extraction procedure was performed three times for each sample.

3.4. Total Phenolic Content Using the Folin–Ciocalteu Method

The TPC of the extracts was determined using the Folin–Ciocalteu colorimetric method as described by Singleton et al. [41]. Prior to spectrophotometric analysis, the lyophilized extracts were redissolved in 20 mL of 57% acetone and filtered through a 0.45- μm pore size membrane filter to remove any insoluble material. Subsequently, 2 mL of properly diluted extract (1:10 (*v/v*) with distilled water) were mixed with 10 mL of Folin–Ciocalteu reagent (diluted to a concentration of 10% (*v/v*) with distilled water). After waiting 1–5 min, 8 mL of a saturated solution of Na_2CO_3 , 7.5% (*w/v*) were added to the reaction mixture and shaken vigorously. The absorbance of the samples was measured after 1 h of incubation in the dark and at room temperature at 765 nm using a UV-Vis spectrophotometer (UV-1900, Shimadzu, Tokyo, Japan). The results were expressed in terms of the mg of the gallic acid equivalents (GAE) per g of dry extract and for

comparison purposes as mg GAE/g of the carob pulp or product. All analyses were performed in triplicate and the results were expressed as the mean value \pm standard deviation.

3.5. Radical Scavenging Activity by DPPH Assay

The antioxidant capacity of the carob pulp and carob products was determined using the DPPH method, as previously described by Roseiro et al. [18]. The increased DPPH radical scavenging activity is indicated by the decrease in absorbance of the DPPH solution. When DPPH reacts with antioxidant components, which can donate a hydrogen atom to the DPPH radical, discoloration of the free radical is observed, from deep violet to light-yellow. The color changes are detected at 515 nm. Prior to the UV-Vis measurements, a methanolic solution of DPPH (60 μ M) was freshly prepared. A volume of 1950 μ L of DPPH solution was mixed with various concentrations of extract (50 μ L) and vortexed. The samples were kept in the dark at room temperature for 30 min and the decrease in absorbance was measured using a UV-Vis spectrophotometer (UV-1900, Shimadzu, Tokyo, Japan). The same volumes of methanol and DPPH solution were mixed to prepare the blank sample. All the analyses were performed in triplicate. The radical scavenging activity (%) was calculated as $((1 - \text{Abs}_{\text{ext}})/\text{Abs}_{\text{bl}}) \times 100$, where Abs_{ext} is the absorbance of the tested extract solution ($t = 30$ min) and Abs_{bl} is the absorption of the blank sample ($t = 0$ min). The concentration of extract (mg/mL) required to scavenge 50% of the DPPH radical was used to determine the half-maximal inhibitory concentration (IC_{50}) value.

3.6. Ferric Reducing Antioxidant Power (FRAP Assay)

The antioxidant potential of the extracts was determined via FRAP assay through the reduction of ferric ion (Fe^{3+}) to ferrous ion (Fe^{2+}), using the method described by Benzie and Strain [42]. The FRAP working solution was prepared by mixing the solutions of 300 mM acetate buffer (pH 3.6), 10 mM TPTZ in 40 mM HCl, and 20 mM $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ in the ratio of 10:1:1 and incubated at 37 $^{\circ}\text{C}$ for 30 min. The extract (150 μ L) was allowed to react with the FRAP working solution (2850 μ L) for 30 min in the dark. The absorbance of the blue-colored mixture was then measured at 593 nm using a UV-Vis spectrophotometer (UV-1900, Shimadzu, Tokyo, Japan). All experiments were performed in triplicate. Results were expressed as mM Trolox equivalents (TE) per gram of dry extract, using a Trolox standard curve.

3.7. Solid-Phase Extraction

An additional step to purify the extracts was necessary, before HPLC analysis. Solid-phase extraction (SPE) is a suitable technique for purifying extracts to remove co-extracted compounds. SPE was performed according to the method described by Christou et al. [8]. A vacuum manifold system with a 12-positions rack (VisiprepTM SPE Vacuum Manifold, SUPELCO, St. Louis, MO, USA) was utilized for all SPE experiments. Discovery DPA-6S (500 mg, SUPELCO, St. Louis, MO, USA) SPE polyamide cartridges were used for the purification of the extracts. The polyamide cartridges were preconditioned with 5 mL of methanol (MeOH) and equilibrated with 5 mL of acidified methanolic solution, 20:80 (v/v) MeOH: H_2O , acidified to pH 2.0 with HCl. The obtained lyophilized extracts were re-dissolved in acidified methanolic solution (20 mL) and then loaded into the cartridges. Subsequently, the cartridges were washed with 5 mL of acidified water (pH 2.0) to remove co-extracted compounds. Finally, the purified extracts were obtained after eluting the bound phenolics from the cartridges with an aqueous acetone mixture (2×5 mL, 80% v/v). The purified extracts were then dried using a rotary evaporator and lyophilizer. The dry residues were redissolved in an appropriate volume of MeOH, filtrated through a 0.45- μ m pore size membrane filter, properly diluted, and injected into the HPLC system.

3.8. HPLC-DAD

The phenolic profile of the carob pulp and its derived products was obtained with an HPLC system from Shimadzu (Kyoto, Japan), equipped with a photodiode array detector (PDA) (SPD-M20A), a pump (LC-10AD), an autosampler (SIL-20AHT), and a thermostat column compartment (CTO-10ASVP). Chromatographic separation conditions were performed according to the procedure described by Kumar et al. [43]. The PDA detector was set at 280 nm for the determination of all polyphenolic compounds. The Venusil XSP C18 (Radnor, PA, USA) analytical column (150 × 4.6 mm, 5 µm) was equipped with a pre-column composed of the same material and kept constant at 25 °C. Milli Q water (mobile phase A) and 0.02% (*v/v*) TFA in acetonitrile (mobile phase B) were used as solvents with a flow rate of 1 mL/min. A gradient elution program was used, starting at 80% A for the first five min; mobile phase A was decreased linearly to 60% at 8 min and then to 50% at 12 min. Finally, mobile phase A increased linearly to 60% until 17 min and then returned to the initial composition at 21 min, where it remained constant until 25 min. Twenty microliters of each extract were injected into the system and assayed in triplicate.

3.9. Data Analysis

All experimental assays were performed in triplicate. The results obtained were expressed as mean values ± standard deviation (SD). The means were compared, and statistical differences were obtained through a one-way analysis of variance (ANOVA) followed by Duncan's multiple range test at a 95% of confidence level.

For sample differentiation, the results obtained via HPLC and spectrophotometric assays were statistically processed using SIMCA 17.0.1 software (Umetrics, Umea, Sweden). Each sample was considered as an assembly of twelve variables represented by antioxidant activity and polyphenolic content. The data matrix (Dataset 12 variables × 30 observations × 3 groups) is provided in Supporting Information, Table S1.

For the initial overview of the data, a preliminary PCA was carried out. This procedure was mainly applied to estimate the systematic variation of the observations and to reduce the dimensionality of the data matrix. The input data were mean-centered with Unit Variate Scaling (UV) and the PCA model was extracted at a confidence level of 95%.

OPLS-DA increases the quality of the classification model by separating the systematic variation of X into two parts: one that is linearly related to Y (predictive information) and one that is unrelated to Y (orthogonal information) [44]. The extracted OPLS-DA model at a confidence level of 95% was UV-scaled and log-transformed. The effectiveness of the model was described using the goodness-of-fit R^2 ($0 \leq R^2 \leq 1$) and the predictive ability Q^2 values ($0 \leq Q^2 \leq 1$). The OPLS-DA model was validated using a cross-validation of variance (CV-ANOVA), with a *p*-value < 0.05.

4. Conclusions

It has over the years been demonstrated that the consumption of phenolic compounds present in foods may lower the risk of health disorders due to their antioxidant activity. In recent years, carob and its derived products have gained wide attention due to their health-promoting effects, which are mainly attributed to their phenolic compounds. In this work, the polyphenolic components of carob samples (carob pulps, powders, syrups) were analyzed via HPLC, the TPC via the Folin–Ciocalteu assay, and their antioxidant capacity via DPPH and FRAP tests. The thermal treatment of the samples significantly affected their phenolic composition. Particularly, the roasting of carob pulp and powder increased the concentrations of the polyphenolic compounds, while in thermally treated syrups, the temperature had a negative impact on the polyphenolic composition. The TPC, antioxidant activity, and phenolic concentrations were utilized as chemical markers in chemometric analysis. Both PCA and OPLS-DA were performed, with 100% successful classification of samples according to their matrix in OPLS-DA. As a future perspective, the use of high-resolution mass spectrometry will reveal more polyphenolic

compounds that will therefore provide more informative datasets used in chemometric analysis.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/molecules28052269/s1>, Figure S1: Loading score plot of the PCA model; Figure S2: Loading score plot of the OPLS-DA model; Table S1: Total phenolic content and antioxidant activity of carobs and carob-derived products; Table S2: Data matrix; Table S3: Correlation matrix of original variables (phenolics, TPC, antioxidant activity); Table S4: Misclassification table of OPLS-DA.

Author Contributions: Conceptualization, G.D.I. and A.C.; methodology, G.D.I., I.K.S. and A.C.; software, G.D.I. and A.C.; validation, G.D.I. and A.C.; investigation, G.D.I., I.K.S., A.C., and I.J.S.; resources, C.P.K.-C.; data curation, G.D.I. and A.C.; writing—original draft preparation, G.D.I.; writing—review and editing, A.C., I.J.S. and C.P.K.-C.; visualization, G.D.I., A.C., I.J.S. and C.P.K.-C.; supervision, C.P.K.-C.; project administration, C.P.K.-C.; funding acquisition, C.P.K.-C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was co-funded by the European Regional Development Fund and the Republic of Cyprus through the Research & Innovation Foundation (Project: BlackGold INTEGRATED /0916/0019).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available upon request from the corresponding author.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kyratzis, A.C.; Antoniou, C.; Papayiannis, L.C.; Graziani, G.; Rouphael, Y.; Kyriacou, M.C. Pod Morphology, Primary and Secondary Metabolite Profiles in Non-Grafted and Grafted Carob Germplasm Are Configured by Agro-Environmental Zone, Genotype, and Growing Season. *Front. Plant Sci.* **2021**, *11*, 612376. <https://doi.org/10.3389/fpls.2020.612376>.
2. Goulas, V.; Stylos, E.; Chatziathanasiadou, M.V.; Mavromoustakos, T.; Tzakos, A.G. Functional Components of Carob Fruit: Linking the Chemical and Biological Space. *Int. J. Mol. Sci.* **2016**, *17*, 1875. <https://doi.org/10.3390/ijms17111875>.
3. Antoniou, C.; Kyratzis, A.; Rouphael, Y.; Stylianou, S.; Kyriacou, M.C. Heat- And Ultrasound-Assisted Aqueous Extraction of Soluble Carbohydrates and Phenolics from Carob Kibbles of Variable Size and Source Material. *Foods* **2020**, *9*, 364. <https://doi.org/10.3390/foods9101364>.
4. Van Rijs, P.; Fogliano, V. Roasting Carob Flour Decreases the Capacity to Bind Glycoconjugates of Bile Acids. *Food Funct.* **2020**, *11*, 5924–5932. <https://doi.org/10.1039/d0fo01158d>.
5. Cavdarova, M.; Makris, D.P. Extraction Kinetics of Phenolics from Carob (*Ceratonia siliqua* L.) Kibbles Using Environmentally Benign Solvents. *Waste Biomass Valorization* **2014**, *5*, 773–779. <https://doi.org/10.1007/s12649-014-9298-3>.
6. Christou, A.; Martinez-piernas, A.B.; Stavrou, I.J.; Garcia-reyes, J.F.; Kapnissi-christodoulou, C.P. HPLC-ESI-HRMS and Chemometric Analysis of Carobs Polyphenols—Technological and Geographical Parameters Affecting Their Phenolic Composition. *J. Food Compos. Anal.* **2022**, *114*, 104744. <https://doi.org/10.1016/j.jfca.2022.104744>.
7. Kokkinofa, R.; Yiannopoulos, S.; Stylianou, M.A.; Agapiou, A. Use of Chemometrics for Correlating Carobs Nutritional Compositional Values with Geographic Origin. *Metabolites* **2020**, *10*, 62. <https://doi.org/10.3390/metabo10020062>.
8. Christou, A.; Stavrou, I.J.; Kapnissi-Christodoulou, C.P. Continuous and Pulsed Ultrasound-Assisted Extraction of Carob's Antioxidants: Processing Parameters Optimization and Identification of Polyphenolic Composition. *Ultrason. Sonochem.* **2021**, *76*, 105630. <https://doi.org/10.1016/j.ultsonch.2021.105630>.
9. Chait, Y.A.; Gunenc, A.; Bendali, F.; Hosseinian, F. Simulated Gastrointestinal Digestion and in Vitro Colonic Fermentation of Carob Polyphenols: Bioaccessibility and Bioactivity. *Lwt* **2020**, *117*, 108623. <https://doi.org/10.1016/j.lwt.2019.108623>.
10. Stavrou, I.J.; Christou, A.; Kapnissi-Christodoulou, C.P. Polyphenols in Carobs: A Review on Their Composition, Antioxidant Capacity and Cytotoxic Effects, and Health Impact. *Food Chem.* **2018**, *269*, 355–374. <https://doi.org/10.1016/j.foodchem.2018.06.152>.
11. Christodoulou, M.C.; Orellana Palacios, J.C.; Hesami, G.; Jafarzadeh, S.; Lorenzo, J.M.; Domínguez, R.; Moreno, A.; Hadidi, M. Spectrophotometric Methods for Measurement of Antioxidant Activity in Food and Pharmaceuticals. *Antioxidants* **2022**, *11*, 2213. <https://doi.org/10.3390/antiox11112213>.
12. Scalbert, A.; Manach, C.; Morand, C.; Rémésy, C.; Jiménez, L. Dietary Polyphenols and the Prevention of Diseases. *Crit. Rev. Food Sci. Nutr.* **2005**, *45*, 287–306. <https://doi.org/10.1080/1040869059096>.

13. Ruiz-Roso, B.; Quintela, J.C.; de la Fuente, E.; Haya, J.; Pérez-Olleros, L. Insoluble Carob Fiber Rich in Polyphenols Lowers Total and Ldl Cholesterol in Hypercholesterolemic Subjects. *Plant Foods Hum. Nutr.* **2010**, *65*, 50–56. <https://doi.org/10.1007/s11130-009-0153-9>.
14. Gregoriou, G.; Neophytou, C.M.; Vasincu, A.; Gregoriou, Y.; Hadjipakkou, H.; Pinakoulaki, E.; Christodoulou, M.C.; Ioannou, G.D.; Stavrou, I.J.; Christou, A.; et al. Anti-Cancer Activity and Phenolic Content of Extracts Derived from Cypriot Carob (*Ceratonia siliqua* L.) Pods Using Different Solvents. *Molecules* **2021**, *26*, 5017. <https://doi.org/10.3390/molecules26165017>.
15. Vekari, S.A.; Ouzounidou, G.; Ozturk, M.; Gök, G. Variation of Quality Characteristics in Greek and Turkish Carob Pods during Fruit Development. *Procedia-Soc. Behav. Sci.* **2011**, *19*, 750–755. <https://doi.org/10.1016/j.sbspro.2011.05.194>.
16. Saci, F.; Bachir bey, M.; Louaileche, H.; Gali, L.; Bensouici, C. Changes in Anticholinesterase, Antioxidant Activities and Related Bioactive Compounds of Carob Pulp (*Ceratonia siliqua* L.) during Ripening Stages. *J. Food Meas. Charact.* **2020**, *14*, 937–945. <https://doi.org/10.1007/s11694-019-00344-9>.
17. Quiles-Carrillo, L.; Mellinas, C.; Garrigos, M.C.; Balart, R.; Torres-Giner, S. Optimization of Microwave-Assisted Extraction of Phenolic Compounds with Antioxidant Activity from Carob Pods. *Food Anal. Methods* **2019**, *12*, 2480–2490. <https://doi.org/10.1007/s12161-019-01596-3>.
18. Roseiro, L.B.; Tavares, C.S.; Roseiro, J.C.; Rauter, A.P. Antioxidants from Aqueous Decoction of Carob Pods Biomass (*Ceratonia siliqua* L.): Optimisation Using Response Surface Methodology and Phenolic Profile by Capillary Electrophoresis. *Ind. Crops Prod.* **2013**, *44*, 119–126. <https://doi.org/10.1016/j.indcrop.2012.11.006>.
19. Petkova, N.; Petrova, I.; Ivanov, I.; Mihov, R.; Hadjikinova, R.; Ognyanov, M.; Nikolova, V. Nutritional and Antioxidant Potential of Carob (*Ceratonia siliqua*) Flour and Evaluation of Functional Properties of Its Polysaccharide Fraction. *J. Pharm. Sci. Res.* **2017**, *9*, 2189–2195.
20. Roseiro, L.B.; Duarte, L.C.; Oliveira, D.L.; Roque, R.; Bernardo-Gil, M.G.; Martins, A.I.; Sepúlveda, C.; Almeida, J.; Meireles, M.; Gírio, F.M.; et al. Supercritical, Ultrasound and Conventional Extracts from Carob (*Ceratonia siliqua* L.) Biomass: Effect on the Phenolic Profile and Antiproliferative Activity. *Ind. Crops Prod.* **2013**, *47*, 132–138. <https://doi.org/10.1016/j.indcrop.2013.02.026>.
21. Benchikh, Y.; Louaileche, H.; George, B.; Merlin, A. Changes in Bioactive Phytochemical Content and in Vitro Antioxidant Activity of Carob (*Ceratonia siliqua* L.) as Influenced by Fruit Ripening. *Ind. Crops Prod.* **2014**, *60*, 298–303. <https://doi.org/10.1016/j.indcrop.2014.05.048>.
22. Goulas, V.; Georgiou, E. Utilization of Carob Fruit as Sources of Phenolic Compounds with Antioxidant Potential: Extraction Optimization and Application in Food Models. *Foods* **2020**, *9*, 20. <https://doi.org/10.3390/foods9010020>.
23. Owen, R.W.; Haubner, R.; Hull, W.E.; Erben, G.; Spiegelhalder, B.; Bartsch, H.; Haber, B. Isolation and Structure Elucidation of the Major Individual Polyphenols in Carob Fibre. *Food Chem. Toxicol.* **2003**, *41*, 1727–1738. [https://doi.org/10.1016/S0278-6915\(03\)00200-X](https://doi.org/10.1016/S0278-6915(03)00200-X).
24. Rakib, E.M.; Chicha, H.; Abouricha, S.; Alaoui, M. Determination of Phenolic Composition of Carob Pods Grown in Different Regions of Morocco. *J. Nat. Prod.* **2010**, *3*, 134–140.
25. Christou, A.; Stavrou, I.J.; Kapnissi-Christodoulou, C.P. Combined Use of β -Cyclodextrin and Ionic Liquid as Electrolyte Additives in EKC for Separation and Determination of Carob's Phenolics—A Study of the Synergistic Effect. *Electrophoresis* **2021**, *42*, 1945–1955. <https://doi.org/10.1002/elps.202100085>.
26. Papagiannopoulos, M.; Wollseifen, H.R.; Mellenthin, A.; Haber, B.; Galensa, R. Identification and Quantification of Polyphenols in Carob Fruits (*Ceratonia siliqua* L.) and Derived Products by HPLC-UV-ESI/MSn. *J. Agric. Food Chem.* **2004**, *52*, 3784–3791. <https://doi.org/10.1021/jf030660y>.
27. Krokou, A.; Kokkinofa, R.; Stylianou, M.; Agapiou, A. Decoding Carob Flavor Aroma Using HS-SPME-GC-MS and Chemometrics. *Eur. Food Res. Technol.* **2020**, *246*, 1419–1428. <https://doi.org/10.1007/s00217-020-03500-4>.
28. Papaefstathiou, E.; Agapiou, A.; Giannopoulos, S.; Kokkinofa, R. Nutritional Characterization of Carobs and Traditional Carob Products. *Food Sci. Nutr.* **2018**, *6*, 2151–2161. <https://doi.org/10.1002/fsn3.776>.
29. Alabdi, F.; Elharfi, N.; Balouki, A.; Kzaiber, F.; Oussama, A. Carob Origin Classification by FTIR Spectroscopy and Chemometrics. *J. Chem. Chem. Eng* **2011**, *5*, 1020–1029.
30. Christou, C.; Agapiou, A.; Kokkinofa, R. Use of FTIR Spectroscopy and Chemometrics for the Classification of Carobs Origin. *J. Adv. Res.* **2018**, *10*, 1–8. <https://doi.org/10.1016/j.jare.2017.12.001>.
31. Farag, M.A.; El-Kersh, D.M.; Ehrlich, A.; Choucry, M.A.; El-Seedi, H.; Frolov, A.; Wessjohann, L.A. Variation in *Ceratonia Siliqua* Pod Metabolome in Context of Its Different Geographical Origin, Ripening Stage and Roasting Process. *Food Chem.* **2019**, *283*, 675–687. <https://doi.org/10.1016/j.foodchem.2018.12.118>.
32. Eldeeb, G.S.S.; Mosilhey, S.H. Roasting Temperature Impact on Bioactive Compounds and PAHs in Carob Powder (*Ceratonia siliqua* L.). *J. Food Sci. Technol.* **2022**, *59*, 105–113. <https://doi.org/10.1007/s13197-021-04989-7>.
33. Şahin, H.; Topuz, A.; Pischetsrieder, M.; Özdemir, F. Effect of Roasting Process on Phenolic, Antioxidant and Browning Properties of Carob Powder. *Eur. Food Res. Technol.* **2009**, *230*, 155–161. <https://doi.org/10.1007/s00217-009-1152-7>.
34. Kyriacou, M.C.; Antoniou, C.; Roupheal, Y.; Graziani, G.; Kyrtzias, A. Mapping the Primary and Secondary Metabolomes of Carob (*Ceratonia siliqua* L.) Fruit and Its Postharvest Antioxidant Potential at Critical Stages of Ripening. *Antioxidants* **2021**, *10*, 57. <https://doi.org/10.3390/antiox10010057>.
35. Žugić, A.; Dorđević, S.; Arsić, I.; Marković, G.; Živković, J.; Jovanović, S.; Tadić, V. Antioxidant Activity and Phenolic Compounds in 10 Selected Herbs from Vrujci Spa, Serbia. *Ind. Crops Prod.* **2014**, *52*, 519–527. <https://doi.org/10.1016/j.indcrop.2013.11.027>.

36. Sethi, S.; Joshi, A.; Arora, B.; Bhowmik, A.; Sharma, R.R.; Kumar, P. Significance of FRAP, DPPH, and CUPRAC Assays for Antioxidant Activity Determination in Apple Fruit Extracts. *Eur. Food Res. Technol.* **2020**, *246*, 591–598. <https://doi.org/10.1007/s00217-020-03432-z>.
37. Ou, B.; Huang, D.; Hampsch-Woodill, M.; Flanagan, J.A.; Deemer, E.K. Analysis of Antioxidant Activities of Common Vegetables Employing Oxygen Radical Absorbance Capacity (ORAC) and Ferric Reducing Antioxidant Power (FRAP) Assays: A Comparative Study. *J. Agric. Food Chem.* **2002**, *50*, 3122–3128. <https://doi.org/10.1021/jf0116606>.
38. Huma, Z.E.; Jayasena, V.; Nasar-Abbas, S.M.; Imran, M.; Khan, M.K. Process Optimization of Polyphenol Extraction from Carob (*Ceratonia siliqua*) Kibbles Using Microwave-Assisted Technique. *J. Food Process. Preserv.* **2018**, *42*, e13450. <https://doi.org/10.1111/jfpp.13450>.
39. Benchikh, Y.; Louailèche, H. Effects of Extraction Conditions on the Recovery of Phenolic Compounds and in Vitro Antioxidant Activity of Carob (*Ceratonia siliqua* L.) Pulp. *Acta Bot. Gall.* **2014**, *161*, 175–181. <https://doi.org/10.1080/12538078.2014.909325>.
40. Buchner, N.; Krumbein, A.; Rohn, S.; Kroh, L.W. Effect of Thermal Processing on the Flavonols Rutin and Quercetin. *Rapid Commun. Mass Spectrom.* **2006**, *20*, 3229–3235. <https://doi.org/10.1002/rcm.2720>.
41. Singleton, V.L.; Rossi, J.A. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am. J. Enol. Vitic.* **1965**, *16*, 144–158.
42. Benzie, I.F.F.; Strain, J.J. The Ferric Reducing Ability of Plasma (FRAP) as a Measure of ‘Antioxidant Power’: The FRAP Assay. *Anal. Biochem.* **1996**, *239*, 70–76. <https://doi.org/10.1006/abio.1996.0292>.
43. Kumar, N.; Bhandari, P.; Singh, B.; Gupta, A.P.; Kaul, V.K. Reversed Phase-HPLC for Rapid Determination of Polyphenols in Flowers of Rose Species. *J. Sep. Sci.* **2008**, *31*, 262–267. <https://doi.org/10.1002/jssc.200700372>.
44. Trygg, J.; Holmes, E.; Lundstedt, T. Chemometrics in Metabonomics. *J. Proteome Res.* **2007**, *6*, 469–479. <https://doi.org/10.1021/pr060594q>.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.