



Article Periphery Exploration around 2,6-Diazaspiro[3.4]Octane Core Identifies a Potent Nitrofuran Antitubercular Lead

Alexei Lukin¹, Kristina Komarova¹, Lyubov Vinogradova¹, Marine Dogonadze², Tatiana Vinogradova², Piotr Yablonsky², Alexander Kazantsev³ and Mikhail Krasavin^{3,4,*}

- ¹ Lomonosov Institute of Fine Chemical Technologies, MIREA—Russian Technological University, 119454 Moscow, Russia
- ² Saint Petersburg Research Institute of Phthisiopulmonology, 2-4 Ligovsky Prospekt, 191036 Saint Petersburg, Russia
- ³ Institute of Chemistry, Saint Petersburg State University, 26 Universitetskii Prospect, 198504 Peterhof, Russia
- ⁴ Institute of Living Systems, Immanuel Kant Baltic Federal University, 236041 Kaliningrad, Russia
- * Correspondence: m.krasavin@spbu.ru

Abstract: A small set of twelve compounds of a nitrofuran carboxamide chemotype was elaborated from a readily available 2,6-diazaspiro[3.4] octane building block, exploring diverse variants of the molecular periphery, including various azole substituents. The in vitro inhibitory activities of the synthesized compounds were assessed against *Mycobacterium tuberculosis* H37Rv. As a result, a remarkably potent antitubercular lead displaying a minimal inhibitory concentration of 0.016 μ g/mL was identified.

Keywords: 2,6-diazaspiro[3.4]octane; diversity-oriented synthesis; molecular periphery; azoles; antitubercular activity



Citation: Lukin, A.; Komarova, K.; Vinogradova, L.; Dogonadze, M.; Vinogradova, T.; Yablonsky, P.; Kazantsev, A.; Krasavin, M. Periphery Exploration around 2,6-Diazaspiro[3.4]Octane Core Identifies a Potent Nitrofuran Antitubercular Lead. *Molecules* **2023**, *28*, 2529. https://doi.org/10.3390/ molecules28062529

Academic Editor: László Somsák

Received: 5 February 2023 Revised: 7 March 2023 Accepted: 7 March 2023 Published: 10 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

1. Introduction

5-Nitrofuryl-substituted chemotypes have been a prolific source of compounds endowed with antimycobacterial activity due to the ability of the nitrofuran moiety to undergo reduction with the bacterial enzyme machinery and generate reactive intermediates that are lethal to the bacterium itself [1]. The same mode of biotransformation of nitrofurans by mammalian cells is considered the likely origin of the potential toxicity associated with this and other nitroheterocycles [2]. However, the toxicity of nitrofurans is highly dependent on the specific molecular moiety to which these heterocycles are conjugated, and by optimizing the nitrofuran's molecular periphery, it is possible not only to increase the potency of nitrofuran antibacterials but also alleviate the systemic toxicity of this class of drugs. Indeed, we have previously shown that by linking the 5-nitrofuryl moiety to various heterocyclic motifs, potent antimycobacterial compounds efficacious against multidrug-resistant strains can be developed that are also non-toxic to rodents [3]. Later, it became evident that the conjugation of the 5-nitrofuryl fragment with saturated spirocyclic piperidines was very important for the implementation of this strategy [4]. This motivated us to continue studies along this line of inquiry.

The fully saturated, high- F_{sp3} 2,6-diazaspiro[3.4]octane core is an emerging privileged structure, taking into account its recent frequent appearance in compounds possessing diverse biological activities. Notable recently reported examples include a hepatitis B capsid protein inhibitor [5], a menin-MLL1 interaction inhibitor for the treatment of cancer [6], a MAP and PI3K signaling modulator [7], a selective dopamine D₃ receptor antagonist [8], and VDAC1 inhibitors for the treatment of diabetes [9] (Figure 1).



Figure 1. Examples of diverse biologically active compounds containing 2,6-diazaspiro[3.4]octane motif.

Considering the significance of the 2,6-diazaspiro[3.4]octane motif in recent drug discovery efforts and the availability of convenient procedures in the literature to access this spirocycle in a functionalized form, we were prompted to synthesize a diverse set of nitrofuran derivatives based on this core that would contain various periphery groups, including a range of azoles. The idea was to demonstrate that by broadly exploring the molecular periphery, potent antitubercular compound(s) could be identified within a relatively small screening set. Herein, we report our findings gathered in the course of realizing this idea.

2. Results and Discussion

The synthesis of core building block **3** on a multigram scale was achieved from commercially available *N*-Boc-protected azetidine-3-one **1**, as described in the literature [10]. The initial Horner-Wadsworth-Emmons olefination proceeded in high yield and gave α , β -unsaturated ester **2** following chromatographic purification. A [3 + 2] cycloaddition with **2** and *N*-(methoxymethyl)-*N*-(trimethylsilylmethyl) benzylamine in the presence of lithium fluoride afforded intermediate core building block **3** at 56% yield, which was sufficiently pure without chromatography. The ester functionality of **3** was manipulated first by one-pot alkaline hydrolysis and then transformation to amides **4a–d**. Amides **4a–c** were transformed into 5-nitrofuroyl-substituted compounds **5a–c** via a two-step deprotection-acylation protocol. Weinreb amide **4d** was reduced to the respective aldehyde, which was subjected to reductive amination with dimethylamine to give compound **5d** at 72% yield over two steps. Finally, compound **5d** was transformed into 5-nitrofuroyl derivative **6d** analogously to amides **4a–c** (Scheme 1).

Next, we aimed to elaborate the ester functionality in **3** into an imidazole and an oxazole moiety. In order to reduce the lipophilicity of the scaffold, we removed the benzyl group by hydrogenation (to obtain **7**) and replaced it with a methyl group using a reductive alkylation procedure. After ester hydrolysis in intermediate **8**, we prepared propargylamide **9**, which was viewed as a starting material for both imidazole and oxazole. Following the protocol from the Beller group [11], propargylamide **9**, without purification, underwent $Zn(OTf)_2$ -catalyzed hydroamination with benzylamine, which led to cyclodehydrative aromatization and the formation of imidazole **10** at 75% yield. In the absence of benzylamine (or any other primary amine), cycloaromatization under the same conditions led to the formation of oxazole **11** [12]. Compounds **10** and **11** were transformed into their *N*-(5-nitrofuroyl) derivatives using the same one-pot deprotection-acylation protocol as described above to furnish nitrofurans **12** and **13**, respectively (Scheme 2).



Scheme 1. Synthesis of core building block **3** and manipulation of the ester group via amidation. *Reagents and conditions:* i. $EtO_2CCH_2P(O)(OEt)_2$, NaH, THF, 0 °C \rightarrow r. t. (92%); ii. MeOCH₂N(Bn)CH₂SiMe₃, LiF, MeCN, 60 °C, 18 h (56%); iii. LiOH·H2O, aq. 1,4-dioxane, r. t., 12 h; iv. RNHR', HOBt, EDC·HCl, r. t., 12 h (4a 66%, 4b 68%, 4c 72%, 4d 72%, 2 steps); v. TFA, CH₂Cl₂, 0 °C \rightarrow r. t., 1 h; vi. 5-nitro-2-furoic acid, CDI, DMF, r. t., 1 h (5a 56%, 5b 56%, 5c 48%, 6d 38%, 2 steps); vii. LiAlH₄, THF, -70 °C \rightarrow -5 °C; viii. NaBH(OAc)₃, CH₂Cl₂, r. t., 18 h (72%, 2 steps).



Scheme 2. Synthesis of imidazole (12) and oxazole (13) derivatives. *Reagents and conditions:* i. H₂, 10% Pd-C, 100 atm, r. t., 12 h (quant.); ii. 37% aq. formaldehyde, CH_2Cl_2 , $NaBH(OAc)_3$, r. t., 18 h (72%); iii. LiOH·H₂O, aq. 1,4-dioxane, r. t., 12 h; iv. propargylamine, HOBt, EDC·HCl, r. t., 12 h (66%, 2 steps); v. BnNH₂, Zn(OTf)₂ (0.25 mol.%), toluene, reflux with Dean-Stark trap, 8 h (75%); vi. Zn(OTf)₂ (0.25 mol.%), toluene, reflux with Dean-Stark trap, 0 °C \rightarrow r. t., 1 h; viii. 5-nitro-2-furoic acid, CDI, DMF, r. t., 1 h (28%, 2 steps).

Next, we turned our attention to elaborating the ester functionality in **3** into the 4methyl-1,2,4-triazole group while also varying the substituent on the pyrrolidine nitrogen atom. Compound **3** was converted to hydrazide **14** by refluxing with hydrazine in ethanol and used without purification. Hydrazide **14** was reacted with methyl isothiocyanate and the intermediate hydrazine carbothioamide was treated with base and the cyclized intermediate with Raney nickel, which triggered the conversion into 1,2,4-triazole **15**. On the one hand, the latter was converted into 5-nitrofuroyl derivative **18**, and on the other hand, **15** was subjected to benzyl-to-mesyl group swap on the nitrogen atom to give *N*-mesyl derivative **16**, which was converted into 5-nitrofuroyl compound **17** (Scheme 3).



Scheme 3. Synthesis of 4-methyl-1,2,4-triazol-3-yl derivatives. *Reagents and conditions:* i. 64% aq. N₂H₄, ethanol, reflux, 8 h (quant.); ii. MeNCS, ethanol, reflux, 2 h; iii. sat. aq. K₂CO₃, reflux, 8 h iv. Raney Ni, ethanol, reflux, 12 h (58%, 3 steps); v. TFA, CH₂Cl₂, 0 °C \rightarrow r. t., 1 h; vi. 5-nitro-2-furoic acid, CDI, DMF, r. t., 1 h (17 56%, 18 62%, 2 steps); vii. H₂, 10% Pd-C, 100 atm, r. t., 12 h; viii. MsCl, Et₃N, CH₂Cl₂, 0 °C \rightarrow r. t., 18 h (53%, 2 steps).

Further, we were interested in replacing the ester functionality in **3** with a differently substituted 1,2,4-triazole moiety. To this end, hydrazide **14** was reacted with two amidines (acetamidine and cyclopropane carboxamidine). The initial adduct was cyclized on heating to 170 °C to give 1,2,4-triazoles **19** and **20**, respectively. Each of the latter was converted into a 5-nitrofuroyl derivative (**21** and **22**, respectively) using the already established one-pot, two-step protocol (Scheme 4).



Scheme 4. Synthesis of 5-substituted 1,2,4-triazol-3-yl derivatives. *Reagents and conditions:* i. acetamidine (cyclopropane carboxamidine) HCl, MeONa, MeOH, r. t., 18 h; ii 170 °C in argon stream (**19** 48%, **20** 42%, 2 steps); iii. TFA, CH₂Cl₂, 0 °C \rightarrow r. t., 1 h; iv. 5-nitro-2-furoic acid, CDI, DMF, r. t., 1 h (**21** 43%, **22** 38%, 2 steps).

Finally, we were interested in grafting a 1,2,4-oxadiazole moiety in lieu of the ester group on the 2,6-diazaspiro[3.4]octane building block **3** while also varying the substituent on the pyrrolidine nitrogen atom. To this end, we hydrolyzed the ester functionality in **3**, acylated acetamidoxime with the resulting acid, and subjected the resulting intermediate to TBAF-promoted cyclodehydration. The resulting 1,2,4-oxadiazole **23** was transformed into 5-nitrofuroyl derivative **24**. *N*-Unsubstituted ester **7** from our previous synthesis depicted in Scheme **2** was *N*-mesylated and subjected to ester hydrolysis to furnish carboxylic acid



Scheme 5. Synthesis of 1,2,4-oxadiazole derivatives. *Reagents and conditions:* i. LiOH·H₂O, aq. 1,4-dioxane, r. t., 12 h; ii. acetamidoxime, HOBt, EDC·HCl, r. t., 12 h; iii. TBAF, toluene, Dean-Stark trap, reflux, 6 h (**23** 41%, **26** 63%, 3 steps); iv. TFA, CH₂Cl₂, 0 °C \rightarrow r. t., 1 h; v. 5-nitro-2-furoic acid, CDI, DMF, r. t., 1 h (**24** 48%, **27** 63%, 2 steps); vi. MsCl, Et₃N, CH₂Cl₂, 0 °C \rightarrow r. t., 18 h (quant.).

Having thus synthesized twelve 5-nitrofuroyl derivatives (**5a–c**, **6d**, **12–13**, **17–18**, **21–22**, **24**, and **27**), we proceeded to evaluate their activity against the drug-sensitive strain of *Mycobacterium tuberculosis* H37Rv using the resazurin microtiter plate assay (REMA) [13]. The resulting data expressed as minimum inhibitory concentrations (MIC) are collated in Table 1.

| Compound | Yield, % | MIC, μg/mL * |
|----------|----------|--------------|
| 5a | 56 | 25 |
| 5b | 56 | 50 |
| 5c | 48 | 100 |
| 6d | 38 | >100 |
| 12 | 28 | 25 |
| 13 | 48 | >100 |
| 17 | 56 | 0.016 |
| 18 | 62 | 100 |
| 21 | 43 | 50 |
| 22 | 38 | 50 |
| 24 | 48 | 6.2 |
| 27 | 63 | >100 |

Table 1. Yields and growth inhibitory activity of 5-nitrofuroyl derivatives against drug-sensitive *M. tuberculosis* H37Rv strain.

* Determined chromogenically in 6 repeated experiments.

As follows from the data summarized in Table 1, most of the compounds were either weakly active or inactive against this strain of *M. tuberculosis*. However, two specific compounds (out of only twelve)—1,2,4-oxadiazole **24** and especially 1,2,4-triazole **17**—displayed quite respectable levels of activity. In fact, compound **17** was very active and caused the bacteria to die at a minimum inhibitory concentration of 0.016 μ g/mL.

Comparing the activity of compounds **17** and **18**, it would be assumed that the N-mesyl group strongly promoted activity. On the other hand, when comparing compounds **18** and **24**, one can note an increase in activity when replacing 1,2,4- triazole with 1,2,4-

oxadiazole. Based on these two comparisons, one should expect outstanding results from compound **27**, but this was not reflected in the data in Table 1. Therefore, there was no unambiguous correlation between structure and property, and further research is required.

3. Material and Methods

3.1. Synthesis—General

All commercial reagents were used without purification. NMR spectra were recorded using a Bruker DPX-300 spectrometer in CDCl₃ (¹H: 300 MHz; ¹³C: 75 MHz). Chemical shifts are reported as parts per million (δ , ppm). The residual solvent peak (CHCl₃ or DMSO-*d*₆) was used as the internal standard: 7.28 or 2.51 for ¹H and 77.07 or 40.00 ppm for ¹³C. Multiplicities are abbreviated as follows: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad, dd = doublet of doublets, dt = doublet of triplets, ddd = doublet/doublets of doublets (see Supplementary Materials). Coupling constants, J, are reported in Hz. Mass spectra were recorded using a Bruker microTOF spectrometer (ionization by electrospray, positive ion detection). Melting points were determined in open capillary tubes using a Stuart SMP50 Automatic Melting Point Apparatus. Analytical thinlayer chromatography was carried out on UV-254 silica gel plates using appropriate eluents. Compounds were visualized with short-wavelength UV light. Column chromatography was performed using silica gel Merk grade 60 (0.040–0.063 mm) 230–400 mesh. All reactions were conducted under an atmosphere of argon.

3.2. 2-(tert-Butyl) 8-ethyl 6-benzyl-2,6-diazaspiro[3.4]octane-2,8-dicarboxylate (3)

Prepared according to the literature procedure [10] with minor modifications. To a suspension of NaH (60% dispersion in mineral oil, 1.88 g, 0.047 mol, 1.15 equiv.) in THF (150 mL) at 0 °C, triethylphosphonoacetate (11 g, 0.054 mol, 1.2 equiv.) was added. The resulting mixture was allowed to warm up to r. t. and stirred at that temperature for 30 min. It was cooled down to 0 °C, at which point *tert*-butyl 3-oxazetidine-1-carboxylate (7 g, 0.041 mol, 1.0 equiv.) in THF (50 mL) was added. The reaction mixture was allowed to reach r. t. again and stirred at that temperature overnight. It was then diluted with ethyl acetate and washed with sat. aq. NaHCO₃, water, and brine. The organic phase was separated and dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by column chromatography on silica, eluting with $0\rightarrow10\%$ ethyl acetate in hexane. Yield of **2**—9 g (92%), colorless oil. The spectral data of compound **2** were in accordance with those reported in the literature: ¹H NMR (300 MHz, CDCl₃) δ 5.74 (dd, *J* = 4.5, 2.2 Hz, 1H), 4.80 (dd, *J* = 6.3, 2.9 Hz, 2H), 4.57 (dt, *J* = 5.3, 2.7 Hz, 2H), 4.31–4.02 (m, 2H), 1.45 (s, 9H), 1.26 (t, *J* = 7.1 Hz, 3H); LCMS (ESI): *m/z* (M + H) calcd, 242.3; found, 242.2.

To a solution of **2** (5 g, 20.75 mmol, 1 equiv.) in acetonitrile (50 mL), LiF (2.15 g, 83 mmol, 4 equiv.) and (methoxymethyl)-1-phenyl-*N*-(trimethylsilylmethyl)methanamine (6.25 g, 25 mmol, 1.2 equiv.) were added and the resulting mixture was stirred at 60 °C overnight. The solvent was removed in vacuo and the residue was dissolved in ethyl acetate (50 mL). The solution was washed with sat. aq. citric acid (3 × 25 mL). The combined aqueous phases were extracted with ethyl acetate (2 × 100 mL). The pH of the aqueous phase was brought to 8 with sat. aq. K₂CO₃ and it was extracted with ethyl acetate (2 × 100 mL). The combined organic phases were dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo to give 4.3 g (56%) of the title product as a colorless oil. This product was used in subsequent steps without further purification. The spectral data of compound **3** were in accordance with those reported in the literature: ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.46–7.14 (m, 5H), 4.22–4.00 (m, 2H), 3.94 (dd, *J* = 16.6, 8.4 Hz, 2H), 3.78–3.51 (m, 5H), 3.12 (t, *J* = 7.6 Hz, 1H), 2.83–2.66 (m, 4H), 1.36 (s, 9H), 1.19 (t, *J* = 7.1 Hz, 3H); LCMS (ESI): *m/z* (M + H⁺) calcd, 375.5; found, 375.4.

3.3. 2-tert-Butoxycarbonyl-6-benzyl-8-(methylcarbamoyl)-2,6-diazaspiro[3.4]octane (4a)

General Procedure A. To a solution of **3** (1 g, 3.35 mmol, 1.0 equiv.) in 1,4-dioxane (10 mL) was added aqueous solution (1 mL) of LiOH·H₂O (0.175 g, 4.2 mmol, 1.25 equiv.) and the resulting mixture was stirred for 12 h. HOBt (0.56 g, 4.2 mmol,1.25 equiv.), EDC·HCl (0.81 g, 4.2 mmol, 1.25 equiv.), and methylamine (1.4 equiv.) were added and the resulting mixture was stirred for 12 h. The solvent was removed in vacuo and the residue was dissolved in ethyl acetate (50 mL). The solution was washed with 10% aqueous K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo to obtain 0.68 g (66%) of the title compound as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.38–7.19 (m, 5H), 7.00 (s, 1H), 3.88 (d, *J* = 4.4 Hz, 1H), 3.85 (d, *J* = 5.0 Hz, 1H), 3.79 (d, *J* = 8.4 Hz, 1H), 3.69 (d, *J* = 9.1 Hz, 1H), 3.63 (s, 1H), 3.01 (d, *J* = 9.7 Hz, 1H), 2.89–2.62 (m, 6H), 1.39 (s, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 173.52, 156.08, 138.07, 128.53, 128.45, 127.33, 79.56, 65.33, 59.44, 57.16, 53.85, 42.07, 28.28, 25.99; HRMS (ESI) *m/z* calcd for C₂₀H₃₀N₃O₃ [M + H⁺] 360.2287, found 360.2291.

3.4. 2-tert-Butoxycarbonyl-6-benzyl-8-(isobutylcarbamoyl)-2,6-diazaspiro[3.4]octane (4b)

This compound was synthesized according to general procedure A from **3** (0.5 g, 1.33 mmol). Yield 0.36 g (68%), colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.40–7.10 (m, 6H), 3.90 (d, *J* = 8.9 Hz, 2H), 3.80 (d, *J* = 8.5 Hz, 1H), 3.69 (d, *J* = 9.1 Hz, 1H), 3.63 (s, 2H), 3.15–2.98 (m, 3H), 2.92 (dd, *J* = 9.6, 2.7 Hz, 1H), 2.80 (dd, *J* = 6.5, 2.7 Hz, 1H), 2.73–2.52 (m, 2H), 1.82–1.60 (m, 1H), 1.39 (s, 9H), 0.87 (dd, *J* = 6.7, 1.0 Hz, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 178.85, 167.17, 155.96, 138.18, 128.53, 128.37, 127.24, 79.69, 64.33, 59.41, 57.47, 43.93, 43.22, 29.65, 28.26, 11.60, -0.05; HRMS (ESI) *m*/*z* calcd for C₂₃H₃₆N₃O₃ [M + H⁺] 402.2756, found 402.2761.

3.5. 2-tert-Butoxycarbonyl-6-benzyl-N-(cyclopropylmethyl)-2,6-diazaspiro[3.4]octane-8-carboxamide (**4c**)

This compound was synthesized according to general procedure A from **3** (0.5 g, 1.33 mmol). Yield 0.38 g (72%), colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.40–7.14 (m, 5H), 3.91 (d, *J* = 9.2 Hz, 2H), 3.81 (d, *J* = 8.5 Hz, 1H), 3.70 (d, *J* = 9.0 Hz, 1H), 3.64 (d, *J* = 1.8 Hz, 1H), 3.14–3.02 (m, 2H), 2.92 (dd, *J* = 9.5, 2.8 Hz, 1H), 2.80 (dd, *J* = 6.6, 2.9 Hz, 1H), 2.72–2.57 (m, 2H), 1.03–0.77 (m, 1H), 0.57–0.35 (m, 2H), 0.25–0.08 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 172.72, 138.13, 128.48, 128.44, 127.35, 79.52, 65.67, 59.49, 57.17, 54.05, 43.98, 41.98, 28.27, 10.71, 3.39; HRMS (ESI) *m*/*z* calcd for C₂₃H₃₄N₃O₃ [M + H⁺] 400.2600, found 400.2597.

3.6. 2-tert-Butoxycarbonyl-1-(6-benzyl-2,6-diazaspiro[3.4]oct-8-yl)-N,N-dimethylmethanamine (**5d**)

Compound 4d was synthesized according to general procedure A from 3 (1 g, 2.66 mmol) and used in the next step without further purification. A solution of this compound (0.7 g, 1.8 mmol, 1 equiv.) in absolute THF (3 mL) was added to a suspension of LAH (0.07 g, 1.8 mmol, 1 equiv.) in THF (10 mL) at -70 °C. The reaction mixture was stirred at that temperature for 30 min and allowed to reach -5 °C. Then, it was cooled to -30 °C and decomposed by adding water (0.1 mL), 15% aq. NaOH (0.1 mL), and water (0.3 mL). The resulting precipitate was filtered off and the filtrate was concentrated in vacuo. The resulting aldehyde was used immediately in the next step. It was dissolved in CH_2Cl_2 (10 mL) and the solution was treated, on vigorous stirring, with 33% aq. dimethylamine (0.5 mL). Sodium triacetoxyborohydride (0.96 g, 4.5 mmol, 2.5 equiv.) was added in portions and the resulting mixture was stirred overnight. Sat. aq. NaHCO₃ was added and the resulting mixture was washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The residue was purified by column chromatography on silica, eluting with 10% methanol in CH_2Cl_2 to give 0.25 g (72%) of the title compound as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.47–7.06 (m, 5H), 4.07 (d, *J* = 8.8 Hz, 1H), 3.78 (s, 2H), 3.66–3.48 (m, 3H), 3.01–2.91 (m, 1H), 2.89 (d, J = 9.3 Hz, 1H), 2.55 (d, J = 9.3 Hz, 1H), 2.45–2.30 (m, 1H), 2.30–2.15 (m, 9H), 1.42 (s, 9H); 13 C NMR (75 MHz, CDCl₃) δ 156.38, 138.61, 128.65, 128.22, 126.97, 79.28, 65.56, 61.35, 60.07, 59.18, 45.67, 41.99, 41.73, 28.33; HRMS (ESI) *m*/*z* calcd for C₂₁H₃₄N₃O₂ [M + H⁺] 360.2651, found 360.2655.

3.7. 2-tert-Butoxycarbonyl-8-(1-benzyl-5-methyl-1H-imidazol-2-yl)-6-methyl-2,6diazaspiro[3.4]octane (**10**)

To a solution of **3** (2 g, 5.3 mmol) in ethanol (25 mL) was added 10% Pd/C (0.25 g) and the mixture was hydrogenated in an autoclave at a start pressure of 100 atm and r. t. for 12 h. The reaction mixture was filtered through a plug of Celite and the filtrate was concentrated in vacuo. The yield of compound 7 was 1.6 g (quantitative, assuming analytical purity), obtained as a colorless oil. It was used directly in the next step without further purification.

To a vigorously stirred solution of 7 (1.6 g, 5.3 mmol, 1 equiv.) in CH_2Cl_2 (25 mL), 37% aq. formaldehyde solution (1 mL) was added followed by portion-wise addition of sodium triacetoxyborohydride (2.8 g, 13.25 mmol, 2.5 equiv.). The resulting mixture was stirred overnight. Sat. aq. NaHCO₃ (10 mL) was added, whereupon the mixture was washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The resulting product **8** (1.2 g, 72%, assuming analytical purity) was used in the next step without further purification.

To a solution of **8** (1 g, 3.35 mmol, 1.0 equiv.) in 1,4-dioxane (10 mL), aq. solution of LiOH·H₂O (1 mL, 0.175 g, 4.2 mmol, 1.25 equiv.) was added and the resulting mixture was stirred for 12 h. Thereupon HOBt (0.56 g, 4.2 mmol, 1.25 equiv.), EDC·HCl (0.81 g, 4.2 mmol, 1.25 equiv.), and propargylamine (0.29 g, 4.7 mmol, 1.4 equiv.) were added. The resulting mixture was stirred for 12 h. The solvent was removed in vacuo and the residue was dissolved in ethyl acetate (50 mL). The solution was washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo to give 0.68 g (66%, assuming analytical purity) of product **9** as a colorless oil, which was used in the next step without further purification.

To a solution of compound **9** (0.3 g, 0.97 mmol, 1 equiv.) in toluene (10 mL), benzylamine (0.13 g, 0.24 mmol, 1.25 equiv.) was added followed by $Zn(OTf)_2$ (0.088 g, 1.21 mmol, 0.25 equiv.). The mixture was heated under reflux with a Dean-Stark trap over 8 h. Upon cooling to r. t., the reaction mixture was washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The residue was purified by column chromatography on silica gel using 10% methanol in CH₂Cl₂ to give the title compound (0.29 g, 75%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.35–7.19 (m, 3H), 6.96–6.86 (m, 2H), 6.83 (s, 1H), 5.19–4.93 (m, 2H), 3.77 (d, *J* = 8.6 Hz, 1H), 3.60 (d, *J* = 9.5 Hz, 1H), 3.47–3.29 (m, 3H), 3.11 (d, *J* = 9.3 Hz, 1H), 3.05 (t, *J* = 8.6 Hz, 1H), 2.33 (s, 3H), 2.15 (d, *J* = 0.7 Hz, 3H), 1.39 (s, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 155.97, 147.40, 136.58, 129.02, 127.85, 127.40, 125.84, 125.47, 79.24, 68.07, 62.57, 46.39, 44.59, 44.36, 41.55, 28.31, 9.70; HRMS (ESI) *m/z* calcd for C₂₃H₃₃N₄O₂ [M + H⁺] 397.2603, found 397.2598.

3.8. 2-tert-Butoxycarbonyl-6-methyl-8-(5-methyloxazol-2-yl)-2,6-diazaspiro[3.4]octane (11)

To a solution of compound **9** (0.3 g, 0.97 mmol, 1 equiv.) in toluene (10 mL), Zn(OTf)₂ (0.088 g, 1.21 mmol, 0.25 equiv.) was added and the mixture was heated under reflux with a Dean-Stark trap over 16 h. Upon cooling to r. t., the mixture was washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The residue was purified by column chromatography on silica gel using 10% methanol in CH₂Cl₂ to give the title compound (0.14 g, 47%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 6.63 (d, *J* = 1.2 Hz, 1H), 4.00 (d, *J* = 8.6 Hz, 1H), 3.89 (d, *J* = 8.6 Hz, 1H), 3.60 (d, *J* = 9.2 Hz, 1H), 3.51 (t, *J* = 8.0 Hz, 1H), 3.41 (d, *J* = 9.2 Hz, 1H), 3.18–3.10 (m, 1H), 3.06 (d, *J* = 9.4 Hz, 1H), 2.82–2.70 (m, 2H), 2.39 (s, 3H), 2.27 (s, 3H), 1.40 (s, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 162.33, 156.09, 148.93, 122.69, 79.38, 67.35, 59.58, 46.23, 43.93, 41.86, 28.26, 10.79; HRMS (ESI) *m*/z calcd for C₁₆H₂₆N₃O₃ [M + H⁺] 308.1974, found 308.1980.

3.9. 2-tert-Butyl 6-benzyl-8-(4-methyl-4H-1,2,4-triazol-3-yl)-2,6-diazaspiro[3.4]octane-2-carboxylate (15)

To a solution of **3** (3 g, 8 mmol) in ethanol (25 mL), N_2H_4 (64% aq. solution, 1 mL) was added. The reaction mixture was heated under reflux for 8 h. Upon cooling to r. t., the mixture was concentrated in vacuo to give compound **14** (2.8 g, quant., assuming analytical purity), which was used in the next step without further purification.

To a solution of 14 (2 g, 5.5 mmol, 1 equiv.) in ethanol (25 mL), CH_3NCS (0.5 g, 6.8 mmol, 1.25 equiv.) was added dropwise and the resulting mixture was heated under reflux for 2 h. Sat. aq. K_2CO_3 (5 mL) was added and refluxing continued for 8 h. The reaction mixture was concentrated in vacuo, the residue was dissolved in water (25 mL), and the solution was acidified to pH 7 with 5% aq. HCl. The resulting precipitate was filtered off and dissolved in ethanol (25 mL). A suspension of freshly prepared Raney nickel in a minimum amount of ethanol was added and the mixture was heated under reflux with vigorous stirring for 12 h. Upon cooling to r. t., the reaction mixture was filtered through a plug of Celite and the filtrate was concentrated in vacuo. The residue was purified by column chromatography on silica gel using 10% methanol in CH_2Cl_2 to give the title compound (1.22 g, 58%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 8.08 (s, 1H), 7.39–7.13 (m, 5H), 4.01–3.84 (m, 2H), 3.74–3.59 (m, 6H), 3.53 (t, J = 8.0 Hz, 1H), 3.35 (d, J = 9.5 Hz, 1H), 3.28 (t, J = 8.8 Hz, 1H), 3.12 (d, J = 9.4 Hz, 1H), 2.97–2.86 (m, 1H), 2.80 (d, J = 9.4 Hz, 1H), 1.37 (s, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 156.25, 154.24, 144.60, 138.35, 128.65, 128.34, 127.19, 79.74, 64.86, 59.57, 58.88, 43.59, 42.08, 30.78, 28.23; HRMS (ESI) m/z calcd for C₂₁H₃₀N₅O₂ [M + H⁺] 384.2399, found 384.2405.

3.10. 2-tert-Butoxycarbonyl-6-(methylsulfonyl)-8-(4-methyl-4H-1,2,4-triazol-3-yl)-2,6diazaspiro[3.4]octane (**16**)

Compound **15** was hydrogenated as described for the preparation of compound **7** to give 0.76 g of a colorless oil. To a solution of this oil (0.35 g, 1.1 mmol, 1.0 equiv.) in CH₂Cl₂ (10 mL), Et₃N (0.14 g, 1.37 mmol, 1.25 equiv.) was added dropwise. The mixture was cooled to 0 °C and treated with MsCl (0.16 g, 1.37 mmol, 1.25 equiv.). After stirring the mixture overnight, it was washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The residue was purified by column chromatography on silica gel using 10% methanol in CH₂Cl₂ to give the title compound (0.22 g, 53%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 8.11 (s, 1H), 3.04–3.95 (m, 2H), 3.91–3.75 (m, 4H), 3.64–3.52 (m, 2H), 1.40 (s, 9H).; ¹³C NMR (75 MHz, CDCl₃) δ 156.03, 152.70, 144.60, 80.13, 61.23, 55.73, 53.35, 51.38, 43.74, 41.23, 35.65, 30.95, 28.20; HRMS (ESI) *m/z* calcd for C₁₅H₂₆N₅O₄S [M + H⁺] 372.1705, found 372.1701.

3.11. 2-tert-Butoxycarbonyl-6-benzyl-8-(5-methyl-4H-1,2,4-triazol-3-yl)-2,6-diazaspiro[3.4]octane (**19**)

To a solution of acetamidine hydrochloride (0.16 g, 1.73 mmol, 1.25 equiv.) in absolute methanol (10 mL), MeONa (0.093 g, 1.73 mmol, 1.25 equiv.) was added. The mixture was stirred for 30 min, whereupon crude **14** (0.5 g, 1.39 mmol, 1.0 equiv.) was added and the stirring continued overnight. The solvent was removed in vacuo and the residue was heated at 170 °C under an argon stream for 15 min. The residue was dissolved in ethyl acetate (25 mL), washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The residue was purified by column chromatography on silica gel using 10% methanol in CH₂Cl₂ to give the title compound (0.26 g, 48%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.39–7.13 (m, 5H), 3.92 (dd, *J* = 37.4, 8.6 Hz, 2H), 3.70 (q, *J* = 12.8 Hz, 2H), 3.59–3.41 (m, 3H), 3.20–2.99 (m, 2H), 2.91 (t, *J* = 8.0 Hz, 2H), 2.38 (s, 3H), 1.37 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 156.29, 138.23, 128.69, 128.42, 127.27, 79.68, 64.71, 59.87, 53.40, 45.16, 43.19, 28.30, 8.08, -0.05; HRMS (ESI) *m/z* calcd for C₂₁H₃₀N₅O₂ [M + H⁺] 384.2399, found 384.2404.

3.12. 2-tert-Butoxycarbonyl-6-benzyl-8-(3-cyclopropyl-1H-1,2,4-triazol-5-yl)-2,6-diazaspiro[3.4]octane (**20**)

Synthesized analogously to **19**. Yield 0.23 g (42%), colorless oil. ¹H NMR 1H NMR (300 MHz, CDCl3) δ 7.39–7.16 (m, 5H), 3.91 (dd, *J* = 41.5, 8.6 Hz, 2H), 3.69 (q, *J* = 12.8 Hz, 2H), 3.57–3.41 (m, 3H), 2.04–1.86 (m, 1H), 1.38 (s, 9H), 1.00–0.94 (m, 4H).; ¹³C NMR (75 MHz, DMSO-*d*₆) δ 156.29, 138.23, 128.69, 128.42, 127.27, 79.68, 64.71, 59.87, 53.40, 45.16, 43.19, 28.30, 8.08, -0.05; HRMS (ESI) *m*/*z* calcd for C₂₃H₃₂N₅O₂ [M + H⁺] 410.2556, found 410.2561.

3.13. 2-tert-Butoxycarbonyl-6-benzyl-8-(3-methyl-1,2,4-oxadiazol-5-yl)-2,6diazaspiro[3.4]octane (**23**)

To a solution of **3** (1 g, 2.29 mmol, 1 equiv.) in 1,4-dioxane, an aqueous solution of LiOH·H₂O (1 mL, 0.175 g, 4.2 mmol, 1.25 equiv.) was added and the mixture was stirred for 12 h. HOBt (0.56 g, 4.2 mmol, 1.25 equiv.), EDC·HCl (0.81 g, 4.2 mmol, 1.25 equiv.), and acetamidoxime (0.35 g, 4.7 mmol, 1.4 equiv.) were added and the resulting mixture was stirred for 12 h. The solvent was removed in vacuo and the residue was dissolved in ethyl acetate (50 mL). The solution was washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The residue was dissolved in toluene (25 mL), TBAF (100 mg) was added, and the resulting mixture was heated under reflux with a Dean-Stark trap for 6 h. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography on silica gel using 10% methanol in CH₂Cl₂ to give the title compound (0.48 g, 41%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.41–7.17 (m, 5H), 3.94 (dd, *J* = 33.0, 8.6 Hz, 2H), 3.78–3.59 (m, 4H), 3.49 (d, *J* = 9.3 Hz, 1H), 3.33–3.21 (m, 1H), 3.09 (d, J = 9.4 Hz, 1H), 2.90–2.72 (m, 2H), 2.40 (s, 3H), 1.39 (s, 9H).; ¹³C NMR (75 MHz, DMSO-*d*₆) δ 178.85, 167.17, 155.96, 138.18, 128.53, 128.37, 127.24, 79.69, 59.41, 57.47, 43.93, 43.22, 28.26, 11.60, -0.05; HRMS (ESI) m/z calcd for $C_{21}H_{29}N_4O_3$ [M + H⁺] 385.2239, found 385.2244.

3.14. 2-tert-Butoxycarbonyl-6-(methylsulfonyl)-2,6-diazaspiro[3.4]octane-8-carboxylic acid (25)

To a solution of crude 7 (0.31 g, 1.1 mmol, 1 equiv.) in CH₂Cl₂ (10 mL), Et₃N (0.14 g, 1.37 mmol, 1.25 equiv.) was added dropwise. The mixture was cooled to 0 °C, treated with dropwise addition of MsCl (0.16 g, 1.37 mmol, 1.25 equiv.), and stirred overnight. It was then washed with 10% aq. K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The residue was dissolved in methanol (10 mL) and treated with dropwise addition of 25% aq. KOH (1 mL). The mixture was stirred for 1 h, concentrated in vacuo, and the residue was dissolved in water. The solution was acidified to pH 6 with 5% aq. HCl. The resulting precipitate was filtered off and dried over NaOH pellets to give the title compound. Yield 0.76 g (100%), white solid, m.p. 102–103 °C. ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.94 (s, 1H), 3.99–3.68 (m, 4H), 3.46–3.29 (m, 3H), 3.22 (dd, *J* = 7.4, 6.0 Hz, 1H), 2.91 (s, 3H), 1.36 (s, 9H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 172.53, 155.92, 79.12, 56.42, 49.37, 48.78, 41.60, 34.30, 28.45; HRMS (ESI) *m/z* calcd for C₁₃H₂₃N₂O₆S [M + H⁺] 335.1276, found 335.1280.

3.15. 2-tert-Butoxycarbonyl-8-(3-cyclopropyl-1,2,4-oxadiazol-5-yl)-6-(methylsulfonyl)-2,6diazaspiro[3.4]octane (**26**)

To a solution of **25** (0.25 g, 0.75 mmol, 1 equiv.) in CH₂Cl₂ (10 mL), CDI (0.15 g, 0.94 mmol, 1.25 equiv.) was added and the mixture was stirred for 1 h, whereupon *N*[']-hydroxycyclopropanecarboximidamide (0.094 g, 0.94 mmol, 1.25 equiv.) was added and the stirring was continued overnight. The reaction mixture was washed with 1% aq. HCl (2 × 15 mL) and concentrated in vacuo. The residue was dissolved in toluene (25 mL), TBAF (100 mg) was added, and the mixture was heated under reflux with a Dean-Stark trap for 6 h. The reaction mixture was evaporated to dryness. The residue was purified by column chromatography on silica gel using 10% methanol in CH₂Cl₂ to give the title compound (0.19 g, 63%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 4.02–3.91 (m, 2H), 3.85 (t, *J* = 7.6 Hz, 1H), 3.81–3.65 (m, 5H), 3.56 (d, *J* = 9.9 Hz, 1H), 2.08 (dt, *J* = 8.3, 4.9 Hz, 1H), 1.41 (s, 9H), 1.13–0.88 (m, 4H); ¹³C NMR (75 MHz, CDCl₃) δ 156.03, 152.70, 144.60,

80.13, 55.73, 51.38, 43.74, 41.23, 35.65, 30.95, 28.20; HRMS (ESI) m/z calcd for C₁₇H₂₇N₄O₅S [M + H⁺] 399.1702, found 399.1697.

3.16. 6-Benzyl-N-methyl-2-(5-nitro-2-furoyl)-2,6-diazaspiro[3.4]octane-8-carboxamide (5a)

General Procedure B. For the preparation of compounds **5a–c**, **6d**, **12–3**, **17–18**, **21–22**, **24** and **27**. To a solution of 5-nitro-2-furoic acid (75 mg, 0.47 mmol) in DMF (3 mL), CDI (97 mg, 0.6 mmol) was added at 0 °C and the solution was stirred for 1 h.

To a solution of 4a (0.22 g, 0.6 mmol) in CH₂Cl₂ (5 mL) at 0 °C, trifluoroacetic acid (1 mL) was added and the mixture was stirred for 1 h. The solution was concentrated in vacuo while keeping the bath temperature under 30 °C. The residue was dissolved in DMF (3 mL), triethylamine (0.19 g, 1.9 mmol) was added dropwise, and after 30 min stirring, the mixture was added to the solution of 5-nitro-2-furoic acid imidazolide prepared as described above. The reaction mixture was stirred at r. t. overnight, poured into water (25 mL), and extracted with ethyl acetate (3 \times 20 mL). The combined organic phases were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The residue was purified by column chromatography on silica gel using 10% methanol in CH_2Cl_2 to give the title compound. Yield 133 mg (56%), white solid, m.p. 132–135 °C. ¹H NMR (300 MHz, DMSO-*d*₆) mixture of rotamers δ 8.06 (s, 1H), 7.75 (dd, *J* = 8.8, 3.9 Hz, 1H), 7.40–7.15 (m, 7H), 4.45 (dd, J = 27.0, 9.3 Hz, 1H), 4.31 (s, 1H), 4.10–3.74 (m, 3H), 3.67–3.51 (m, 2H), 3.07–2.82 (m, 4H), 2.75–2.52 (m, 7H), 1.37–1.14 (m, 2H), 0.96–0.73 (m, 1H); ¹³C NMR (75 MHz, DMSO-*d*₆) mixture of rotamers δ 172.28, 156.47, 151.81, 147.95, 147.82, 139.15, 128.88, 128.85, 128.59, 127.27, 117.35, 117.31, 113.49, 113.43, 65.50, 64.72, 61.17, 59.29, 59.21, 56.69, 55.20, 51.47, 51.33, 43.53, 25.91, 25.88; HRMS (ESI) m/z calcd for $C_{20}H_{23}N_4O_5$ [M + H⁺] 399.1668, found 399.1673.

3.17. 6-Benzyl-N-isobutyl-2-(5-nitro-2-furoyl)-2,6-diazaspiro[3.4]octane-8-carboxamide (5b)

This compound was synthesized according to general procedure B from **4b** (0.24 g, 0.6 mmol). Yield 148 mg (56%), white solid, m.p. 166–165 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 8.13 (dd, J = 10.8, 5.4 Hz, 1H), 7.73 (dt, J = 16.3, 8.2 Hz, 1H), 7.42–7.09 (m, 7H), 4.47 (dd, J = 33.9, 9.4 Hz, 1H), 4.37–4.24 (m, 1H), 4.12 (d, J = 10.3 Hz, 1H), 4.00–3.75 (m, 2H), 3.64–3.50 (m, 2H), 3.08–2.68 (m, 6H), 2.67–2.53 (m, 2H), 1.62 (dd, J = 11.3, 6.7 Hz, 1H), 0.89–0.61 (m, 7H); ¹³C NMR (75 MHz, DMSO- d_6) mixture of rotamers, δ 196.40, 193.60, 174.10, 173.50, 171.66, 168.54, 161.80, 159.89, 156.37, 151.78, 147.97, 147.91, 146.43, 146.33, 145.57, 139.77, 139.14, 134.96, 133.64, 129.86, 129.54, 128.97, 128.85, 128.59, 128.45, 128.31, 127.87, 127.28, 121.61, 121.53, 121.50, 121.45, 120.72, 118.14, 117.31, 113.49, 113.45, 105.49, 65.02, 64.74, 60.90, 59.40, 59.30, 59.23, 56.64, 56.51, 55.38, 55.26, 51.51, 51.44, 46.53, 43.47, 28.38, 28.34, 20.44, 20.35; HRMS (ESI) m/z calcd for C₂₃H₂₉N₄O₅ [M + H⁺] 441.2137, found 441.2134.

3.18. 6-Benzyl-N-(cyclopropylmethyl)-2-(5-nitro-2-furoyl)-2,6-diazaspiro[3.4]octane-8-carboxamide (**5c**)

This compound was synthesized according to general procedure B from **4c** (0.185 g, 0.6 mmol). Yield 126 mg (48%), white solid, m.p. 180–181 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 8.42–8.08 (m, 1H), 7.76 (dd, J = 8.9, 3.9 Hz, 1H), 7.45–7.11 (m, 8H), 4.62–4.26 (m, 2H), 4.14 (d, J = 10.4 Hz, 1H), 3.94 (d, J = 9.2 Hz, 1H), 3.81 (d, J = 11.1 Hz, 1H), 3.69–3.47 (m, 3H), 3.14–2.77 (m, 6H), 2.62 (dd, J = 16.4, 8.9 Hz, 2H), 1.00–0.70 (m, 1H), 0.33 (dd, J = 18.6, 7.1 Hz, 2H), 0.10 (dd, J = 9.1, 4.1 Hz, 2H).; ¹³C NMR (75 MHz, DMSO- d_6) mixture of rotamers, δ 171.59, 156.38, 151.79, 147.97, 147.91, 139.16, 128.88, 128.85, 128.74, 128.60, 127.27, 117.32, 113.46, 65.22, 64.65, 61.03, 59.30, 59.22, 56.51, 56.35, 55.26, 51.34, 43.54, 43.34, 43.30, 11.33, 11.30, 3.66, 3.55, 3.46; HRMS (ESI) m/z calcd for C₂₃H₂₇N₄O₅ [M + H⁺] 439.1981, found 439.1986.

3.19. 1-[6-Benzyl-2-(5-nitro-2-furoyl)-2,6-diazaspiro[3.4]oct-8-yl]-N,N-dimethylmethanamine (6d)

This compound was synthesized according to general procedure B from **5d** (0.215 g, 0.6 mmol). Yield 91 mg (38%), white solid, m.p. 154–155 °C. ¹H NMR (300 MHz, DMSO-*d*₆) mixture of rotamers δ 7.77 (d, *J* = 1.9 Hz, 1H), 7.76 (d, *J* = 1.9 Hz, 1H), 7.43–7.15 (m, 14H), 4.68 (d, *J* = 9.5 Hz, 1H), 4.39 (s, 2H), 4.31 (d, *J* = 10.4 Hz, 1H), 4.17 (d, *J* = 9.6 Hz, 1H), 3.94 (q, *J* = 10.1 Hz, 2H), 3.70 (d, *J* = 10.4 Hz, 1H), 3.63–3.48 (m, 5H), 2.98–2.55 (m, 8H), 2.44–2.25 (m, 5H), 2.10 (t, *J* = 10.9 Hz, 17H); ¹³C NMR (75 MHz, DMSO-*d*₆) mixture of rotamers, δ 156.64, 156.48, 151.80, 148.01, 147.96, 139.28, 135.52, 128.84, 128.57, 127.21, 117.20, 113.45, 65.42, 65.36, 64.58, 61.25, 61.13, 60.33, 59.67, 58.65, 58.48, 58.27, 54.17, 45.74, 45.69, 43.23, 43.14, 41.77, 41.47.; HRMS (ESI) *m*/*z* calcd for C₂₁H₂₇N₄O₅ [M + H⁺] 399.2032, found 399.2028.

3.20. 8-(1-Benzyl-5-methyl-1H-imidazol-2-yl)-6-methyl-2-(5-nitro-2-furoyl)-2,6-diazaspiro[3.4]octane (12)

This compound was synthesized according to general procedure B from **10** (0.237 g, 0.6 mmol). Yield 74 mg (28%), white solid, m.p. 165–167 °C. ¹H NMR (300 MHz, DMSO-*d*₆) mixture of rotamers δ 7.74 (dd, *J* = 13.4, 3.8 Hz, 1H), 7.28–7.06 (m, 3H), 7.02–6.81 (m, 3H), 6.73 (d, *J* = 7.7 Hz, 1H), 5.37–5.06 (m, 2H), 4.24 (d, *J* = 9.5 Hz, 1H), 4.15 (d, *J* = 10.2 Hz, 1H), 3.94–3.53 (m, 4H), 3.38 (d, *J* = 19.3 Hz, 4H), 3.17–2.94 (m, 2H), 2.76–2.54 (m, 2H), 2.46–2.36 (m, 1H), 2.23 (s, 3H), 2.17–2.00 (m, 4H); ¹³C NMR (75 MHz, DMSO-*d*₆) mixture of rotamers, δ 155.99, 151.74, 151.57, 148.01, 147.85, 147.65, 138.48, 138.23, 128.98, 128.91, 127.69, 127.51, 127.38, 127.19, 126.15, 125.97, 125.77, 125.72, 117.17, 117.12, 113.46, 68.02, 67.52, 65.66, 63.11, 62.97, 62.02, 58.47, 55.04, 45.96, 45.87, 45.24, 45.10, 43.51, 43.40, 41.95, 41.87.; HRMS (ESI) *m*/*z* calcd for C₂₃H₂₆N₅O₄ [M + H⁺] 436.1984, found 436.1988.

3.21. 6-Methyl-8-(5-methyl-1,3-oxazol-2-yl)-2-(5-nitro-2-furoyl)-2,6-diazaspiro[3.4]octane (13)

This compound was synthesized according to general procedure B from **11** (0.184 g, 0.6 mmol). Yield 99 mg (48%), white solid, m.p. 175–176 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 7.77 (d, *J* = 3.9 Hz, 1H), 7.72 (d, *J* = 3.9 Hz, 1H), 7.32 (d, *J* = 3.9 Hz, 1H), 7.27 (d, *J* = 3.9 Hz, 1H), 7.02 (s, 1H), 6.80–6.73 (m, 2H), 4.57 (q, *J* = 9.6 Hz, 2H), 4.31 (d, *J* = 10.1 Hz, 1H), 4.17 (d, *J* = 10.4 Hz, 1H), 4.05 (d, *J* = 10.4 Hz, 1H), 3.97 (d, *J* = 10.0 Hz, 1H), 3.84–3.64 (m, 3H), 3.55–3.44 (m, 4H), 3.12–2.94 (m, 4H), 2.76 (ddd, *J* = 17.3, 7.5, 4.9 Hz, 4H), 2.30 (s, 5H), 2.25 (s, 3H), 2.22 (s, 2H); ¹³C NMR (75 MHz, DMSO- d_6) mixture of rotamers, δ 162.41, 162.31, 156.55, 156.43, 151.85, 149.25, 149.19, 147.67, 147.61, 122.93, 117.41, 113.42, 113.36, 66.17, 64.16, 59.94, 59.25, 58.90, 58.79, 55.23, 45.46, 45.41, 44.98, 41.89, 10.82.; HRMS (ESI) *m*/z calcd for C₁₆H₁₉N₄O₅ [M + H⁺] 347.1355, found 347.1360.

3.22. 6-(Methylsulfonyl)-8-(4-methyl-4H-1,2,4-triazol-3-yl)-2-(5-nitro-2-furoyl)-2,6diazaspiro[3.4]octane (**17**)

This compound was synthesized according to general procedure B from **16** (0.22 g, 0.6 mmol). Yield 138 mg (56%), white solid, m.p. 180–182 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 8.44 (s, 1H), 7.78 (d, *J* = 3.9 Hz, 1H), 7.73 (d, *J* = 3.9 Hz, 1H), 7.30 (t, *J* = 4.1 Hz, 1H), 4.57 (q, *J* = 9.5 Hz, 2H), 4.21–3.84 (m, 5H), 3.84–3.58 (m, 6H), 3.56–3.37 (m, 1H), 2.85 (d, *J* = 9.7 Hz, 3H); ¹³C NMR (75 MHz, DMSO- d_6) mixture of rotamers, δ 156.67, 152.31, 151.83, 147.80, 147.70, 145.65, 117.39, 113.50, 113.45, 62.50, 59.36, 57.62, 56.55, 55.27, 53.97, 51.20, 51.05, 44.12, 43.99, 34.59, 34.52, 30.92.; HRMS (ESI) *m*/*z* calcd for C₁₅H₁₉N₆O₆S [M + H⁺] 411.1086, found 411.1091.

3.23. 6-Benzyl-8-(4-methyl-4H-1,2,4-triazol-3-yl)-2-(5-nitro-2-furoyl)-2,6diazaspiro[3.4]octane (**18**)

This compound was synthesized according to general procedure B from **15** (0.23 g, 0.6 mmol). Yield 157 mg (62%), white solid, m.p. 152–153 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 8.65–8.20 (m, 1H), 7.84–7.59 (m, 2H), 7.57–7.16 (m, 5H), 7.02 (s, 1H), 4.77–3.96 (m, 4H), 3.94–3.06 (m, 13H), 2.85 (dd, *J* = 21.6, 9.0 Hz, 1H), 2.68 (dd, *J* = 16.7, 8.9 Hz, 1H); ¹³C NMR (75 MHz, DMSO- d_6) mixture of rotamers, δ 156.60, 154.43, 154.32,

13 of 15

151.79, 147.75, 145.30, 139.09, 135.51, 129.86, 129.54, 128.91, 128.86, 128.62, 127.32, 122.04, 117.32, 113.43, 65.48, 64.63, 64.49, 61.43, 59.33, 58.92, 58.72, 54.82, 44.51, 30.77; HRMS (ESI) m/z calcd for C₂₁H₂₃N₆O₄ [M + H⁺] 423.1780, found 423.1777.

3.24. 6-Benzyl-8-(5-methyl-4H-1,2,4-triazol-3-yl)-2-(5-nitro-2-furoyl)-2,6diazaspiro[3.4]octane (**21**)

This compound was synthesized according to general procedure B from **19** (0.23 g, 0.6 mmol). Yield 108 mg (43%), white solid, m.p. 168–170 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 8.57–8.26 (m, 1H), 7.87–7.60 (m, 2H), 7.52–7.13 (m, 6H), 7.02 (s, 1H), 4.82–3.93 (m, 4H), 3.89–3.52 (m, 7H), 3.27–3.04 (m, 2H), 2.99–2.55 (m, 3H); ¹³C NMR (75 MHz, DMSO- d_6) mixture of rotamers δ 158.23, 156.60, 154.43, 154.32, 151.79, 147.91, 147.75, 145.30, 139.09, 135.51, 129.86, 129.54, 128.91, 128.86, 128.62, 127.82, 127.32, 122.04, 117.32, 113.43, 66.85, 65.48, 64.63, 64.49, 61.43, 59.33, 58.92, 58.72, 54.82, 45.93, 44.51, 43.31, 30.77.; HRMS (ESI) *m/z* calcd for C₂₁H₂₃N₆O₄ [M + H⁺] 423.1780, found 423.1784.

3.25. 6-Benzyl-8-(5-cyclopropyl-4H-1,2,4-triazol-3-yl)-2-(5-nitro-2-furoyl)-2,6diazaspiro[3.4]octane (**22**)

This compound was synthesized according to general procedure B from **20** (0.24 g, 0.6 mmol). Yield 102 mg (38%), white solid, m.p. 155–157 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 13.39 (s, 1H), 7.82–7.58 (m, 1H), 7.41–7.10 (m, 5H), 4.59–4.39 (m, 1H), 4.26–3.85 (m, 2H), 3.73–3.57 (m, 2H), 3.14–2.60 (m, 3H), 2.06–1.69 (m, 1H), 1.06–0.49 (m, 4H); 13C NMR (75 MHz, DMSO- d_6) mixture of rotamers δ 193.60, 167.18, 161.07, 156.43, 156.30, 151.73, 147.85, 147.80, 139.26, 135.51, 134.96, 129.86, 129.54, 128.87, 128.60, 127.26, 121.70, 121.63, 121.57, 121.54, 121.50, 121.48, 118.46, 117.24, 113.44, 113.38, 64.18, 64.10, 63.66, 59.53, 59.38, 57.37, 57.29, 55.38, 55.26, 44.29, 44.20, 9.48, 8.21, 7.60; HRMS (ESI) *m/z* calcd for C₂₃H₂₅N₆O₄ [M + H⁺] 449.1937, found 449.1942.

3.26. 6-Benzyl-8-(3-methyl-1,2,4-oxadiazol-5-yl)-2-(5-nitro-2-furoyl)-2,6diazaspiro[3.4]octane (**24**)

This compound was synthesized according to general procedure B from **23** (0.245 g, 0.6 mmol). Yield 121 mg (48%), white solid, m.p. 180–182 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 7.84–7.63 (m, 1H), 7.41–7.09 (m, 5H), 4.75–4.42 (m, 1H), 4.27 (dd, J = 36.8, 10.2 Hz, 1H), 4.10–3.93 (m, 2H), 3.82 (t, J = 11.6 Hz, 0H), 3.74–3.50 (m, 2H), 3.10 (dt, J = 26.4, 20.0, 9.4 Hz, 2H), 2.92–2.75 (m, 2H), 2.31 (d, J = 8.0 Hz, 3H); ¹³C NMR (75 MHz, DMSO- d_6) mixture of rotamers δ 179.23, 179.17, 167.29, 167.22, 162.72, 156.61, 151.85, 147.72, 147.58, 138.82, 129.86, 129.54, 129.00, 128.87, 128.66, 127.39, 121.59, 117.42, 113.44, 63.90, 63.78, 59.79, 59.04, 56.78, 56.66, 55.25, 55.16, 44.20, 43.12, 36.17, 31.17, 11.52.; HRMS (ESI) m/z calcd for C₂₁H₂₂N₅O₅ [M + H⁺] 424.1620, found 424.1616.

3.27. 8-(3-Cyclopropyl-1,2,4-oxadiazol-5-yl)-6-(methylsulfonyl)-2-(5-nitro-2-furoyl)-2,6-diazaspiro[3.4]octane (**27**)

This compound was synthesized according to general procedure B from **26** (0.24 g, 0.6 mmol). Yield 165 mg (63%), white solid, m.p. 171–173 °C. ¹H NMR (300 MHz, DMSO- d_6) mixture of rotamers δ 7.77 (dd, J = 9.8, 3.9 Hz, 1H), 7.31 (dd, J = 8.0, 3.9 Hz, 1H), 4.65 (q, J = 9.9 Hz, 1H), 4.45 (dd, J = 25.5, 10.0 Hz, 1H), 4.26–4.07 (m, 2H), 3.97 (dt, J = 35.2, 9.1 Hz, 1H), 3.82–3.52 (m, 4H), 2.97 (d, J = 1.7 Hz, 3H), 2.18–2.02 (m, 1H), 1.07–0.68 (m, 4H); ¹³C NMR (75 MHz, DMSO- d_6) mixture of rotamers δ 177.00, 176.87, 172.19, 156.67, 151.86, 147.63, 147.52, 122.80, 122.67, 122.56, 117.49, 113.48, 60.70, 57.56, 55.56, 55.26, 54.21, 49.35, 49.15, 43.77, 43.71, 42.72, 42.54, 34.65, 34.59, 7.89, 6.62; HRMS (ESI) *m/z* calcd for C₁₇H₂₀N₅O₇S [M + H⁺] 438.1083, found 438.1088.

3.28. Method to Evaluate Antitubercular Activity of Compounds 5a–c, 6d, 12–13, 17–18, 21–22, 24, and 27

Mycobacterium tuberculosis H₃₇Rv strain (originated from the Institute of Hygiene and Epidemiology in Prague, 1976) was obtained on 7 August 2013 from the Federal Scientific

14 of 15

Center for Expertise of Medical Products (RF Ministry of Health Care). The lyophilized strain was seeded in Löwenstein–Jensen growth medium. After 3 weeks, the culture was suspended in physiological solution containing glycerol (15%), transferred into cryotubes, and kept at -80 °C. Three weeks prior to the experiment, the culture was brought to ambient temperature and re-seeded into Löwenstein–Jensen growth medium. Thus, the 2nd generation of the original *M. tuberculosis* culture was used in the present study.

The minimal inhibitory concentration (MIC) of each compound was determined using the REMA (resazurin microtiter plate assay) [13]. A 3-week M. tuberculosis culture was transferred into a dry, sterile tube containing 8–9 3 mm glass beads. The tube was placed on a Vortex shaker for 30-40 s and then 5 mL Middlebrook 7H9 Broth (Becton Dickinson, catalogue No. 271310) was introduced. The turbidity of the bacterial suspension was adjusted to 1.0 McFarland units (corresponding to approximately 3×10^8 bacteria/mL) and diluted 20-fold with Middlebrook 7H9 Broth containing OADC enrichment (Becton Dickinson, catalogue No. 245116). The same culture medium was used to prepare the 1:100 *M. tuberculosis* (1% population) control. The stock solution of the compound in DMSO (10 mg/mL) was diluted with Middlebrook 7H9 Broth (containing OADC enrichment) to a concentration of 100 μ g/mL. A 200 μ L aliquot of the solution thus obtained was introduced into the 1st row of a 96-well microtiter plate. This row was used to perform 2-fold serial dilutions using an 8-channel pipette to obtain final concentrations of 0.4, 0.8, 1.6, 3.1, 6.2, 12.5, 25, and 50 μ g/mL of the compound in rows 2–9 (accounting for 100 μ L of bacterial suspension introduced for testing). Similarly, an experiment with 0.004, 0.008, 0.016, 0.031, 0.062, 0.125, 0.25, 0.50 and 1 μ g/mL concentrations of the compounds in rows 1–9 was carried out. Row 10 contained the MTb suspension control while row 11 contained the same culture diluted 10-fold (1% control). Row 12 was used as a blank control for the optical density reading (200 μ L of growth medium). The bacterial suspension (100 μ L) was introduced into each well, except rows 11 (1% population control) and 12 (blank culture medium), to give a total volume of 200 μ L in each well. The plates were incubated at 35 °C for 7 days. At that point, 0.01% aqueous solution (30 μ L) of resazurin (Sigma, product No. R7017) was introduced in each well and the incubation continued for 18 h at 35 °C. The fluorescence reading was performed using a FLUOstar Optima plate reader operating at λ_{ex} = 520 nm and λ_{em} = 590 nm. The bacterial viability was determined by comparing the mean values (\pm SD at *p* = 0.05) of fluorescence in the control wells (row 12, blank and row 11, 1% control) and the wells containing the compound tested. The MIC was determined as the compound concentration at which the fluorescence reached a plateau or was statistically (t criterion) similar to that of the 1% control (see Supplementary Materials).

4. Conclusions

In conclusion, by elaborating a small set of twelve compounds from readily available 2,6-diazaspiro[3.4]octane building block **3** and exploring diverse variants of the molecular periphery, including various azole substituents, we identified a remarkably potent antitubercular lead displaying a minimal inhibitory concentration of 0.016 μ g/mL. This compound will be tested in vitro at a lower concentration range and in vivo to determine its efficacy and safety. The results of these studies will be reported in due course.

Supplementary Materials: The following supporting information can be downloaded at https: //www.mdpi.com/article/10.3390/molecules28062529/s1: copies of the NMR spectra, images of exemplary assay plates.

Author Contributions: Conceptualization, A.L. and M.K.; methodology, A.L. and P.Y.; investigation, K.K., L.V. and M.D.; data curation, A.L. and T.V.; writing—original draft preparation, M.K.; writing—review and editing, A.K., A.L. and M.K.; supervision, A.L.; funding acquisition, M.K. All authors have read and agreed to the published version of the manuscript.

Funding: We gratefully acknowledge the financial support from the Russian Foundation for Basic Research (grant no. 21-53-12001).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data available from the corresponding authors upon reasonable request.

Acknowledgments: We are grateful to the Centre for Chemical Analysis and Materials Research of Saint Petersburg State University Research Park for the high-resolution mass spectrometry data. This work was also performed using the equipment of the Shared Science and Training Center for Collective Use RTU MIREA and supported by the Ministry of Science and Higher Education of the Russian Federation within the framework of agreement No. 075-15-2021-689 dated 1 September 2021.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

- Elsaman, T.; Mohamed, M.S.; Mohamed, M.A. Current development of 5-nitrofuran-2-yl derivatives as antitubercular agents. Bioorg. Chem. 2019, 88, 102969. [CrossRef] [PubMed]
- Stewart, M.L.; Bueno, G.J.; Baliani, A.; Klenke, B.; Brun, R.; Brock, J.M.; Gilbert, I.H.; Barrett, M.B. Trypanocidal activity of melamine-based nitroheterocycles. *Antimicrob. Agents Chemother.* 2004, 48, 1733–1738. [CrossRef] [PubMed]
- Krasavin, M.; Lukin, A.; Vedekhina, T.; Manicheva, O.; Dogonadze, M.; Vinogradova, T.; Zabolotnykh, N.; Rogacheva, E.; Kraeva, L.; Yablonsky, P. Conjugation of a 5-nitrofuran-2-oyl moiety to aminoalkylimidazoles produces non-toxic nitrofurans that are efficacious in vitro and in vivo against multidrug-resistant *Mycobacterium tuberculosis*. *Eur. J. Med. Chem.* 2018, 157, 1115–1126. [CrossRef] [PubMed]
- Krasavin, M.; Lukin, A.; Vedekhina, T.; Manicheva, O.; Dogonadze, M.; Vinogradova, T.; Zabolotnykh, N.; Rogacheva, E.; Kraeva, L.; Sharoyko, V.; et al. Attachment of a 5-nitrofuroyl moiety to spirocyclic piperidines produces non-toxic nitrofurans that are efficacious in vitro against multidrug-resistant *Mycobacterium tuberculosis*. *Eur. J. Med. Chem.* 2019, *166*, 125–135. [CrossRef] [PubMed]
- Ma, Y.; Zhao, S.; Ren, Y.; Cherukupalli, S.; Li, Q.; Woodson, M.E.; Bradley, D.P.; Tavis, J.E.; Liu, X.; Zhan, P. Design, synthesis and evaluation of heteroaryldihydropyrimidine analogues bearing spiro ring as hepatitis B virus capsid protein inhibitors. *Eur. J. Med. Chem.* 2021, 225, 113780. [CrossRef] [PubMed]
- 6. Burrows, F.; Tomkinson, B. Treatment of Hematological Malignancies with Inhibitors of Menin. PCT Int. Appl. WO 2022086986A1, 28 April 2022.
- Akan, D.T.; Howes, J.E.; Sai, J.; Arnold, A.L.; Beesetty, Y.; Phan, J.; Olejniczak, E.T.; Waterson, A.G.; Fesik, S.W. Small Molecule SOS1 Agonists Modulate MAPK and PI3K Signaling via Independent Cellular Responses. ACS Chem. Biol. 2019, 14, 325–331. [CrossRef] [PubMed]
- Reilly, S.W.; Griffin, S.; Taylor, M.; Sahlholm, K.; Weng, C.-C.; Xu, K.; Jacome, D.A.; Luedtke, R.R.; Mach, R.H. Highly Selective Dopamine D3 Receptor Antagonists with Arylated Diazaspiro Alkane Cores. J. Med. Chem. 2017, 60, 9905–9910. [CrossRef] [PubMed]
- 9. Brown, W.D.; Hansen, K.T.H.; Salehi, S.A. Novel Vdac1 Inhibitors. PCT Int. Appl. WO 2022022846A1, 28 April 2022.
- Le Manach, C.; Dam, J.; Woodland, J.G.; Kaur, G.; Khonde, L.P.; Brunschwig, C.; Njoroge, M.; Wicht, K.J.; Horatscheck, A.; Paquet, T.; et al. Identification and Profiling of a Novel Diazaspiro[3.4]octane Chemical Series Active against Multiple Stages of the Human Malaria Parasite Plasmodium falciparum and Optimization Efforts. *J. Med. Chem.* 2021, 64, 2291–2309. [CrossRef] [PubMed]
- 11. Pews-Davtyan, A.; Beller, M. A novel Zn-catalyzed hydroamination of propargylamides: A general synthesis of di- and trisubstituted imidazoles. *Chem. Commun.* 2011, 47, 2152–2154. [CrossRef] [PubMed]
- 12. Safrygin, A.; Dar'in, D.; Lukin, A.; Bakholdina, A.; Sapegin, A.; Krasavin, M. Zn(OTf)2-catalyzed, microwave-promoted synthesis of 2-substituted 5-methyloxazoles from propargylic amides. *Tetrahedron Lett.* **2019**, *60*, 777–779. [CrossRef]
- Martin, A.; Camacho, M.; Portaels, F.; Palomino, C.C. Resazurin microtiter assay plate testing of *Mycobacterium tuberculosis* susceptibility to second line drugs: Rapid simple and inexpensive method. *Antimicrob. Agents Chemother.* 2003, 47, 3616–3619. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.