

Host-Guest Interactions of Zirconium-based Metal-Organic Framework with Ionic Liquid

Mohd. Faridzuan Majid ^{1,2}, Hayyiratul Fatimah Mohd Zaid ^{2,3,*}, Muhammad Fadhlullah Abd Shukur ^{1,2}, Azizan Ahmad ^{4,5} and Khairulazhar Jumbri ^{1,6}

¹ Department of Fundamental and Applied Sciences, Universiti Teknologi PETRONAS, 32610 Seri Iskandar, Perak Darul Ridzuan, Malaysia; mohd._17006281@utp.edu.my (M.F.M.); mfadhlullah.ashukur@utp.edu.my (M.F.A.S.); khairulazhar.jumbri@utp.edu.my (K.J.)

² Centre of Innovative Nanostructures & Nanodevices (COINN), Universiti Teknologi PETRONAS, 32610 Seri Iskandar, Perak Darul Ridzuan, Malaysia

³ Chemical Engineering Department, Universiti Teknologi PETRONAS, 32610 Seri Iskandar, Perak, Malaysia

⁴ Department of Chemical Sciences, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia; mfadhlullah.ashukur@utp.edu.my

⁵ Department of Physics, Faculty of Science and Technology, Airlangga University (Campus C), Mulyorejo Road, Surabaya 60115, Indonesia

⁶ Centre for Research in Ionic Liquids (CORIL), Universiti Teknologi PETRONAS, 32610 Seri Iskandar, Perak Darul Ridzuan, Malaysia

* Correspondence: hayyiratul.mzaid@utp.edu.my; Tel.: +605-3687618




Figure S1. The surface of pristine UiO-66 at 5 μm.




Figure S2. The surface of pristine UiO-66 at 2 μm.




Figure S3. The surface UiO-66@IL at 20 μm.




Figure S4. The surface of UiO-66@IL at 10 μm.