

Cervical Squamous Cell Carcinoma Diagnosis by FTIR Microspectroscopy

Maria M. Félix ¹, Mariana V. Tavares ^{1,2}, Inês P. Santos ¹, Ana L. M. Batista de Carvalho ¹,
Luís A. E. Batista de Carvalho ^{1,*} and Maria Paula M. Marques ^{1,3}

¹ Molecular Physical-Chemistry R&D Unit, Department of Chemistry, University of Coimbra, 3004-535 Coimbra, Portugal; mmfelix@uc.pt (M.M.F.); mariana.vide.tavares@ipoporto.min-saude.pt (M.V.T.); ips@uc.pt (I.P.S.); pmc@ci.uc.pt (M.P.M.M.)

² Gynaecology Department, Portuguese Oncology Institute of Porto, 4200-072 Porto, Portugal

³ Department of Life Sciences, Faculty of Science and Technology, University of Coimbra, 3000-456 Coimbra, Portugal

* Correspondence: labc@ci.uc.pt; Tel.: +351-239-854-462

Supplementary Information

TABLE OF CONTENTS

1. Figures




Figure S1. Principal Component Analysis score of FTIR data for cryopreserved normal vs. SCC cervical tissues; **(A)** PC1 vs. PC2; **(B)** PC1 vs. PC3 and **(C)** PC2 vs. PC3