


Molecules **1999**, *4*, M119

3-Cyano-2-cyanomethylthio-4-methoxymethyl-6-methylpyridine

Yelena Kaigorodova, Vladimir Vasilin and Gennady Krapivin

Department of organic chemistry, Kuban State Technological University, Moskovskaya, 2, Krasnodar, 350072, Russian Federation. Phone +7 8612 559556, E-mail: organics@kubstu.ru

Received: 13 October 1999 / Accepted: 14 October 1999 / Published: 23 November 1999



2-Cyanomethylthiopyridine (**2**) was prepared by the reaction of the pyridinethione **1** with chloroacetonitrile using literature procedures [1, 2].

To a suspension of pyridinethione **1** (1.94 g, 10 mmol) in 20 ml of DMF aqueous solution of potassium hydroxide (10%, 5.6 ml, 10 mmol) and chloroacetonitrile (0.64 ml, 10 mmol) were added. The mixture was kept at room temperature for 3 hours, then diluted with twice the volume of water. The precipitate of product (**2**) was separated, washed with a small volume of water and recrystallized from ethanol. Compound **2** - colourless solid. Yield 2.24g (96%).

M.p.: 105-106°C (ethanol).

¹H NMR ((CD₃)₂CO, 60 MHz): 2.73 (s, 3H, CH₃); 2.77 (s, 3H, CH₂OCH₃); 4.30 (s, 2H, S-CH₂); 4.58 (s, 2H, CH₂OCH₃); 7.28 (s, 1H, H_{Het}).

IR (vaseline oil, cm⁻¹): 2240s, 2210vs, 1575vs, 1540s, 1415m, 1390w, 1340s, 1305w, 1265s, 1230m, 1190s, 1140s, 1010vs, 1005vs, 1030w, 990w, 970m, 960m, 930w, 890s, 870s, 850m, 760w, 725m.

Anal. calc. for C₁₁H₁₁N₃OS (233.291): C 56.63, H 4.75, N 18.01; found: C 56.59, H 4.82, N 17.92.

References

1. Kaigorodova, Ye. A.; Konyushkin, L.D.; Niyazymbetov, M.E.; Kvak, S.N.; Zaplishny, V.N.; Litvinov, V.P. *Russ. Chem. Bull.* **1994**, *43*, 2095.
2. Kaigorodova, Ye. A.; Konyushkin, L.D.; Michailichenko, S.N.; Vasilin, V.K.; Kulnevich, V.G. *Khimiya Geterotsiklicneskikh Soedineny (Chemistry of heterocyclic compounds)* **1996**, *10*, 1432.

Sample availability: available from the authors and MDPI.

©1999 MDPI. All rights reserved. *Molecules* website www.mdpi.org/molecules/