



Review

Aux/IAA Gene Family in Plants: Molecular Structure, Regulation, and Function

Jie Luo ¹, Jing-Jing Zhou ^{1,*} and Jin-Zhi Zhang ^{2,*}

¹ College of Horticulture and Forestry Science, Hubei Engineering Technology Research Center for Forestry Information, Huazhong Agricultural University, Wuhan 430070, China; luojie@mail.hzau.edu.cn

² Key Laboratory of Horticultural Plant Biology (Ministry of Education), College of Horticulture and Forestry Science, Huazhong Agricultural University, Wuhan 430070, China

* Correspondence: hupodingxiangyu@mail.hzau.edu.cn (J.-J.Z.); jinzhizhang@mail.hzau.edu.cn (J.-Z.Z.); Tel.: +86-27-8728-1826 (J.-Z.Z.); Fax: +86-27-8728-2010 (J.-Z.Z.)

Received: 1 December 2017; Accepted: 13 January 2018; Published: 16 January 2018

Abstract: Auxin plays a crucial role in the diverse cellular and developmental responses of plants across their lifespan. Plants can quickly sense and respond to changes in auxin levels, and these responses involve several major classes of auxin-responsive genes, including the *Auxin/Indole-3-Acetic Acid (Aux/IAA)* family, the *auxin response factor (ARF)* family, *small auxin upregulated RNA (SAUR)*, and the *auxin-responsive Gretchen Hagen3 (GH3)* family. Aux/IAA proteins are short-lived nuclear proteins comprising several highly conserved domains that are encoded by the auxin early response gene family. These proteins have specific domains that interact with ARFs and inhibit the transcription of genes activated by ARFs. Molecular studies have revealed that Aux/IAA family members can form diverse dimers with ARFs to regulate genes in various ways. Functional analyses of Aux/IAA family members have indicated that they have various roles in plant development, such as root development, shoot growth, and fruit ripening. In this review, recently discovered details regarding the molecular characteristics, regulation, and protein–protein interactions of the Aux/IAA proteins are discussed. These details provide new insights into the molecular basis of the Aux/IAA protein functions in plant developmental processes.

Keywords: auxin; *Aux/IAA* gene family; regulation; function

1. Introduction

In plants, many developmental processes are finely tuned by auxin, such as vascular tissue formation, adventitious root initiation, tropistic responses, apical dominance, and flower and fruit development. Auxin also affects cellular processes, such as cell division, enlargement, and differentiation [1–4]. Dynamic spatial and temporal changes in auxin levels can trigger gene reprogramming precisely and rapidly, which requires auxin early response genes, such as the *Auxin/Indole-3-Acetic Acid (Aux/IAA)* family, the *auxin response factor (ARF)* family, *small auxin upregulated RNA (SAUR)*, *aminocyclopropane-1-carboxylic acid synthase (ACS)*, *glutathione-S-transferase (GH2/4-like)*, and the *auxin-responsive Gretchen Hagen3 (GH3)* family [3,5,6]. Among these genes, Aux/IAA family members have been identified as short-lived nuclear proteins that play a crucial role in repressing the expression levels of genes activated by ARFs [7,8]. It has also been demonstrated that auxin-mediated transcriptional regulation is exclusively dependent on the functions of Aux/IAA [9]. Aux/IAA proteins have been suggested to bind with ARFs and prevent activation of auxin-responsive genes in the absence of auxin. At high auxin levels, these proteins can be ubiquitinated by interacting with TRANSPORT INHIBITOR RESPONSE 1/AUXIN SIGNALING F-BOX (TIR1/AFB) receptors and subsequently degraded via the 26S proteasome [10–12]. The released ARFs regulate the expression of auxin-responsive genes [10].

The different TIR1/AFB-Aux/IAA protein combinations usually have differential auxin-binding affinities, and the auxin levels are different in different tissues and developmental phases, which results in distinct auxin-sensing effects [9,11,13,14]. Thus, the spatial-temporal dynamic change in auxin levels can be finely transformed into gene reprogramming signaling, thereby regulating the processes of plant growth and development in a precise manner.

Over the past two decades, mutations in *Aux/IAA* genes have been intensely studied in *Arabidopsis*, and these studies have revealed that the members of this family display distinct functions in plant growth and development processes [15–18]. To date, 29 *Aux/IAA* gene family members in *Arabidopsis* have been isolated [18–20]. Meanwhile, many candidate genes that potentially regulate the stability of *Aux/IAA* proteins have also been identified [21,22]. It is noteworthy that a large number of *Aux/IAA* family members have also been identified in other plants, including *Eucalyptus grandis*, *Solanum lycopersicon*, *Cucumis sativus*, *Populus trichocarpa*, *Zea mays*, and *Oryza sativa* [23–29]. Recently, through the combined application of physiological, genetic, molecular, and biochemical methods, increasingly abundant new information regarding the mechanisms of *Aux/IAA* in regulating auxin signal transduction and auxin degradation has been obtained, and this information provides valuable opportunities to clarify the developmental processes fine-tuned by auxin signaling in plants. In this review, recent advances in the roles of *Aux/IAA* in regulating plant growth and development, as well as global and new insights into the underlying molecular mechanisms of these processes, will be presented.

2. Identification, Gene Duplication, and Molecular Structure of *Aux/IAA* Proteins in Plants

The first isolated *Aux/IAA* genes were the *PS-IAA4/5* and *PS-IAA6* genes from pea [30,31]. Subsequently, 14 *Aux/IAA* genes were isolated from *Arabidopsis* based on the homologues to *PS-IAA4/5* and *PS-IAA6* [6]. With the advent of genome sequencing, the *IAA/Aux* gene family has been identified in 30 plant species by genome-wide analysis, including 29 genes from *Arabidopsis*, 55 from *Brassica rapa*, 26 from Citrus, 26 from *Eucalyptus grandis*, 26 from *Solanum lycopersicum*, 26 from *Solanum tuberosum*, 26 from *Vitis vinifera*, and 35 from *Populus trichocarpa* (Table 1) [19,23,24,26,32–35]. Among these species, the largest *Aux/IAA* gene family (119 members) was observed in the genome of *Brassica napus*, largely as a result of duplication in different subgenomes [36]. In addition, a large proportion of *Aux/IAA* genes was observed to be generated from gene duplication events during the course of evolution, such as segmental duplication, tandem duplication, or both [29,37]. For example, 76% and 90% of *Aux/IAA* genes were segmentally duplicated in the genomes of *Arabidopsis* and *Glycine max*, respectively [29,37]. Curiously, only 6% (2 out of 31) of *Aux/IAA* genes were tandemly correlated in *Picea abies* [29], even though this species has undergone numerous duplications [38]. The duplication events of the *Aux/IAA* gene family seem to coincide with whole-genome duplication (WGD) [29,37]. For instance, two WGD events and one whole-gene triplication (WGT) have been found in the genome of *Glycine max*, but no WGD event was observed during the genome evolution of *Picea abies* [37–39]. Compared with higher-order plants, no or fewer *Aux/IAA* genes were identified in lower plants, such as *Chlamydomonas reinhardtii* (0), *Marchantia polymorpha* (1), *Physcomitrella patens* (3), and *Selaginella moellendorffii* (9) [29,40–42]. Thus, the expanded *Aux/IAA* gene members in land plants create functional redundancy and may be associated with new functions to adapt to environmental changes [43–45].

Table 1. Genome-wide identification of the Aux/IAA protein family in 32 plant species.

Species	Protein No.	Truncated Protein No.	Complete Protein No.	Reference
<i>Amborella trichopoda</i>	16	8 (50%)	8 (50%)	[29]
<i>Arabidopsis thaliana</i>	29	11 (38%)	18 (62%)	[19]
<i>Brachypodium distachyon</i>	27	8 (30%)	19 (70%)	[29]
<i>Brassica rapa</i>	55	18 (33%)	37 (67%)	[46]
<i>Brassica napus</i>	119	75 (63%)	44 (37%)	[36]
<i>Carica papaya</i>	18	8 (44%)	10 (56%)	[47]
<i>Cicer arietinum</i>	22	10 (45%)	12 (55%)	[37]
Citrus	26	3 (12%)	23 (88%)	[33]
<i>Cucumis sativus</i>	27	9 (33%)	18 (67%)	[25]
<i>Eucalyptus grandis</i>	26	9 (35%)	17 (65%)	[23]
<i>Glycine max</i>	63	22 (35%)	41 (65%)	[37]
<i>Gossypium raimondii</i>	44	14 (32%)	30 (68%)	[29]
<i>Marchantia polymorpha</i>	1	-	1 (100%)	[41,42]
<i>Medicago truncatula</i>	17	6 (35%)	11 (65%)	[48]
<i>Oryza sativa</i>	31	5 (16%)	26 (84%)	[28]
<i>Petunia hybrida</i>	16	-	-	[40]
<i>Phalaenopsis equestris</i>	16	8 (50%)	8 (50%)	[29]
<i>Physcomitrella patens</i>	3	1 (33%)	2 (67%)	[29]
<i>Picea abies</i>	31	23 (74%)	8 (26%)	[29]
<i>Populus trichocarpa</i>	35	11 (31%)	24 (69%)	[26]
<i>Prunus persica</i>	24	8 (33%)	16 (67%)	[29]
<i>Ricinus communis</i>	21	8 (38%)	13 (62%)	[29]
<i>Selaginella moellendorffii</i>	9	7 (77%)	2 (22%)	[40]
<i>Solanum lycopersicum</i>	26	3 (12%)	23 (88%)	[24]
<i>Solanum melongena</i>	16	-	-	[40]
<i>Solanum tuberosum</i>	26	5 (19%)	21 (81%)	[34]
<i>Sorghum bicolor</i>	26	2 (8%)	24 (92%)	[49]
<i>Triticum aestivum</i>	84	34 (40%)	50 (60%)	[50]
<i>Triticum urartu</i>	27	-	-	[50]
<i>Utricularia gibba</i>	25	7 (28%)	18 (72%)	[29]
<i>Vitis vinifera</i>	26	0 (0%)	26 (100%)	[35]
<i>Zea mays</i>	34	10 (29%)	24 (71%)	[27]

Bioinformatic and molecular studies have relabeled most Aux/IAA proteins to indicate that they generally harbor four conserved characteristic domains: Domain I, II, III, and IV (Figure 1) [51]. Domain I has been identified as a repression domain bearing an ethylene response factor (ERF)-associated amphiphilic repression (EAR) motif, “LxLxL”, which can recruit a TOPLESS (TPL) co-repressor [52]. Domain II is the auxin degron with a conserved “GWPPV” motif, which can directly interact with SCF^{TIR1} and plays a role in rapid protein turnover [32,53]. Domain II also contains a bipartite structural nuclear localization signal (NLS) with the invariant basic doublet KR [6]. Recently, a conserved lysine–arginine (KR) motif located between the ERF-EAR motif and Domain II was identified as a rate motif that aids in tuning the dynamic turnover of Aux/IAA proteins [54,55]. More recently, flexible hotspots with certain specific lysine residues have been identified in the degron-flanking regions, which are closely linked to ubiquitylation mediated by TIR1 [56]. Domain III contains an amphipathic $\beta\alpha\alpha$ -fold with a structure and function similar to the DNA recognition motif in Arc and MetJ repressors [7,57]. Further studies confirmed the roles of the $\beta\alpha\alpha$ -fold in homo- and heterodimerization with Aux/IAA and/or ARF proteins [51,57,58]. Domain IV comprises an acidic region and an SV40 type NLS (PKKKRKV) [59]. In addition, a conserved “GDVP” motif between $\beta 4$ and $\alpha 2$ in Domain IV may contribute to electrostatic protein interactions [59]. Domain III and IV (namely, Domain III/IV) share homology with the carboxy-terminal dimerization domain (CTD) in ARF proteins, which is involved in regulating ARF activity by interacting with Aux/IAA and ARF family proteins via the type I/II Phox and Bem1p (PB1) protein–protein interaction domain [51,58–60]. The type I/II PB1 domains share both an acidic motif (type I) and basic motif (type II), and the type I

motif contains the conserved octicosapeptide (OPCA)-like motif (Dx(D/E)GDx8[E/D], the Phox and Cdc motif, and the atypical protein kinase C interaction domain motif [59,61]. These identified motifs provide physical binding sites for regulating the auxin signaling pathway.




Figure 1. Domain architecture of the canonical Aux/IAA proteins. The canonical Aux/IAA proteins consist of four typical domains, namely, Domains I–IV. Domain I contains an “LxLxL” motif, which is called the ethylene response factor (ERF)-associated amphiphilic repression (EAR) motif. Between Domain I and II, a conserved “KR” motif was identified as a rate motif, and a bipartite nuclear localization signal (NLS) was located between the KR motif and Domain II. The “GWPPV” motif in Domain II is a degron, which controls the turnover of Aux/IAA proteins. Domain III and IV together form type I/II Phox and Bem1p (PB1) domains. Domain III comprises $\beta 1$, $\beta 2$, and $\alpha 1$, and the conserved “K” residue in $\beta 1$ has been identified as a type II basic motif of PB1. Domain IV contains three β -sheets ($\beta 3$ – $\beta 5$) and two α -helices ($\alpha 2$ and $\alpha 3$). One OPCA-like motif in this domain forms the other face of the ePB1 domain, and a conserved “GDVP” motif between $\beta 4$ and $\alpha 2$ may facilitate electrostatic protein interactions. Another NLS was also observed in this domain [5,44,60].

Although most Aux/IAA proteins harbor the typical four domains, a considerable number of truncated Aux/IAA proteins have also been found in different plant species (Table 1). For instance, two Aux/IAA proteins (AtIAA29 and AtIAA33) in *Arabidopsis*, three Aux/IAA proteins (SlIAA13, SlIAA16, and SlIAA20) in tomato, and five Aux/IAA proteins (StIAA13, StIAA15, StIAA16, StIAA18, and StIAA20) in potato lack Domains I and II [29,34]. Meanwhile, some truncated proteins without Domain III or IV have been identified, such as CpIAA11, CpIAA19, CpIAA27, and CpIAA31 from *Carica papaya* [47]. Additionally, a large proportion of truncated Aux/IAA proteins have been found in *Amborella trichopoda*, *Cicer arietinum*, *Phalaenopsis equestris*, *Picea abies*, *Selaginella moellendorffii*, and *Triticum aestivum* (Table 1) [29,37,40,50]. In addition to lacking some essential domains, a tandem duplication of domain II was exclusively identified in PoptrIAA7.1 (a homolog of *Arabidopsis* IAA7) and conserved only in *Populus* species [26]. Overall, the variations in these domains may contribute to the diverse functions in the auxin signaling pathway, which further facilitate the multiple roles played by Aux/IAA in response to changes in the environment among plants.

3. Interaction of Aux/IAA Proteins in Plants

Plants can sense and respond to auxin signaling in a precise and rapid manner, and this response involves a group of genes called the early auxin response genes [44]. The Aux/IAA proteins are well-known as the early auxin response proteins and participate in auxin signaling through interacting with ARF proteins as transcriptional repressors [7,9,62]. The functions of Aux/IAA proteins in auxin signaling have been well summarized [63,64]. In brief, Aux/IAA and ARF have the same type I/II PB1 domain in their homologues, Domain III/IV, and the type I/II PB1 domain carries both positive and negative charges, which facilitate head-to-tail homo- and heterodimerization of Aux/IAA and ARF via electrostatic interactions [58,61,65,66]. At low auxin levels, Aux/IAA proteins directly dimerize with ARF proteins to prevent their physical interaction with transcription initiation complexes and recruit TPL/TPR co-repressors in the EAR domain [52,67]. The recruited TPL/TPR proteins interact with histone deacetylases to induce chromatin condensation [52]. At high auxin levels, the Aux/IAA proteins are brought together with TIR1/AFB proteins by auxin. The F-box in TIR1/AFB proteins can be recognized by SCF-type ubiquitin protein ligase complexes,

and subsequently, the Aux/IAA proteins are poly-ubiquitinated by SCF proteins and degraded through the 26S proteasome proteolytic pathway [12,63,64]. Next, the released ARF proteins may recruit SWITCH/SUCROSE NONFERMENTING (SWI/SNF) to remodel the chromatin into an active state and bind to the auxin response DNA *cis*-elements (AuxREs) to trigger changes in their target auxin response genes at the transcriptional level [5,62]. Based on this mechanism, auxin signaling can result in changes in gene expression levels.

The interaction and regulation of Aux/IAA proteins are closely associated with their modular structure. Domains I and II are well known for their roles in stabilizing Aux/IAA proteins, and Domain II contains a GWPP-core degron sequence, which is the core binding site with TIR/AFB proteins and auxin [68]. Crystal structures have shown that the leucine-rich repeat (LRR) domain of TIR/AFB proteins can form an auxin binding pocket, which allows auxin to anchor at the bottom and “glue” its “GWPP” motif in the degron of Aux/IAA proteins and to occupy the rest of the area of this pocket [12]. Thus, even single-point mutations in the degron region reduce the affinity to TIR/AFB proteins and hamper the degradation of Aux/IAA proteins [69]. In addition, the conserved rate motif, KR, also facilitates the turnover of Aux/IAA proteins, and plants without this rate motif exhibit insensitive auxin response phenotypes similar to plants with degron mutations [54]. Recent evidence showed that auxin can regulate the 26S proteasome-mediated degradation of ubiquitinated Aux/IAA by remobilizing PROTEASOME REGULATOR1 (PTRE1) from the nucleus to the plasma membrane [70,71]. The degradation rates and ubiquitination sites are varied among Aux/IAA proteins, and lysine residues are not necessary for degradation [72]. Variations in the ubiquitylation sites in flexible hotspots of the degron-flanking regions also lead to distinct auxin affinities, which correspond to the degradation dynamics among different IAA proteins [56]. The negative feedback regulation of 26S proteasome activity may contribute to avoiding over-enhancement of auxin signaling and excessive degradation of Aux/IAA proteins [70]. In addition to Domains I and II, Domains III and IV, which are homologous to the carboxy-terminal dimerization domain (CTD) of ARFs, contain the well-known PB1 domain [61]. Based on thermodynamic and structural approaches, a two-pronged electrostatic interaction was identified in the PB1 domain, and the roles of the charged residues on the basic and acidic faces in stabilizing the protein–protein interaction have been revealed [60]. Although Aux/IAA and ARF can form homo- or hetero-oligomers via their PB1 domains, case studies of ARF5 and IAA17 have revealed that their hetero-oligomers have up to 100-fold greater affinity than their homo-oligomers [73]. These studies shed new light on how these PB1-containing proteins interact with each other at the atomic level to facilitate auxin signaling.

Based on large-scale interactome analyses conducted in *Arabidopsis* using affinity capture-Western/MS, yeast two-hybrid, reconstituted complex, and protein-fragment complementation assay (PCA) approaches, a comprehensive physical interactome map of Aux/IAA proteins has been developed in recent years (Figure 2) [13,17,20,74]. This interactome map consists of the essential auxin signaling transduction components mediated by Aux/IAA proteins (Figure 2). For instance, 29 Aux/IAA proteins had physical contact with 20 ARF proteins via 544 interactions, and Aux/IAA homo-interactions and Aux/IAA-ARF hetero-interactions occupied 47% and 45% of interactions, respectively (Figure 2). All 29 Aux/IAA proteins interacted with each other with 253 interactions, and only 13 ARF shared 22 homo-interactions (Figure 2). TPL/TPR proteins are co-repressors that bind to the EAR domain of Aux/IAA. In the interaction network, four TPL/TPR proteins were identified, and most Aux/IAA proteins had physical interactions with these four TPL/TPR proteins, except IAA5, IAA6, IAA9, IAA14, IAA15, IAA20, IAA29, IAA30, IAA31, IAA32, IAA33, and IAA34 (Figure 2). Recently, a report demonstrated that the homo-oligomerization of TIR1 in *Arabidopsis* was another pathway for regulating auxin signaling by affecting Aux/IAA protein turnover [75]. In the interactome map, four out of six TIR1/AFB proteins were involved in this interactome, but only 11 Aux/IAA proteins had direct physical interactions with the four TIR1/AFB proteins (Figure 2). Thus, the large number of homo-interactions among Aux/IAA proteins and the lack of physical interaction with the TIR1/AFB proteins imply that many Aux/IAA proteins may only function as heterodimerization

IAA17-mediated repression of ARF7/19 activity and for lateral root initiation [83]. In addition, *IAA3* has been shown to be a target of two different histone acetyltransferases, specifically, auxin influx carrier LIKE AUXIN RESISTANT 2 (LAX2) and general control nonderepressible 5 (GCN5), which indicates that the *Aux/IAA* genes can also be regulated by epigenetic modifications [84,85]. Thus, the epigenetic modifications also play an important role in regulating the expression levels of *Aux/IAA* genes. Furthermore, the transcription factor PHYTOCHROME INTERACTING FACTOR 4 (PIF4) can promote the expression level of *IAA19* and *IAA29* by directly binding to their promoters to repress the activity of ARF7, thereby negatively regulating phototropism and auxin signaling [86]. Recent studies have uncovered that 21 of 29 *Aux/IAA* genes are the targets of three PIFs (PIF3, PIF4, and PIF5), and 12 *Aux/IAA* genes are upregulated in response to natural shade [87,88]. These results highlight the crucial roles of *Aux/IAA* genes in auxin-mediated light responses. Another transcription factor, *MYB77*, had in vitro interactions with ARF7 or *IAA19* to control lateral root formation [74,89]. In addition, *IAA29* can competitively interact with WRKY57 to antagonize jasmonic acid-induced leaf senescence [90,91]. Recently, other stress-related transcription factors, such as the dehydration-responsive element binding proteins (DREB1)/C-repeat (CRT) binding factor (CBF) family, have been shown to directly bind to the promoters of *IAAs* [92]. For instance, CBF1 and DREB2A can directly regulate the transcription level of *IAA5* and *IAA19*; the latter genes are essential for overcoming desiccation stress in *Arabidopsis* [92]. The basic leucine zipper (bZIP) transcription factor bZIP11 balances energy homeostasis and primary root growth by directly activating the transcriptional reprogramming of *IAA3*, which is a suppressor of root growth [93]. In addition, a nuclear import receptor IMPORTIN-ALPHA 6 (IMP α 6) can efficiently take up *IAA12* into the nucleus, and this process is required for maintaining enough short-lived *IAA12* in the nucleus to regulate primary root initiation [94]. In rice, an inositol polyphosphate kinase (OsIPK2) stabilized OsIAA11 protein by interacting with its Domain II, and overexpression of OsIPK2 caused typical auxin insensitivity and failure to form the lateral root [95]. OsCYP2 (a cyclophilin gene) directly interacts with its co-chaperone suppressor of the G2 allele of *skp1* (OsSGT1) to regulate lateral root initiation by promoting the turnover of *Aux/IAA* proteins [96]. Further study identified a C2HC-type zinc finger protein (OsZFP) that can directly interact with OsCYP2 to regulate lateral root development via the *IAA* pathway, and *OsIAA23* might participate in this *OsZFP*-mediated auxin signaling process [97]. Interestingly, the capsid protein P2 from the rice dwarf virus (RDV) can directly interact with OsIAA10 and block the degradation processes mediated by OsTIR1, and mutations in *OsIAA10* enhanced rice resistance to this virus [98].




Figure 3. *Aux/IAA* proteins might be regulated at different levels. PKL, PICKLE; LAX2, LIKE AUXIN RESISTANT 2; GCN5, general control nonderepressible 5; PIF3, PHYTOCHROME INTERACTING FACTOR 3; PIF4, PHYTOCHROME INTERACTING FACTOR 4; PIF5, PHYTOCHROME INTERACTING FACTOR 5; CBF1, C-repeat binding factor 1; DREB2A, dehydration-responsive element binding protein 2A; bZIP11, basic leucine zipper 11; PHYA, PHYTOCHROME A; RGL3, RGA-LIKE3; UBC13, UBIQUITIN-CONJUGATING ENZYME 13; TIR1, TRANSPORT INHIBITOR RESPONSE 1/AUXIN SIGNALING F-BOX.

Meanwhile, post-translational modifications also play crucial roles in regulating Aux/IAA proteins (Figure 3) [99]. For instance, previous experiments verified that IAA1, IAA3, IAA9, and IAA17 of *Arabidopsis* can be phosphorylated by PHYTOCHROME A (PHYA) in vitro, which provides evidence of auxin involvement in light signaling [100]. In another recent paper, a cyclophilin-type peptidyl-prolyl *cis/trans* isomerase, LATERAL ROOTLESS2 (LRT2), was shown to catalyze the peptidyl-prolyl isomerization of OsIAA11, which facilitates OsIAA11-OsTIR1 interactions and the turnover of OsIAA11 via the 26S proteasome [101]. Recently, a large number of microRNAs (miRNAs) have also been shown to target the upstream regulators of Aux/IAA, but information on posttranscriptional regulation of Aux/IAA proteins by miRNA is still very limited [9,102]. The first miRNA found to cleave *Aux/IAA* has been identified in *Arabidopsis*, and this miRNA, *miR847*, targets *IAA28* mRNA for degradation and positively regulates lateral organ development as a positive regulator [103]. However, the information about post-translational regulation of Aux/IAA proteins is still very limited; further studies are needed to shed new light on this area.

5. Functional Roles of Aux/IAA Genes during Plant Growth and Development Processes

Although there are 29 Aux/IAA proteins in *Arabidopsis* (with some functional redundancy), most mutations in Aux/IAA proteins display similar phenotypes in which auxin signaling transduction is hampered, leading to defects in diverse developmental processes such as embryo development, lateral root initiation and elongation, hypocotyl growth, tropisms, flower organ development, and other processes (Figure 4) [17–19]. For example, a gain-of-function mutant, *iaa1/axr5*, showed defects in root and shoot tropisms [104], but gain-of-function mutations in *iaa2* and *iaa6* lead to similar phenotypes, which exhibited a short hypocotyl and upcurled leaves, in *Arabidopsis* [15,105]. IAA3, IAA14, and IAA18 act redundantly in controlling lateral root formation by interacting with two ARF proteins (ARF7 and ARF19). Additionally, phenotypic analysis of transgenic *Arabidopsis* variants differing in the IAA14 degradation rate indicated that the timing of lateral root initiation is finely tuned by the turnover rate of IAA14 [106]. *IAA7/AXR2* controls the morphological responses induced by light, such as inhibiting hypocotyl elongation and promoting leaf development, as observed in *iaa7/axr2-1* mutants cultured in the dark [107]. *IAA8* can interact with TIR1 to regulate lateral root formation, and overexpression of this gene caused abnormal gravitropism [108]. A mutation in domain II of *IAA8* resulted in abnormal flower organs with low jasmonic acid levels, which was mediated by its interactions with *ARF6* and *AFR8* [17,109]. *IAA12* regulates root meristem initiation during early embryogenesis by interacting with the ARF5 protein, the latter of which is essential for embryonic root and embryo patterning [110,111]. Stabilized *iaa16* mutants display smaller rosettes, shorter root hairs, fewer lateral roots and shorter anther filament elongations than the wild type, and the homozygous *iaa16* mutants are not fertile [16]. *IAA17* is involved in some typical phenotypes controlled by auxin signaling, such as hypocotyl elongation, root gravitropism, and root hair and adventitious root formation [112,113]. A recent report showed that *IAA17* was also involved in cytosolic glutamine synthetase (*GLN1;2*)-mediated ammonium assimilation in *Arabidopsis* root [114]. A gain-of-function analysis verified the roles that *IAA18* plays in embryonic apical patterning by repressing the activity of ARF5 and other ARFs [115]. Interestingly, transcripts of *IAA18* and *IAA28* can be transported from mature leaves to the root via phloem to negatively regulate lateral root formation [116]. *IAA19* participated in hypocotyl growth and lateral root formation by interacting with ARF7, and the *IAA19* mutant failed to maintain hypocotyl gravitropism and regular lateral roots [117]. Gain-of-function *iaa28-1* mutants show a strong auxin phenotype with failure to form lateral roots and a reduction in plant size and apical dominance [118]. Meanwhile, *IAA28* also participates in the process of lateral root founder cell identity by negatively regulating the activity of its downstream transcription factor, GATA23 [119]. Overexpression of three non-canonical *Aux/IAA* genes (*IAA20*, *IAA30*, and *IAA31*) results in similar aberrant phenotypes, and plants with *IAA20* overexpression show modified gravitropic root and hypocotyl growth and collapse of root apical meristem after germination [120].




Figure 4. Summary of the functions of Aux/IAA genes in growth and development processes of *Arabidopsis thaliana* [18,19].

In addition to *Arabidopsis*, significant achievements have also been made in understanding the roles of Aux/IAA proteins in auxin signaling in other plants, such as tomato, potato, rice, and poplar [95,121–124]. In tomato, *SiIAA3* has been verified as a molecular bridge between auxin and ethylene signaling pathways, and *SiIAA3* RNAi plants display decreased petiole epinasty in light-grown seedlings and an exaggerated curvature of the apical hook under dark conditions [121]. *SiIAA9* controls multiple processes mediated by auxin signaling, such as apical dominance, leaf morphogenesis, flower organ development, and fruit set and development [125–127]. Mutations in *SiIAA9* lead to abnormal leaf shape and parthenocarpic fruits [125–127]. Recently, the CRISPR/Cas9 system was optimized to obtain *SiIAA9-crispr* in tomato plant (*Solanum lycopersicum*) mutants, which have phenotypes similar to *SiIAA9* mutant plants, e.g., abnormal leaves and parthenocarpy [128]. *SiIAA15* plays multiple roles during tomato developmental processes, and *SiIAA15*-suppressed plants exhibit enhanced lateral root formation, modified development of axillary shoots, decreased trichome number and fruit set, and increased leaf thickness [129]. *SiIAA17* plays crucial roles in controlling fruit quality, and *SiIAA17*-silenced lines display larger fruit with a thicker pericarp than the wild-type line [130]. *SiIAA27* RNAi plants display modified reproductive organ anatomy and reduced expression levels of genes involved in chlorophyll synthesis [131]. More recently, *SiIAA27* has been shown to positively regulate the formation of arbuscular mycorrhiza and participate in strigolactone biosynthesis by regulating the expression level of *NODULATION SIGNALING PATHWAY1 (NSP1)* [132]. In potato, *StIAA2* repression results in altered shoot morphogenesis, increased plant height, and petiole hyponasty, and transcriptional reprogramming of other *StIAA* genes [122]. For members of the *Aux/IAA* gene family in rice, *OsIAA1* is induced by auxin and plays a role in light response and coleoptile elongation in rice [133]. Overexpression of *OsIAA4* in rice leads to less sensitivity to exogenous auxin, dwarfism, and more tiller angles in comparison with wild-type plants [134]. Overexpression of *OsIAA6* enhances tolerance to drought, however, loss-of-function mutants of *OsIAA6* display much more tiller outgrowth than does the wild type [135]. Conversely,

overexpression of *OsIAA10* increases the number of tillers [98]. Degradation of *OsIAA11* is essential for initiating lateral root formation, and mutations in the degron of *OsIAA11* stabilize this protein and produce defects in the lateral root [95,96,136]. Gain-of-function *OsIAA13* mutants also display phenotypes of reduced lateral roots and defects in the gravitropic response, which are closely related to transcriptional reprogramming of the set of genes involved in lateral root initiation [137]. *OsIAA23* is essential for postembryonic maintenance of the quiescent center by auxin signaling in rice [138]. In maize, *IAA10* can repress the expression level of the *LATERAL ROOT PRIMORDIA 1 (LRP1)* gene by directly binding to its promoter, and the gain-of-function mutant *iaa11 (rum1)* cannot activate the expression of *LRP1* and fails to initiate seminal and lateral roots [139].

In woody plants, overexpression of *PttIAA3m* causes global auxin-mediated gene reprogramming, which is closely linked to cambial cell division and secondary xylem development in hybrid aspen [123]. Overexpression of *PtrIAA14.1* (a homolog of *Arabidopsis IAA7*) in *Arabidopsis* confirms its roles in regulating vascular patterning via interaction with *AFR5* [140]. Recently, another study verified the role of *Eucalyptus EgrIAA4* in fiber formation and secondary cell wall deposition by heterologous expression of this gene in *Arabidopsis* [23].

6. Conclusions and Perspectives

Over the last decade, remarkable progress has been made in understanding the mechanism of auxin perception and signaling transduction processes through forward and reverse genetic approaches. These advances reveal that *Aux/IAA* genes act as hub factors that regulate gene expression in auxin signaling transduction, and understanding of their molecular functions and regulatory machinery in plant growth and development has been greatly enhanced. However, concerning the processes of the *Aux/IAA*-mediated auxin signaling pathway, there are still many gaps that remain to be resolved. For example, there is still very limited information regarding the regulation of *Aux/IAA* genes at both transcriptional and post-transcriptional levels. *Aux/IAA* proteins have many interactions with other proteins in interactome maps, and the specific roles of these interactions are largely unknown. Conversely, current studies of *Aux/IAA* in the auxin signaling pathway have mainly focused on aspects of plant growth and development, but little attention has been paid to understanding the roles of *Aux/IAA* in the auxin-mediated response to environmental interactions, such as drought, nutrient deficiency, and other abiotic stresses. In the future, genetic approaches combined with computational modeling approaches will continue to make important contributions to identifying new candidate genes that regulate *Aux/IAA*, and a new, broader layer of auxin signaling regulatory networks may be discovered. Apart from its function in *Arabidopsis*, the model plant, our knowledge regarding *Aux/IAA* regulation of plant growth and development is very limited, but it is very clear that our examination of these processes in other plants will greatly benefit from the knowledge that is being obtained in model plants.

Acknowledgments: This research was financed by the Fundamental Research Funds for the Central Universities (No. 2662017QD001 and 2662015BQ009), Hubei Province Natural Science Foundation of China (No. 2017CFB338), the National Natural Science Foundation of China (No. 31772252 and 31471863), and the International Foundation for Science No. C/5148-2.

Author Contributions: Jie Luo, Jing-Jing Zhou and Jin-Zhi Zhang wrote the paper.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Goldfarb, B.; Lanz-Garcia, C.; Lian, Z.; Whetten, R. *Aux/IAA* gene family is conserved in the gymnosperm, loblolly pine (*Pinus taeda*). *Tree Physiol.* **2003**, *23*, 1181–1192. [[CrossRef](#)] [[PubMed](#)]
2. Sundberg, E.; Østergaard, L. Distinct and dynamic auxin activities during reproductive development. *Cold Spring Harb. Perspect. Biol.* **2009**, *1*, a001628. [[CrossRef](#)] [[PubMed](#)]
3. Goldental-Cohen, S.; Israeli, A.; Ori, N.; Yasuor, H. Auxin response dynamics during wild-type and entire flower development in tomato. *Plant Cell Physiol.* **2017**, *58*, 1661–1672. [[CrossRef](#)] [[PubMed](#)]

4. Perrotrechenmann, C. Cellular responses to auxin: Division versus expansion. *Cold Spring Harb. Perspect. Biol.* **2010**, *2*, a001446.
5. Guilfoyle, T.J.; Hagen, G. Auxin response factors. *Curr. Opin. Plant Biol.* **2007**, *10*, 453–460. [[CrossRef](#)] [[PubMed](#)]
6. Abel, S.; Nguyen, M.D.; Theologis, A. The PS-IAA4/5-like family of early Auxin-inducible mRNAs in *Arabidopsis thaliana*. *J. Mol. Biol.* **1995**, *251*, 533–549. [[CrossRef](#)] [[PubMed](#)]
7. Abel, S.; Oeller, P.W.; Theologis, A. Early auxin-induced genes encode short-lived nuclear proteins. *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 326–330. [[CrossRef](#)] [[PubMed](#)]
8. Dreher, K.A.; Brown, J.; Saw, R.E.; Callis, J. The *Arabidopsis* Aux/IAA protein family has diversified in degradation and auxin responsiveness. *Plant Cell* **2006**, *18*, 699–714. [[CrossRef](#)] [[PubMed](#)]
9. Lavy, M.; Estelle, M. Mechanisms of auxin signaling. *Development* **2016**, *143*, 3226–3229. [[CrossRef](#)] [[PubMed](#)]
10. Weijers, D.; Benkova, E.; Jäger, K.E.; Schlereth, A.; Hamann, T.; Kientz, M.; Wilmoth, J.C.; Reed, J.W.; Jürgens, G. Developmental specificity of auxin response by pairs of ARF and Aux/IAA transcriptional regulators. *EMBO J.* **2005**, *24*, 1874–1885. [[CrossRef](#)] [[PubMed](#)]
11. Parry, G.; Calderon-Villalobos, L.I.; Prigge, M.; Peret, B.; Dharmasiri, S.; Itoh, H.; Lechner, E.; Gray, W.M.; Bennett, M.; Estelle, M. Complex regulation of the TIR1/AFB family of auxin receptors. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 22540–22545. [[CrossRef](#)] [[PubMed](#)]
12. Tan, X.; Calderon-Villalobos, L.I.A.; Sharon, M.; Zheng, C.; Robinson, C.V.; Estelle, M.; Zheng, N. Mechanism of auxin perception by the TIR1 ubiquitin ligase. *Nature* **2007**, *446*, 640–645. [[CrossRef](#)] [[PubMed](#)]
13. Calderon Villalobos, L.I.; Lee, S.; De Oliveira, C.; Ivetac, A.; Brandt, W.; Armitage, L.; Sheard, L.B.; Tan, X.; Parry, G.; Mao, H.; et al. A combinatorial TIR1/AFB-Aux/IAA co-receptor system for differential sensing of auxin. *Nat. Chem. Biol.* **2012**, *8*, 477–485. [[CrossRef](#)] [[PubMed](#)]
14. Trenner, J.; Poeschl, Y.; Grau, J.; Gogol-Döring, A.; Quint, M.; Delker, C. Auxin-induced expression divergence between *Arabidopsis* species may originate within the TIR1/AFB-AUX/IAA-ARF module. *J. Exp. Bot.* **2017**, *68*, 539. [[PubMed](#)]
15. Tian, Q.; Reed, J.W. Control of auxin-regulated root development by the *Arabidopsis thaliana* SHY2/IAA3 gene. *Development* **1999**, *126*, 711–721. [[PubMed](#)]
16. Rinaldi, M.A.; Liu, J.; Enders, T.A.; Bartel, B.; Strader, L.C. A gain-of-function mutation in IAA16 confers reduced responses to auxin and abscisic acid and impedes plant growth and fertility. *Plant Mol. Biol.* **2012**, *79*, 359–373. [[CrossRef](#)] [[PubMed](#)]
17. Wang, J.; Yan, D.-W.; Yuan, T.-T.; Gao, X.; Lu, Y.-T. A gain-of-function mutation in IAA8 alters *Arabidopsis* floral organ development by change of jasmonic acid level. *Plant Mol. Biol.* **2013**, *82*, 71–83. [[CrossRef](#)] [[PubMed](#)]
18. Overvoorde, P.J.; Okushima, Y.; Alonso, J.M.; Chan, A.; Chang, C.; Ecker, J.R.; Hughes, B.; Liu, A.; Onodera, C.; Quach, H.; et al. Functional genomic analysis of the AUXIN/INDOLE-3-ACETIC ACID gene family members in *Arabidopsis thaliana*. *Plant Cell* **2005**, *17*, 3282–3300. [[CrossRef](#)] [[PubMed](#)]
19. Liscum, E.; Reed, J. Genetics of Aux/IAA and ARF action in plant growth and development. *Plant Mol. Biol.* **2002**, *49*, 387–400. [[CrossRef](#)] [[PubMed](#)]
20. Dharmasiri, N.; Dharmasiri, S.; Weijers, D.; Lechner, E.; Yamada, M.; Hobbie, L.; Ehrismann, J.S.; Jürgens, G.; Estelle, M. Plant development is regulated by a family of auxin receptor F box proteins. *Dev. Cell* **2005**, *9*, 109–119. [[CrossRef](#)] [[PubMed](#)]
21. Hayashi, K.-I.; Jones, A.M.; Ogino, K.; Yamazoe, A.; Oono, Y.; Inoguchi, M.; Kondo, H.; Nozaki, H. Yokonolide B, A novel inhibitor of auxin action, blocks degradation of AUX/IAA factors. *J. Biol. Chem.* **2003**, *278*, 23797–23806. [[CrossRef](#)] [[PubMed](#)]
22. Tian, Q.; Nagpal, P.; Reed, J.W. Regulation of *Arabidopsis* SHY2/IAA3 protein turnover. *Plant J.* **2003**, *36*, 643–651. [[CrossRef](#)] [[PubMed](#)]
23. Yu, H.; Soler, M.; Clemente, H.; Mila, I.; Paiva, J.A.P.; Myburg, A.A.; Bouzayen, M.; Grima-Pettenati, J.; Cassan-Wang, H. Comprehensive genome-wide analysis of the Aux/IAA gene family in eucalyptus: Evidence for the role of EgrIAA4 in wood formation. *Plant Cell Physiol.* **2015**, *56*, 700–714. [[CrossRef](#)] [[PubMed](#)]
24. Audran-Delalande, C.; Bassa, C.; Mila, I.; Regad, F.; Zouine, M.; Bouzayen, M. Genome-wide identification, functional analysis and expression profiling of the Aux/IAA gene family in tomato. *Plant Cell Physiol.* **2012**, *53*, 659–672. [[CrossRef](#)] [[PubMed](#)]

25. Gan, D.; Zhuang, D.; Ding, F.; Yu, Z.; Zhao, Y. Identification and expression analysis of primary auxin-responsive Aux/IAA gene family in cucumber (*Cucumis sativus*). *J. Genet.* **2013**, *92*, 513–521. [[CrossRef](#)] [[PubMed](#)]
26. Kalluri, U.C.; DiFazio, S.P.; Brunner, A.M.; Tuskan, G.A. Genome-wide analysis of Aux/IAA and ARF gene families in *Populus trichocarpa*. *BMC Plant Biol.* **2007**, *7*, 59. [[CrossRef](#)] [[PubMed](#)]
27. Ludwig, Y.; Zhang, Y.; Hochholdinger, F. The maize (*Zea mays* L.) AUXIN/INDOLE-3-ACETIC ACID gene family: Phylogeny, synteny, and unique root-type and tissue-specific expression patterns during development. *PLoS ONE* **2013**, *8*, e78859. [[CrossRef](#)] [[PubMed](#)]
28. Jain, M.; Kaur, N.; Garg, R.; Thakur, J.K.; Tyagi, A.K.; Khurana, J.P. Structure and expression analysis of early auxin-responsive Aux/IAA gene family in rice (*Oryza sativa*). *Funct. Integr. Genom.* **2006**, *6*, 47–59. [[CrossRef](#)] [[PubMed](#)]
29. Wu, W.; Liu, Y.; Wang, Y.; Li, H.; Liu, J.; Tan, J.; He, J.; Bai, J.; Ma, H. Evolution analysis of the Aux/IAA gene family in plants shows dual origins and variable nuclear localization signals. *Int. J. Mol. Sci.* **2017**, *18*, 2107. [[CrossRef](#)] [[PubMed](#)]
30. Oeller, P.W.; Keller, J.A.; Parks, J.E.; Silbert, J.E.; Theologis, A. Structural Characterization of the early indoleacetic acid-inducible genes, PS-IAA4/5 and PS-IAA6, of Pea (*Pisum sativum* L.). *J. Mol. Biol.* **1993**, *233*, 789–798. [[CrossRef](#)] [[PubMed](#)]
31. Theologis, A.; Huynh, T.V.; Davis, R.W. Rapid induction of specific mRNAs by auxin in pea epicotyl tissue. *J. Mol. Biol.* **1985**, *183*, 53–68. [[CrossRef](#)]
32. Kepinski, S.; Leyser, O. The Arabidopsis F-box protein TIR1 is an auxin receptor. *Nature* **2005**, *435*, 446–451. [[CrossRef](#)] [[PubMed](#)]
33. Xie, R.; Pang, S.; Ma, Y.; Deng, L.; He, S.; Yi, S.; Lv, Q.; Zheng, Y. The ARF, AUX/IAA and GH3 gene families in citrus: Genome-wide identification and expression analysis during fruitlet drop from abscission zone A. *Mol. Genet. Genom.* **2015**, *290*, 2089–2105. [[CrossRef](#)] [[PubMed](#)]
34. Wu, J.; Peng, Z.; Liu, S.; He, Y.; Cheng, L.; Kong, F.; Wang, J.; Lu, G. Genome-wide analysis of Aux/IAA gene family in Solanaceae species using tomato as a model. *Mol. Genet. Genom.* **2012**, *287*, 295–311. [[CrossRef](#)] [[PubMed](#)]
35. Çakir, B.; Kiliçkaya, O.; Olcay, A.C. Genome-wide analysis of Aux/IAA genes in *Vitis vinifera*: Cloning and expression profiling of a grape Aux/IAA gene in response to phytohormone and abiotic stresses. *Acta Physiol. Plant.* **2013**, *35*, 365–377. [[CrossRef](#)]
36. Wang, B.; King, G.J.; Li, H.; Wang, J.; Liu, K.; Zhang, Q. Genome-wide analysis of the auxin/indoleacetic acid (Aux/IAA) gene family in allotetraploid rapeseed (*Brassica napus* L.). *BMC Plant Biol.* **2017**, *17*, 204.
37. Singh, V.K.; Jain, M. Genome-wide survey and comprehensive expression profiling of Aux/IAA gene family in chickpea and soybean. *Front. Plant Sci.* **2015**, *6*, 918. [[CrossRef](#)] [[PubMed](#)]
38. Nystedt, B.; Street, N.R.; Wetterbom, A.; Zuccolo, A.; Lin, Y.C.; Scofield, D.G.; Vezzi, F.; Delhomme, N.; Giacomello, S.; Alexeyenko, A.; et al. The Norway spruce genome sequence and conifer genome evolution. *Nature* **2013**, *497*, 579–584. [[CrossRef](#)] [[PubMed](#)]
39. Schmutz, J.; Cannon, S.B.; Schlueter, J.; Ma, J.; Mitros, T.; Nelson, W.; Hyten, D.L.; Song, Q.; Thelen, J.J.; Cheng, J. Genome sequence of the palaeopolyploid soybean. *Nature* **2010**, *463*, 178–183. [[CrossRef](#)] [[PubMed](#)]
40. Kumar, R.; Agarwal, P.; Pareek, A.; Tyagi, A.K.; Sharma, A.K. Genomic Survey, Gene Expression, and Interaction Analysis Suggest Diverse Roles of ARF and Aux/IAA Proteins in Solanaceae. *Plant Mol. Biol. Rep.* **2015**, *33*, 1552–1572. [[CrossRef](#)]
41. Bowman, J.L.; Kohchi, T.; Yamato, K.T.; Jenkins, J.; Shu, S.; Ishizaki, K.; Yamaoka, S.; Nishihama, R.; Nakamura, Y.; Berger, F. Insights into land plant evolution garnered from the *Marchantia polymorpha* Genome. *Cell* **2017**, *171*, 287–304. [[CrossRef](#)] [[PubMed](#)]
42. Kato, H.; Ishizaki, K.; Kouno, M.; Shirakawa, M.; Bowman, J.L.; Nishihama, R.; Kohchi, T. Auxin-mediated transcriptional system with a minimal set of components is critical for morphogenesis through the life cycle in *Marchantia polymorpha*. *PLoS Genet.* **2015**, *11*, e1005084.
43. Singh, V.K.; Rajkumar, M.; Garg, R.; Jain, M. Genome-wide identification and co-expression network analysis provide insights into the roles of auxin response factor gene family in chickpea. *Sci. Rep.* **2017**, *7*, 10895. [[CrossRef](#)] [[PubMed](#)]
44. Weijers, D.; Wagner, D. Transcriptional responses to the auxin hormone. *Ann. Rev. Plant Biol.* **2016**, *67*, 539–574. [[CrossRef](#)] [[PubMed](#)]

45. Ishizaki, K. Evolution of land plants: Insights from molecular studies on basal lineages. *Biosci. Biotechnol. Biochem.* **2017**, *81*, 73–80. [[CrossRef](#)] [[PubMed](#)]
46. Paul, P.; Dhandapani, V.; Rameneni, J.J.; Li, X.; Sivanandhan, G.; Choi, S.R.; Pang, W.; Im, S.; Lim, Y.P. Genome-wide analysis and characterization of Aux/IAA family genes in *Brassica rapa*. *PLoS ONE* **2016**, *11*, e0151522. [[CrossRef](#)] [[PubMed](#)]
47. Liu, K.; Yuan, C.; Feng, S.; Zhong, S.; Li, H.; Zhong, J.; Shen, C.; Liu, J. Genome-wide analysis and characterization of Aux/IAA family genes related to fruit ripening in papaya (*Carica papaya* L.). *BMC Genom.* **2017**, *18*, 351. [[CrossRef](#)] [[PubMed](#)]
48. Shen, C.; Yue, R.; Yang, Y.; Zhang, L.; Sun, T.; Xu, L.; Tie, S.; Wang, H. Genome-wide identification and expression profiling analysis of the Aux/IAA gene family in *Medicago truncatula* during the early phase of *Sinorhizobium meliloti* infection. *PLoS ONE* **2014**, *9*, e107495. [[CrossRef](#)] [[PubMed](#)]
49. Wang, S.; Bai, Y.; Shen, C.; Wu, Y.; Zhang, S.; Jiang, D.; Guilfoyle, T.J.; Chen, M.; Qi, Y. Auxin-related gene families in abiotic stress response in *Sorghum bicolor*. *Funct. Integr. Genom.* **2010**, *10*, 533–546. [[CrossRef](#)] [[PubMed](#)]
50. Qiao, L.; Zhang, X.; Han, X.; Zhang, L.; Li, X.; Zhan, H.; Ma, J.; Luo, P.; Zhang, W.; Cui, L.; et al. A genome-wide analysis of the auxin/indole-3-acetic acid gene family in hexaploid bread wheat (*Triticum aestivum* L.). *Front. Plant Sci.* **2015**, *6*, 770. [[CrossRef](#)] [[PubMed](#)]
51. Hagen, G.; Guilfoyle, T. Auxin-responsive gene expression: Genes, promoters and regulatory factors. *Plant Mol. Biol.* **2002**, *49*, 373–385. [[CrossRef](#)] [[PubMed](#)]
52. Szemenyei, H.; Hannon, M.; Long, J.A. TOPLESS mediates auxin-dependent transcriptional repression during Arabidopsis embryogenesis. *Science* **2008**, *319*, 1384–1386. [[CrossRef](#)] [[PubMed](#)]
53. Ramos, J.A.; Zenser, N.; Leyser, O.; Callis, J. Rapid degradation of auxin/indoleacetic acid proteins requires conserved amino acids of domain II and is proteasome dependent. *Plant Cell* **2001**, *13*, 2349–2360. [[CrossRef](#)] [[PubMed](#)]
54. Moss, B.L.; Mao, H.; Guseman, J.M.; Hinds, T.R.; Hellmuth, A.; Kovenock, M.; Noorassa, A.; Lanctot, A.; Villalobos, L.I.A.C.; Zheng, N. Rate motifs tune auxin/indole-3-acetic acid degradation dynamics. *Plant Physiol.* **2015**, *169*, 803–813. [[CrossRef](#)] [[PubMed](#)]
55. Mironova, V.; Teale, W.; Shahriari, M.; Dawson, J.; Palme, K. The systems biology of auxin in developing embryos. *Trends Plant Sci.* **2017**, *22*, 225–235. [[CrossRef](#)] [[PubMed](#)]
56. Winkler, M.; Niemeyer, M.; Hellmuth, A.; Janitzka, P.; Christ, G.; Samodelov, S.L.; Wilde, V.; Majovsky, P.; Trujillo, M.; Zurbriggen, M.D.; et al. Variation in auxin sensing guides AUX/IAA transcriptional repressor ubiquitylation and destruction. *Nat. Commun.* **2017**, *8*, 15706. [[CrossRef](#)] [[PubMed](#)]
57. Morgan, K.E.; Zarembinski, T.L.; Theologis, A.; Abel, S. Biochemical characterization of recombinant polypeptides corresponding to the predicted $\beta\alpha\alpha$ fold in Aux/IAA proteins. *FEBS Lett.* **1999**, *454*, 283–287. [[PubMed](#)]
58. Kim, J.; Harter, K.; Theologis, A. Protein–protein interactions among the Aux/IAA proteins. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 11786–11791. [[CrossRef](#)] [[PubMed](#)]
59. Guilfoyle, T.J.; Hagen, G. Getting a grasp on domain III/IV responsible for auxin response factor–IAA protein interactions. *Plant Sci.* **2012**, *190*, 82–88. [[CrossRef](#)] [[PubMed](#)]
60. Korasick, D.A.; Chatterjee, S.; Tonelli, M.; Dashti, H.; Lee, S.G.; Westfall, C.S.; Fulton, D.B.; Andreotti, A.H.; Amarasinghe, G.K.; Strader, L.C.; et al. Defining a two-pronged structural model for PB1 (Phox/Bem1p) domain interaction in plant auxin responses. *J. Biol. Chem.* **2015**, *290*, 12868–12878. [[CrossRef](#)] [[PubMed](#)]
61. Noda, Y.; Kohjima, M.; Izaki, T.; Ota, K.; Yoshinaga, S.; Inagaki, F.; Ito, T.; Sumimoto, H. Molecular recognition in dimerization between PB1 domains. *J. Biol. Chem.* **2003**, *278*, 43516–43524. [[CrossRef](#)] [[PubMed](#)]
62. Li, S.-B.; Xie, Z.-Z.; Hu, C.-G.; Zhang, J.-Z. A review of auxin response factors (ARFs) in plants. *Front. Plant Sci.* **2016**, *7*, 47. [[CrossRef](#)] [[PubMed](#)]
63. Salehin, M.; Bagchi, R.; Estelle, M. SCFTIR1/AFB-Based auxin perception: Mechanism and role in plant growth and development. *Plant Cell* **2015**, *27*, 9–19. [[CrossRef](#)] [[PubMed](#)]
64. Leyser, O. Auxin Signaling. *Plant Physiol.* **2017**, *176*. [[CrossRef](#)] [[PubMed](#)]
65. Guilfoyle, T.J. The PB1 domain in auxin response factor and Aux/IAA proteins: A versatile protein interaction module in the Auxin response. *Plant Cell* **2015**, *27*, 33–43. [[CrossRef](#)] [[PubMed](#)]
66. Parcy, F.; Vernoux, T.; Dumas, R. A Glimpse beyond structures in auxin-Dependent transcription. *Trends Plant Sci.* **2016**, *21*, 574–583. [[CrossRef](#)] [[PubMed](#)]

67. Ito, J.; Fukaki, H.; Onoda, M.; Li, L.; Li, C.; Tasaka, M.; Furutani, M. Auxin-dependent compositional change in Mediator in ARF7- and ARF19-mediated transcription. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 6562–6567. [[CrossRef](#)] [[PubMed](#)]
68. Dinesh, D.; Villalobos, L.A.; Abel, S. Structural biology of nuclear auxin action. *Trends Plant Sci.* **2016**, *21*, 302–316. [[CrossRef](#)] [[PubMed](#)]
69. Brunoud, G.; Wells, D.M.; Oliva, M.; Larrieu, A.; Mirabet, V.; Burrow, A.H.; Beekman, T.; Kepinski, S.; Traas, J.; Bennett, M.J.; et al. A novel sensor to map auxin response and distribution at high spatio-temporal resolution. *Nature* **2012**, *482*, 103–106. [[CrossRef](#)] [[PubMed](#)]
70. Kong, X.; Zhang, L.; Ding, Z. 26S Proteasome: Hunter and prey in auxin signaling. *Trends Plant Sci.* **2016**, *21*, 546–548. [[CrossRef](#)] [[PubMed](#)]
71. Yang, B.-J.; Han, X.-X.; Yin, L.-L.; Xing, M.-Q.; Xu, Z.-H.; Xue, H.-W. Arabidopsis proteasome regulator1 is required for auxin-mediated suppression of proteasome activity and regulates auxin signalling. *Nat. Commun.* **2016**, *7*, 11388. [[CrossRef](#)] [[PubMed](#)]
72. Gilkerson, J.; Kelley, D.R.; Tam, R.; Estelle, M.; Callis, J. Lysine residues are not required for proteasome-mediated proteolysis of the auxin/indole acidic acid protein IAA1. *Plant Physiol.* **2015**, *168*, 708–720. [[CrossRef](#)] [[PubMed](#)]
73. Han, M.; Park, Y.; Kim, I.; Kim, E.-H.; Yu, T.-K.; Rhee, S.; Suh, J.-Y. Structural basis for the auxin-induced transcriptional regulation by Aux/IAA17. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 18613–18618. [[CrossRef](#)] [[PubMed](#)]
74. Causier, B.; Ashworth, M.; Guo, W.; Davies, B. The TOPLESS interactome: A framework for gene repression in Arabidopsis. *Plant Physiol.* **2012**, *158*, 423–438. [[CrossRef](#)] [[PubMed](#)]
75. Dezfulian, M.H.; Jalili, E.; Roberto, D.A.; Moss, B.L.; Khoo, K.; Nemhauser, J.L.; Crosby, W.L. Oligomerization of SCFTIR1 is essential for aux/iaa degradation and auxin signaling in Arabidopsis. *PLoS Genet.* **2016**, *12*, e1006301. [[CrossRef](#)] [[PubMed](#)]
76. Chatr-aryamontri, A.; Oughtred, R.; Boucher, L.; Rust, J.; Chang, C.; Kolas, N.K.; O'Donnell, L.; Oster, S.; Theesfeld, C.; Sellam, A. The BioGRID interaction database: 2017 update. *Nucl. Acids Res.* **2017**, *45*, D369–D379. [[CrossRef](#)] [[PubMed](#)]
77. Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N.S.; Wang, J.T.; Ramage, D.; Amin, N.; Schwikowski, B.; Ideker, T. Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res.* **2003**, *13*, 2498–2504. [[CrossRef](#)] [[PubMed](#)]
78. Thimm, O.; Bläsing, O.; Gibon, Y.; Nagel, A.; Meyer, S.; Krüger, P.; Selbig, J.; Müller, L.A.; Rhee, S.Y.; Stitt, M. Mapman: A user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *Plant J.* **2004**, *37*, 914–939. [[CrossRef](#)] [[PubMed](#)]
79. Wang, R.; Estelle, M. Diversity and specificity: Auxin perception and signaling through the TIR1/AFB pathway. *Curr. Opin. Plant Biol.* **2014**, *21*, 51–58. [[CrossRef](#)] [[PubMed](#)]
80. Shi, H.; Liu, W.; Wei, Y.; Ye, T. Integration of auxin/indole-3-acetic acid 17 and RGA-LIKE3 confers salt stress resistance through stabilization by nitric oxide in Arabidopsis. *J. Exp. Bot.* **2017**, *68*, 1239–1249. [[CrossRef](#)] [[PubMed](#)]
81. Terrile, M.C.; París, R.; Calderón-Villalobos, L.I.; Iglesias, M.J.; Lamattina, L.; Estelle, M.; Casalougué, C.A. Nitric oxide influences auxin signaling through S-nitrosylation of the Arabidopsis TRANSPORT INHIBITOR RESPONSE 1 auxin receptor. *Plant J.* **2012**, *70*, 492–500. [[CrossRef](#)] [[PubMed](#)]
82. Wen, R.; Wang, S.; Xiang, D.; Venglat, P.; Shi, X.; Zang, Y.; Datla, R.; Xiao, W.; Wang, H. UBC13, an E2 enzyme for Lys63-linked ubiquitination, functions in root development by affecting auxin signaling and Aux/IAA protein stability. *Plant J.* **2014**, *80*, 424–436. [[CrossRef](#)] [[PubMed](#)]
83. Fukaki, H.; Taniguchi, N.; Tasaka, M. PICKLE is required for SOLITARY-ROOT/IAA14-mediated repression of ARF7 and ARF19 activity during Arabidopsis lateral root initiation. *Plant J.* **2006**, *48*, 380–389. [[CrossRef](#)] [[PubMed](#)]
84. Yamamuro, C.; Zhu, J.K.; Yang, Z. Epigenetic modifications and plant hormone action. *Mol. Plant* **2016**, *9*, 57–70. [[CrossRef](#)] [[PubMed](#)]
85. Benhamed, M.; Bertrand, C.; Servet, C.; Zhou, D.-X. Arabidopsis GCN5, HD1, and TAF1/HAF2 Interact to regulate histone acetylation required for light-responsive gene expression. *Plant Cell* **2006**, *18*, 2893–2903. [[CrossRef](#)] [[PubMed](#)]

86. Sun, J.; Qi, L.; Li, Y.; Zhai, Q.; Li, C. PIF4 and PIF5 transcription factors link blue light and auxin to regulate the phototropic response in Arabidopsis. *Plant Cell* **2013**, *25*, 2102–2114. [[CrossRef](#)] [[PubMed](#)]
87. Sellaro, R.; Yanovsky, M.J.; Casal, J.J. Repression of shade-avoidance reactions by sunfleck induction of HY5 expression in Arabidopsis. *Plant J.* **2011**, *68*, 919–928. [[CrossRef](#)] [[PubMed](#)]
88. Iglesias, M.J.; Sellaro, R.; Zurbriggen, M.D.; Casal, J.J. Multiple links between shade avoidance and auxin networks. *J. Exp. Bot.* **2017**, *69*, 213–228. [[CrossRef](#)] [[PubMed](#)]
89. Shin, R.; Burch, A.Y.; Huppert, K.A.; Tiwari, S.B.; Murphy, A.S.; Guilfoyle, T.J.; Schachtman, D.P. The Arabidopsis transcription factor MYB77 modulates auxin signal transduction. *Plant Cell* **2007**, *19*, 2440–2453. [[CrossRef](#)] [[PubMed](#)]
90. Jiang, Y.; Liang, G.; Yang, S.; Yu, D. Arabidopsis WRKY57 functions as a node of convergence for jasmonic acid—And auxin-mediated signaling in jasmonic acid—Induced leaf senescence. *Plant Cell* **2014**, *26*, 230–245. [[CrossRef](#)] [[PubMed](#)]
91. Hu, Y.; Jiang, Y.; Han, X.; Wang, H.; Pan, J.; Yu, D. Jasmonate regulates leaf senescence and tolerance to cold stress: Crosstalk with other phytohormones. *J. Exp. Bot.* **2017**, *68*, 1361–1369. [[CrossRef](#)] [[PubMed](#)]
92. Shani, E.; Salehin, M.; Zhang, Y.; Sanchez, S.E.; Doherty, C.; Wang, R.; Mangado, C.; Song, L.; Tal, I.; Pisanty, O.; et al. Plant stress tolerance requires auxin-sensitive Aux/IAA transcriptional repressors. *Curr. Biol.* **2017**, *27*, 437–444. [[CrossRef](#)] [[PubMed](#)]
93. Weiste, C.; Pedrotti, L.; Selvanayagam, J.; Muralidhara, P.; Fröschel, C.; Novák, O.; Ljung, K.; Hanson, J.; Dröge-Laser, W. The Arabidopsis bZIP11 transcription factor links low-energy signalling to auxin-mediated control of primary root growth. *PLoS Genet.* **2017**, *13*, e1006607. [[CrossRef](#)] [[PubMed](#)]
94. Herud, O.; Weijers, D.; Lau, S.; Jürgens, G. Auxin responsiveness of the MONOPTEROS-BODENLOS module in primary root initiation critically depends on the nuclear import kinetics of the Aux/IAA inhibitor BODENLOS. *Plant J.* **2016**, *85*, 269–277. [[CrossRef](#)] [[PubMed](#)]
95. Chen, Y.; Yang, Q.; Sang, S.; Wei, Z.; Wang, P. Rice inositol polyphosphate kinase (OsIPK2) directly interacts with OsIAA11 to regulate lateral root formation. *Plant Cell Physiol.* **2017**, *58*, 1891–1900. [[CrossRef](#)] [[PubMed](#)]
96. Kang, B.; Zhang, Z.; Wang, L.; Zheng, L.; Mao, W.; Li, M.; Wu, Y.; Wu, P.; Mo, X. OsCYP2, a chaperone involved in degradation of auxin-responsive proteins, plays crucial roles in rice lateral root initiation. *Plant J.* **2013**, *74*, 86–97. [[CrossRef](#)] [[PubMed](#)]
97. Cui, P.; Liu, H.; Ruan, S.; Ali, B.; Gill, R.; Ma, H.; Zheng, Z.; Zhou, W. A zinc finger protein, interacted with cyclophilin, affects root development via IAA pathway in rice. *J. Integr. Plant Biol.* **2017**, *59*, 496–505. [[CrossRef](#)] [[PubMed](#)]
98. Jin, L.; Qin, Q.; Wang, Y.; Pu, Y.; Liu, L.; Wen, X.; Ji, S.; Wu, J.; Wei, C.; Ding, B.; et al. Rice Dwarf Virus P2 protein hijacks auxin signaling by directly targeting the rice OsIAA10 protein, enhancing viral infection and disease development. *PLoS Pathog.* **2016**, *12*, e1005847. [[CrossRef](#)] [[PubMed](#)]
99. Han, S.; Hwang, I. Integration of multiple signaling pathways shapes the auxin response. *J. Exp. Bot.* **2017**, *69*, 189–200. [[CrossRef](#)] [[PubMed](#)]
100. Colón-Carmona, A.; Chen, D.L.; Yeh, K.-C.; Abel, S. Aux/IAA proteins are phosphorylated by phytochrome in vitro. *Plant Physiol.* **2000**, *124*, 1728–1738. [[CrossRef](#)] [[PubMed](#)]
101. Jing, H.; Yang, X.; Zhang, J.; Liu, X.; Zheng, H.; Dong, G.; Nian, J.; Feng, J.; Xia, B.; Qian, Q. Peptidyl-prolyl isomerization targets rice Aux/IAs for proteasomal degradation during auxin signalling. *Nat. Commun.* **2015**, *6*, 7395. [[CrossRef](#)] [[PubMed](#)]
102. Campos-Rivero, G.; Osorio-Montalvo, P.; Sánchez-Borges, R.; Us-Camas, R.; Duarte-Aké, F.; de-la-Peña, C. Plant hormone signaling in flowering: An epigenetic point of view. *J. Plant Physiol.* **2017**, *214*, 16–27. [[CrossRef](#)] [[PubMed](#)]
103. Wang, J.-J.; Guo, H.-S. Cleavage of indole-3-acetic acid inducible28 mRNA by microRNA847 upregulates auxin signaling to modulate cell proliferation and lateral organ growth in Arabidopsis. *Plant Cell* **2015**, *27*, 574–590. [[CrossRef](#)] [[PubMed](#)]
104. Yang, X.; Lee, S.; So, J.H.; Dharmasiri, S.; Dharmasiri, N.; Ge, L.; Jensen, C.; Hangarter, R.; Hobbie, L.; Estelle, M. The IAA1 protein is encoded by AXR5 and is a substrate of SCFTIR1. *Plant J.* **2004**, *40*, 772–782. [[CrossRef](#)] [[PubMed](#)]
105. Kim, B.C.; Soh, M.S.; Kang, B.J.; Furuya, M.; Nam, H.G. Two dominant photomorphogenic mutations of Arabidopsis thaliana identified as suppressor mutations of hy2. *Plant J.* **1996**, *9*, 441–456. [[CrossRef](#)] [[PubMed](#)]

106. Guseman, J.M.; Hellmuth, A.; Lanctot, A.; Feldman, T.P.; Moss, B.L.; Klavins, E.; Calderón Villalobos, L.I.A.; Nemhauser, J.L. Auxin-induced degradation dynamics set the pace for lateral root development. *Development* **2015**, *142*, 905–909. [[CrossRef](#)] [[PubMed](#)]
107. Nagpal, P.; Walker, L.M.; Young, J.C.; Sonawala, A.; Timpte, C.; Estelle, M.; Reed, J.W. AXR2 Encodes a Member of the Aux/IAA Protein Family. *Plant Physiol.* **2000**, *123*, 563–574. [[CrossRef](#)] [[PubMed](#)]
108. Arase, F.; Nishitani, H.; Egusa, M.; Nishimoto, N.; Sakurai, S.; Sakamoto, N.; Kaminaka, H. IAA8 involved in lateral root formation interacts with the TIR1 auxin receptor and ARF transcription factors in Arabidopsis. *PLoS ONE* **2012**, *7*, e43414. [[CrossRef](#)] [[PubMed](#)]
109. Tabata, R.; Ikezaki, M.; Fujibe, T.; Aida, M.; Tian, C.-E.; Ueno, Y.; Yamamoto, K.T.; Machida, Y.; Nakamura, K.; Ishiguro, S. Arabidopsis auxin response factor6 and 8 regulate jasmonic acid biosynthesis and floral organ development via repression of class 1 KNOX genes. *Plant Cell Physiol.* **2009**, *51*, 164–175. [[CrossRef](#)] [[PubMed](#)]
110. Hamann, T.; Benkova, E.; Bäurle, I.; Kientz, M.; Jürgens, G. The Arabidopsis BODENLOS gene encodes an auxin response protein inhibiting MONOPTEROS-mediated embryo patterning. *Genes Dev.* **2002**, *16*, 1610–1615. [[CrossRef](#)] [[PubMed](#)]
111. Hardtke, C.S.; Berleth, T. The Arabidopsis gene MONOPTEROS encodes a transcription factor mediating embryo axis formation and vascular development. *EMBO J.* **1998**, *17*, 1405–1411. [[CrossRef](#)] [[PubMed](#)]
112. Rouse, D.; Mackay, P.; Stirnberg, P.; Estelle, M.; Leyser, O. Changes in auxin response from mutations in an AUX/IAA gene. *Science* **1998**, *279*, 1371–1373. [[CrossRef](#)] [[PubMed](#)]
113. Leyser, H.; Pickett, F.B.; Dharmasiri, S.; Estelle, M. Mutations in the AXR3 gene of Arabidopsis result in altered auxin response including ectopic expression from the SAUR-AC1 promoter. *Plant J.* **1996**, *10*, 403–413. [[CrossRef](#)] [[PubMed](#)]
114. Saito, M.; Konishi, N.; Kanno, K.; Yamaya, T.; Kojima, S. Transcriptional repressor IAA17 is involved in nitrogen use by modulating cytosolic glutamine synthetase GLN1; 2 in Arabidopsis roots. *Soil Sci. Plant Nutr.* **2017**, *63*, 163–170. [[CrossRef](#)]
115. Ploense, S.E.; Wu, M.-F.; Nagpal, P.; Reed, J.W. A gain-of-function mutation in IAA18 alters Arabidopsis embryonic apical patterning. *Development* **2009**, *136*, 1509–1517. [[CrossRef](#)] [[PubMed](#)]
116. Notaguchi, M.; Wolf, S.; Lucas, W.J. Phloem-mobile Aux/IAA transcripts target to the root tip and modify root architecture. *J. Integr. Plant Biol.* **2012**, *54*, 760–772. [[CrossRef](#)] [[PubMed](#)]
117. Tatematsu, K.; Kumagai, S.; Muto, H.; Sato, A.; Watahiki, M.K.; Harper, R.M.; Liscum, E.; Yamamoto, K.T. MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl and formation of lateral roots in Arabidopsis thaliana. *Plant Cell* **2004**, *16*, 379–393. [[CrossRef](#)] [[PubMed](#)]
118. Rogg, L.E.; Lasswell, J.; Bartel, B. A gain-of-function mutation in IAA28 suppresses lateral root development. *Plant Cell* **2001**, *13*, 465–480. [[CrossRef](#)] [[PubMed](#)]
119. De Rybel, B.; Vassileva, V.; Parizot, B.; Demeulenaere, M.; Grunewald, W.; Audenaert, D.; Van Campenhout, J.; Overvoorde, P.; Jansen, L.; Vanneste, S. A novel aux/IAA28 signaling cascade activates GATA23-dependent specification of lateral root founder cell identity. *Curr. Biol.* **2010**, *20*, 1697–1706. [[CrossRef](#)] [[PubMed](#)]
120. Sato, A.; Yamamoto, K.T. Overexpression of the non-canonical Aux/IAA genes causes auxin-related aberrant phenotypes in Arabidopsis. *Physiol. Plant.* **2008**, *133*, 397–405. [[CrossRef](#)] [[PubMed](#)]
121. Chaabouni, S.; Jones, B.; Delalande, C.; Wang, H.; Li, Z.; Mila, I.; Frasse, P.; Latche, A.; Pech, J.-C.; Bouzayen, M. Sl-IAA3, a tomato Aux/IAA at the crossroads of auxin and ethylene signalling involved in differential growth. *J. Exp. Bot.* **2009**, *60*, 1349–1362. [[CrossRef](#)] [[PubMed](#)]
122. Kloosterman, B.; Visser, R.; Bachem, C. Isolation and characterization of a novel potato Auxin/Indole-3-Acetic Acid family member (StIAA2) that is involved in petiole hyponasty and shoot morphogenesis. *Plant Physiol. Biochem.* **2006**, *44*, 766–775. [[CrossRef](#)] [[PubMed](#)]
123. Nilsson, J.; Karlberg, A.; Antti, H.; Lopez-Vernaza, M.; Mellerowicz, E.; Perrot-Rechenmann, C.; Sandberg, G.; Bhalerao, R.P. Dissecting the molecular basis of the regulation of wood formation by auxin in hybrid aspen. *Plant Cell* **2008**, *20*, 843–855. [[CrossRef](#)] [[PubMed](#)]
124. Wang, Y.; Zhang, T.; Wang, R.; Zhao, Y. Recent advances in auxin research in rice and their implications for crop improvement. *J. Exp. Bot.* **2017**, *69*, 255–263. [[CrossRef](#)] [[PubMed](#)]
125. Mazzucato, A.; Cellini, F.; Bouzayen, M.; Zouine, M.; Mila, I.; Minoia, S.; Petrozza, A.; Picarella, M.E.; Ruiu, F.; Carriero, F. A TILLING allele of the tomato Aux/IAA9 gene offers new insights into fruit set mechanisms and perspectives for breeding seedless tomatoes. *Mol. Breed.* **2015**, *35*, 22. [[CrossRef](#)]

126. Wang, H.; Jones, B.; Li, Z.; Frasse, P.; Delalande, C.; Regad, F.; Chaabouni, S.; Latché, A.; Pech, J.-C.; Bouzayen, M. The tomato Aux/IAA transcription factor IAA9 Is involved in fruit development and leaf morphogenesis. *Plant Cell* **2005**, *17*, 2676–2692. [[CrossRef](#)] [[PubMed](#)]
127. Koenig, D.; Bayer, E.; Kang, J.; Kuhlemeier, C.; Sinha, N. Auxin patterns Solanum lycopersicum leaf morphogenesis. *Development* **2009**, *136*, 2997–3006. [[CrossRef](#)] [[PubMed](#)]
128. Ueta, R.; Abe, C.; Watanabe, T.; Sugano, S.S.; Ishihara, R.; Ezura, H.; Osakabe, Y.; Osakabe, K. Rapid breeding of parthenocarpic tomato plants using CRISPR/Cas9. *Sci. Rep.* **2017**, *7*, 507. [[CrossRef](#)] [[PubMed](#)]
129. Deng, W.; Yang, Y.; Ren, Z.; Audran-Delalande, C.; Mila, I.; Wang, X.; Song, H.; Hu, Y.; Bouzayen, M.; Li, Z. The tomato SlIAA15 is involved in trichome formation and axillary shoot development. *New Phytol.* **2012**, *194*, 379–390. [[CrossRef](#)] [[PubMed](#)]
130. Su, L.; Bassa, C.; Audran, C.; Mila, I.; Cheniclet, C.; Chevalier, C.; Bouzayen, M.; Roustan, J.-P.; Chervin, C. The auxin Sl-IAA17 transcriptional repressor controls fruit size via the regulation of endoreduplication-related cell expansion. *Plant Cell Physiol.* **2014**, *55*, 1969–1976. [[CrossRef](#)] [[PubMed](#)]
131. Bassa, C.; Mila, I.; Bouzayen, M.; Audran-Delalande, C. Phenotypes associated with down-regulation of Sl-IAA27 support functional diversity among Aux/IAA family members in tomato. *Plant Cell Physiol.* **2012**, *53*, 1583–1595. [[CrossRef](#)] [[PubMed](#)]
132. Guillotin, B.; Etemadi, M.; Audran, C.; Bouzayen, M.; Becard, G.; Combiér, J.-P. Sl-IAA27 regulates strigolactone biosynthesis and mycorrhization in tomato (var. MicroTom). *New Phytol.* **2017**, *213*, 1124–1132. [[CrossRef](#)] [[PubMed](#)]
133. Thakur, J.K.; Tyagi, A.K.; Khurana, J.P. OsIAA1, an Aux/IAA cDNA from rice, and changes in its expression as influenced by auxin and light. *DNA Res.* **2001**, *8*, 193–203. [[CrossRef](#)] [[PubMed](#)]
134. Song, Y.; Xu, Z.-F. Ectopic Overexpression of an AUXIN/INDOLE-3-ACETIC ACID (Aux/IAA) Gene OsIAA4 in Rice Induces Morphological Changes and Reduces Responsiveness to Auxin. *Int. J. Mol. Sci.* **2013**, *14*, 13645–13656. [[CrossRef](#)] [[PubMed](#)]
135. Jung, H.; Lee, D.-K.; Choi, Y.; Kim, J.-K. OsIAA6, a member of the rice Aux/IAA gene family, is involved in drought tolerance and tiller outgrowth. *Plant Sci.* **2015**, *236*, 304–312. [[CrossRef](#)] [[PubMed](#)]
136. Zhu, Z.X.; Liu, Y.; Liu, S.J.; Mao, C.Z.; Wu, Y.R.; Wu, P. A gain-of-function mutation in OsIAA11 affects lateral root development in rice. *Mol. Plant* **2012**, *5*, 154–161. [[CrossRef](#)] [[PubMed](#)]
137. Kitomi, Y.; Inahashi, H.; Takehisa, H.; Sato, Y.; Inukai, Y. OsIAA13-mediated auxin signaling is involved in lateral root initiation in rice. *Plant Sci.* **2012**, *190*, 116–122. [[CrossRef](#)] [[PubMed](#)]
138. Jun, N.; Gaohang, W.; Zhenxing, Z.; Huanhuan, Z.; Yunrong, W.; Ping, W. OsIAA23-mediated auxin signaling defines postembryonic maintenance of QC in rice. *Plant J.* **2011**, *68*, 433–442. [[CrossRef](#)] [[PubMed](#)]
139. Zhang, Y.; Paschold, A.; Marcon, C.; Liu, S.; Tai, H.; Nestler, J.; Yeh, C.-T.; Opitz, N.; Lanz, C.; Schnable, P.S.; et al. The Aux/IAA gene rum1 involved in seminal and lateral root formation controls vascular patterning in maize (*Zea mays* L.) primary roots. *J. Exp. Bot.* **2014**, *65*, 4919–4930. [[CrossRef](#)] [[PubMed](#)]
140. Liu, S.; Hu, Q.; Luo, S.; Li, Q.; Yang, X.; Wang, X.; Wang, S. Expression of wild-type PtrIAA14.1, a poplar Aux/IAA gene causes morphological changes in Arabidopsis. *Front. Plant Sci.* **2015**, *6*, 388. [[CrossRef](#)] [[PubMed](#)]

