



Article

Na⁺-Dependent High-Affinity Nitrate, Phosphate and Amino Acids Transport in Leaf Cells of the Seagrass *Posidonia oceanica* (L.) Delile

Lourdes Rubio , Delia García-Pérez, María Jesús García-Sánchez and José A. Fernández *

Department of Biología Vegetal, Campus Teatinos, Universidad de Málaga, 29071 Málaga, Spain; lrubio@uma.es (L.R.); delia.garcia.perez@gmail.com (D.G.-P.); mjgs@uma.es (M.J.G.-S.)

* Correspondence: ja_fernandez@uma.es; Tel.: +(34)-952-13-1949

Received: 19 April 2018; Accepted: 22 May 2018; Published: 24 May 2018



Abstract: *Posidonia oceanica* (L.) Delile is a seagrass, the only group of vascular plants to colonize the marine environment. Seawater is an extreme yet stable environment characterized by high salinity, alkaline pH and low availability of essential nutrients, such as nitrate and phosphate. Classical depletion experiments, membrane potential and cytosolic sodium measurements were used to characterize the high-affinity NO₃⁻, Pi and amino acids uptake mechanisms in this species. Net uptake rates of both NO₃⁻ and Pi were reduced by more than 70% in the absence of Na⁺. Micromolar concentrations of NO₃⁻ depolarized mesophyll leaf cells plasma membrane. Depolarizations showed saturation kinetics ($K_m = 8.7 \pm 1 \mu\text{M NO}_3^-$), which were not observed in the absence of Na⁺. NO₃⁻ induced depolarizations at increasing Na⁺ also showed saturation kinetics ($K_m = 7.2 \pm 2 \text{ mM Na}^+$). Cytosolic Na⁺ measured in *P. oceanica* leaf cells ($17 \pm 2 \text{ mM Na}^+$) increased by $0.4 \pm 0.2 \text{ mM Na}^+$ upon the addition of $100 \mu\text{M NO}_3^-$. Na⁺-dependence was also observed for high-affinity L-ala and L-cys uptake and high-affinity Pi transport. All together, these results strongly suggest that NO₃⁻, amino acids and Pi uptake in *P. oceanica* leaf cells are mediated by high-affinity Na⁺-dependent transport systems. This mechanism seems to be a key step in the process of adaptation of seagrasses to the marine environment.

Keywords: *Posidonia*; high-affinity transport; NO₃⁻ uptake; Pi uptake; Na⁺-dependent transport systems; Cytosolic Na⁺; amino acid transport

1. Introduction

Nitrogen and phosphorus are essential nutrients for plants, whose availability may limit biomass maintenance and growth. Seawater is a high salinity and alkaline medium with low concentrations of N and P, often below $5 \mu\text{M}$ in the Mediterranean Sea [1]. The seagrass *Posidonia oceanica*, is a Mediterranean endemism limited by N and P [2], but able to grow under nutrient concentrations low enough (less than $5 \mu\text{M}$ of dissolved inorganic nitrogen and $1 \mu\text{M}$ Pi) to limit the proliferation of other species, as *Cymodocea nodosa* [2,3]. This raises the question of how seagrasses are able to extract nutrients at the low concentrations present in seawater to maintain primary production. To take up nutrients at low external concentrations, plants have to: (i) evolve transport systems with the capability to bind the free ion species with a very low K_m and (ii) a powerful system to energize the movement of the ions across the plasma membranes to compensate the high, outwardly directed, ion motive force. Seagrasses studied so far use the Na⁺ electrochemical gradient to drive the high-affinity uptake of NO₃⁻ and Pi in both roots and leaves [4,5].

In terrestrial vascular plants, where the H⁺-ATPase energizes the plasma membrane [6], NO₃⁻ transport depends on the cellular energy supply and is coupled to the proton electrochemical

gradient [7]. NO_3^- uptake by single cells is associated with depolarization of the plasma membrane, i.e., an increase in the positive charge inside the cell [8]. To account for both the membrane depolarization and the coupling with the proton gradient, it has been proposed that NO_3^- uptake is mediated by a $2\text{H}^+/\text{NO}_3^-$, and therefore electrophoretic, symport mechanism [9,10].

On the other hand, terrestrial plants have evolved a range of physiological and morphological responses which may enhance Pi acquisition under limiting conditions (through symbiotic strategies, root architectural changes, extrusion of organic acids and acid phosphatases by roots, reviewed by [11–13]). As in the case of NO_3^- , transport of inorganic phosphate requires energy and is also driven by the proton electrochemical gradient generated by the plasma membrane H^+ -ATPase [14,15].

In contrast to terrestrial plants, seagrasses can take up mineral nutrients through roots and also through leaves. However, there are several evidences which indicate that leaf tissues have higher affinity for nutrient uptake and can substantially contribute to total nutrient acquisition ([16] and references therein). As a vascular plant, *Z. marina* plasma membrane is energized by a H^+ -ATPase [17]; nevertheless, unlike terrestrial vascular plants, high-affinity NO_3^- and Pi transport mechanisms are fueled not by H^+ but Na^+ in both, root and mesophyll leaf cells [4,5]. Na^+ -coupled transport systems had been previously described in marine organisms, as in the case of the Na^+ -dependent HCO_3^- uptake in marine cyanobacteria [18–20] or the Na^+ -dependent NO_3^- , glucose and amino acids uptake systems in marine diatoms [21,22]. In addition, phosphate transport has been reported to be stimulated by Na^+ in several green algae [23]. However, no functional evidences, apart from the case of *Z. marina*, of Na^+ -dependent high-affinity NO_3^- and Pi uptake systems are found in vascular plant literature.

The use of Na^+ as driving ion for nutrient uptake depends on the maintenance of an inwardly directed electrochemical potential for Na^+ . *Z. marina* maintains homeostatic concentrations of Na^+ around 10 mM [5] and does it in two ways, first by restricting the plasma membrane permeability to Na^+ [17] and second by the operation of an Na^+/H^+ exchanger that takes Na^+ out from the cytosol [24]. Thus, living in the presence of 500 mM Na^+ , a membrane potential below -160 mV guarantees an inwardly directed Na^+ motive force almost 3-fold higher than that for H^+ at the seawater pH [4,5]. Consequently, a Na^+ -coupled transport system could be an important achievement for seagrasses to colonize the marine environment. Recently, we have reported in *P. oceanica*, the operation of a fusicoccin sensitive H^+ -ATPase that provides a highly negative membrane potential (-174 mV) and the direct HCO_3^- uptake in a symport with H^+ [25]. Interestingly, *P. oceanica* mesophyll leaf cells exhibit a cytosolic Na^+ concentration similar to that reported for *Z. marina* (16 ± 1 mM; [25]) sustaining a high inwardly directed Na^+ driving force that could be exploited to fuel high-affinity nutrient uptake.

We have used the information obtained by classical electrophysiology, ion selective intracellular microelectrodes for Na^+ and classical depletion experiments to gain insight into the mechanisms and kinetics of the high-affinity transport systems for NO_3^- , Pi and amino acids (alanine and cysteine) in the mesophyll leaf cells of the marine angiosperm *P. oceanica*. This species shows high-affinity uptake mechanisms for the nutrient investigated, consistent with the ecological nutrient concentrations described for the Mediterranean Sea; the mechanisms for NO_3^- , Pi and amino acids uptake in *P. oceanica*, as previously described for *Z. marina*, are dependent on the presence of Na^+ (not H^+) in the medium, which separates this species from the H^+ -based nutrient uptake model that holds for terrestrial vascular plants. Thus, nutrient transport coupled to the electrochemical gradient for Na^+ could be the general mechanism for high-affinity nutrient uptake energization in seagrasses.

2. Results

2.1. Effect of Na^+ on NO_3^- and Pi Net Uptake in *P. oceanica* Leaves

As aquatic plants, seagrasses can take up nutrients not only through roots but also through the leaves (reviewed by [26]). In the case of *Z. marina*, Na^+ dependence of high-affinity NO_3^- and Pi uptake have been previously demonstrated [4,5]. In order to analyze if this Na^+ dependence also occurs in other seagrasses, NO_3^- and Pi depletion assays were performed in *P. oceanica* leaves incubated

in artificial seawater with or without Na^+ . In artificial seawater (ASW) containing 500 mM NaCl, N-starved *P. oceanica* leaves completely depleted external NO_3^- (100 μM) after 22 h of incubation, while in the absence of Na^+ , the external concentration of NO_3^- remained around 80 μM at the end of incubation. NO_3^- uptake rate, calculated as the slope of the depletion curves, was 4-fold higher in the presence than in the absence of Na^+ (Table 1). Similar results were found in the case Pi uptake assays. In ASW containing 500 mM NaCl, P-starved leaves depleted external Pi (10 μM) at 4 -fold higher uptake rate than in the absence of Na^+ (Table 1). These results strongly suggest the Na^+ -dependence for the NO_3^- and Pi transport in *P. oceanica* leaf cells, to further investigate those transport mechanisms electrophysiological and intracellular ion selective measurements were performed.

Table 1. NO_3^- and Pi net uptake rates in leaves of *P. oceanica*. N- or P-starved plants were incubated in artificial seawater (ASW) containing 500 mM NaCl and Na^+ -free ASW in which NaCl was substituted by 0.8 M Sorbitol, in the case of NO_3^- depletion assays, or 0.5 M Cl-choline in the case of Pi depletion. Net uptake rates were calculated as the slope of the linear phase of NO_3^- and Pi depletion curves within the first 24 h after the addition of 100 μM NO_3K and 10 μM $\text{PO}_4\text{H}_2\text{K}$, respectively. Data are mean \pm SD of three independent replicates.

Net Uptake Rate	ASW	Na^+ -Free ASW
$\mu\text{mol NO}_3^- \cdot \text{g}_{\text{FW}}^{-1} \cdot \text{h}^{-1}$	0.34 ± 0.01	0.09 ± 0.01 *
$\mu\text{mol Pi} \cdot \text{g}_{\text{FW}}^{-1} \cdot \text{h}^{-1}$	0.04 ± 0.003	0.01 ± 0.002 *

* Asterisks indicate significant differences at $p < 0.05$ (Student's *t*-test).

2.2. Na^+ -Dependent High-Affinity NO_3^- and Amino Acid Uptake in Single Mesophyll Leaf Cells

Ion transport across membranes leave an electrical trace such as membrane depolarizations when the net charge that enters the cells is positive. Such membrane depolarizations have been extensively used as a quote of the activity of transport systems [27]. Since the magnitude of the depolarizations reflects a net positive charge that crosses the membrane, in the case of anions, such as NO_3^- or Pi, the membrane depolarization reveals the function of symport mechanisms fueled by a positive driving ion. Thus, to characterize the transport system mechanisms that mediate the high-affinity uptake of NO_3^- , amino acids and Pi in the seagrass *P. oceanica* the effect of micromolar concentration additions on membrane potential of mesophyll leaf cells was analyzed for each nutrient.

Additions of micromolar concentrations of NO_3^- evoked rapid plasma membrane depolarizations in mesophyll leaf cells of N-starved plants (Figure 1A). Measurable changes were evident even at 5 μM NO_3^- and showed saturation at concentrations around 100 μM NO_3^- . As depolarizations are an integral function of the net charge carried by the NO_3^- transport system, the concentration dependence of the depolarizations can be used for the estimation of the affinity of this system for NO_3^- . Fitting the data to the Michaelis-Menten equation yielded a *K_m* value of 8.7 ± 1 μM NO_3^- and a maximum depolarization of 6.3 ± 0.2 mV (Figure 1D, Table 2). On the other hand, the NO_3^- -induced depolarizations were lower in the dark than in the light, but also showed saturation kinetics with a similar *K_m* value, 8.2 ± 1.7 μM NO_3^- but with maximum depolarization reduced by half (Figure 1D, Table 2).

As we previously reported for *Z. marina* [4], NO_3^- -induced depolarizations were not detected in Na^+ -free ASW in *P. oceanica* mesophyll leaf cells. However, after the addition of 30 mM Na^+ to the assay media, the NO_3^- -induced depolarization was recovered (Figure 1B,C). To investigate this Na^+ -dependence, the depolarization induced by 100 μM NO_3^- was assayed at different external Na^+ concentrations. As expected for a Na^+ -dependent transport system, NO_3^- -restored depolarizations showed saturation kinetics at increasing Na^+ concentrations (Figure 1E). Data were fitted to the Michaelis-Menten equation rendering a semisaturation constant of 7.2 ± 1.1 mM Na^+ and a maximum restored depolarization of 4.3 ± 0.1 mV, that was effectively saturated at concentrations around 50 mM Na^+ , supporting the Na^+ -dependence for the high-affinity NO_3^- uptake in *P. oceanica* mesophyll leaf cells.




Figure 1. Effect of the addition of micromolar NO_3^- concentrations on the membrane potential of *P. oceanica* mesophyll leaf cells. Membrane potential changes after the addition of $100 \mu\text{M NO}_3^-$ (downward arrow) in (A) ASW containing 500 mM NaCl, (B) in Na^+ -free ASW and (C) in ASW containing 30 mM NaCl. Vertical lines indicate onset of NO_3^- wash. (D) Mean values of membrane potential depolarizations induced by increasing NO_3^- concentrations in ASW containing 500 mM Na^+ under light (open symbols) or dark (closed symbols) conditions. (E) Mean values of membrane potential depolarizations induced by the addition of $100 \mu\text{M NO}_3^-$ in ASW containing increasing Na^+ concentrations. In D and E data were fitted to the Michaelis-Menten equation and kinetics parameters are shown in Table 2. Values are the means \pm SD of five independent replicates.

To further analyze the $\text{NO}_3^-/\text{Na}^+$ symport, the effect of NO_3^- addition on the cytosolic Na^+ concentration was measured. As shown Figure 2, the addition of $100 \mu\text{M NO}_3^-$ simultaneously evoked a membrane depolarization and the increase of cytosolic Na^+ ($17 \pm 2 \text{ mM Na}^+$) by $0.4 \pm 0.2 \text{ mM Na}^+$. This effect is similar to that formerly found in the case of Na^+ -dependent high-affinity NO_3^- transport in the seagrass *Z. marina* [5] revealing the coupling $\text{NO}_3^-/\text{Na}^+$ uptake and suggesting the use of Na^+ as driving ion for the high-affinity NO_3^- uptake in those species.

To explore the existence of other transport systems fueled by Na^+ in *P. oceanica*, the transport of the amino acids was also analyzed. After a survey among the complete list of them, the two that caused a significant response were alanine (L-ala) and cysteine (L-cys). As in the case of NO_3^- , the addition of micromolar concentrations of L-ala evoked rapid membrane depolarizations in mesophyll leaf cells of N-starved plants. These depolarizations were only recorded in ASW containing Na^+ (Figure 3A). A similar response was observed in the case of micromolar additions of L-cys (data not shown). Since both amino acids exhibit equivalent pH value (6.1), at the pH of seawater (8.2) both solutes show a net negative charge; thus, to depolarize the membrane a net influx of positive charge is needed. As in the case of NO_3^- , Na^+ -dependence of plasma membrane depolarizations strongly suggest that amino acid/ Na^+ symport systems operate at the plasma membrane of *P. oceanica*. In both cases, L-ala and L-cys, induced depolarizations showed saturation kinetics (Figure 3B,C) and values were fitted to the Michaelis-Menten equation, rendering a K_m value of $37 \pm 11 \mu\text{M L-ala}$ and $10 \pm 1 \mu\text{M L-cys}$, respectively

(Table 2). Cysteine transport exhibited lower saturation concentration (around 40 μM L-cys) and lower maximum depolarization than alanine kinetics (Table 2), which saturated at 100 μM L-ala.




Figure 2. Effect of the addition of 100 μM NO_3^- on membrane potential and cytosolic Na^+ concentration of *P. oceanica* mesophyll leaf cells. Black trace shows the membrane potential (E_m , mV) and the blue one corresponds to the cytosolic Na^+ concentration (Na^+_{c} , mM) simultaneously measured. The downward arrow indicates the addition of NO_3^- and the vertical line the onset of the wash. Traces are a representative example of three equivalent measurements.

Table 2. NO_3^- , amino acids and Pi kinetics parameters in mesophyll leaf cells of *P. oceanica*.

Nutrient	K_m (μM)	D_{max} (mV)
NO_3^- (Light)	8.7 ± 1 ($7.2 \pm 1.1 \text{ mM Na}^+$) ¹	6.3 ± 0.2
NO_3^- (Dark)	8.2 ± 1.7	3.1 ± 0.1
L-ala	37 ± 11	16 ± 1.2
L-cys	10 ± 1	5 ± 0.2
Pi	5.8 ± 1 ($4.3 \pm 0.5 \text{ mM Na}^+$) ¹	5.7 ± 0.2

¹ Values in brackets show semisaturation constant for Na^+ as transport driving ion. Data are mean \pm SD of five independent experiments.

2.3. Na^+ -Dependent High-Affinity Pi Uptake in Single Mesophyll Leaf Cells

In P-starved plants, additions of micromolar concentrations of Pi (5–100 μM H_2PO_4^-) evoked rapid membrane depolarizations, unveiling the influx of a net positive charge. As found for other nutrients, no depolarizations were observed in Na^+ -free ASW (data not shown). Pi-induced depolarizations showed saturation kinetics and fitting of the data to the Michaelis-Menten equation rendered a K_m value of $5.8 \pm 1 \mu\text{M}$ H_2PO_4^- and a maximum depolarization of 5.7 ± 0.2 mV (Table 2). On the other hand, depolarizations induced by saturating phosphate concentrations (25 μM Pi) showed Na^+ -dependence and rendered a saturation curve at increasing Na^+ (Figure 4A). Semisaturation constant for Na^+ was 4.3 ± 0.5 mM (Table 2), a lower value than the quoted for NO_3^- transport, suggesting a higher efficiency for Pi transport than for NO_3^- in *P. oceanica* mesophyll leaf cells. Maximum restored depolarization at increasing Na^+ concentrations was 4.5 ± 0.1 mV, which effectively saturated at 50 mM Na^+ (Figure 4B).




Figure 3. Effect of the addition of micromolar amino acid concentrations on the membrane potential of *P. oceanica* mesophyll leaf cells. (A) Membrane potential changes after the addition of 250 μM L-alanine (downward arrows) in ASW containing 500 mM NaCl (black trace) or in Na⁺-free ASW (blue trace); vertical lines show the onset of the wash. Mean values of the membrane potential depolarizations induced by increasing concentrations of L-alanine (B) or L-cysteine (C). Data were fitted to the Michaelis-Menten equation and kinetics parameters are shown in Table 2. Values are the means ± SD of five independent replicates.




Figure 4. Membrane potential depolarizations induced by additions of micromolar H₂PO₄⁻ concentrations. (A) Mean values of membrane depolarizations induced by increasing Pi concentrations in ASW containing 500 mM Na⁺. (B) Mean values of the membrane potential depolarizations induced by the addition of 25 μM H₂PO₄⁻ in ASW containing increasing Na⁺ concentrations. Data were fitted to the Michaelis-Menten equation and kinetics parameters are shown in Table 2. Values are the means ± SD of five independent replicates.

3. Discussion

P. oceanica leaves exhibit higher net rates of NO₃⁻ and Pi uptake in the presence than in the absence of millimolar concentration of Na⁺. Among the seagrasses, this pattern has been only described in the case of *Zostera marina* [4,5]. In full strength Na⁺ artificial seawater, *P. oceanica* NO₃⁻ and Pi net

uptake rates show values in the same range than those quoted at saturating nutrient concentrations in different seagrasses, where, as observed in *P. oceanica*, slightly higher rates are observed for NO_3^- ($3.7\text{--}75 \mu\text{mol NO}_3^- \text{ g}^{-1} \text{ DW h}^{-1}$) than for Pi uptake, $0.014\text{--}43 \mu\text{mol g}^{-1} \text{ DW h}^{-1}$, [28]. The decrease of NO_3^- and Pi uptake rates by *P. oceanica* leaves in the absence of Na^+ strongly suggests the operation of Na^+ -dependent mechanisms for the uptake of both nutrients, as we have previously characterized in *Z. marina* [4,5].

Electrophysiological measurements show that NO_3^- and Pi are incorporated by high-affinity transport systems in mesophyll leaf cells of *P. oceanica*. The semisaturation constant value, obtained from the uptake kinetics is lower for Pi than for NO_3^- , pointing out the higher affinity for Pi uptake in this species. In addition, the semisaturation constant for NO_3^- transport is also higher than the reported value in single mesophyll leaf cells of *Z. marina* ($K_m = 2.3 \pm 0.78 \mu\text{M NO}_3^-$; [4]) but are much lower than those reported for the high-affinity NO_3^- uptake in terrestrial plants, close to $60 \mu\text{M NO}_3^-$ in barley [29] or Arabidopsis [30]; reviewed in [31]. Interestingly, apart from NO_3^- uptake, *P. oceanica* mesophyll leaf cells also show a high-affinity transport of amino acids. Several evidences support that dissolved organic nitrogen utilization seems to be widespread among seagrasses, which provides these species with a competitive advantage over macroalgae in oligotrophic environments [32,33]. Saturation kinetics of amino acids had been previously reported for the tropical seagrasses *Thalassia hemprichii*, *Halodule uninervis*, and *Cymodocea rotundata*, showing K_m values from 3.5 to $10.5 \mu\text{M}$ [34], which are similar to the value obtained in mesophyll leaf cells of *P. oceanica* for cysteine ($10 \pm 1 \mu\text{M}$) but are lower than the K_m for alanine ($37 \pm 11 \mu\text{M}$). However, in contrast to the results of this work, no evidences for Na^+ -dependent amino acids uptake systems have been previously described in seagrasses.

In the case of terrestrial vascular plants, four transport systems have been shown to play a role in root acquisition of amino acids. These are amino acid permeases AtAAP1 [35] and AtAAP5 [36,37], lysine/histidine-type transporter AtLHT1 [38,39], and the compatible solute/proline transporter AtProT2 [40]. LHT1 is probably the most important transporter for root uptake of L-Ala [37]. This high-affinity transporter is also localized at the plasma membrane of mesophyll cells and it has been demonstrated to be responsible for uptake of acidic and neutral amino acids from the leaf apoplast [38]. Interestingly, the K_m values of the high-affinity amino acids transport found in *P. oceanica* leaf cells are similar to those reported in kinetics analysis of LHT1 transport activity, from 7.4 to $44.8 \mu\text{M}$ [37]. In terrestrial vascular plants this transporter functions as a general amino acid permease similar to other AAPs which, when expressed in yeast or oocytes, mediate H^+ -coupled, Na^+ -independent, uptake of a wide variety of amino acids, reviewed in [41].

Thus, Na^+ -dependent high-affinity alanine and cysteine transport mechanisms found in *P. oceanica* mesophyll leaf cells seem to support that secondarily adaptation of vascular plants to alkaline and high Na^+ environments, such seawater, is based, together with other described adaptation processes, on exploiting the high inwardly directed electrochemical gradient for Na^+ , despite the widespread plasma membrane H^+ economy in vascular plants. Na^+ -dependent high-affinity nutrient uptake mechanisms had been found in marine organisms as cyanobacteria [19,20], diatoms and several algae [21–23], but only in the case of our previous work in *Z. marina* it has been demonstrated to operate in vascular plants [4,5].

As observed in *Z. marina*, mesophyll leaf cells of *P. oceanica* also exhibit high-affinity Pi uptake. Semisaturation constant for Pi uptake kinetics is higher than that obtained in epidermal root cells of *Z. marina* ($1.5 \mu\text{M}$; [5]) but similar to the values reported in leaves of the seagrasses *Zostera noltii* ($12.1 \mu\text{M Pi}$; [42]), *Thalassia hemprichii* ($7.7 \mu\text{M Pi}$; [43]) or *Thalassia testudinum* ($11.9 \mu\text{M Pi}$; [44]) and in the range of the high-affinity K_m values described among members of the H^+ -symporter (PHS) family in vascular terrestrial plants [45]. However, as it has been discussed in the case of NO_3^- and amino acids, results shown in this work demonstrated that a Na^+ -coupled high-affinity Pi transport system operates at the plasma membrane of mesophyll leaf cell of *P. oceanica*. Thus, the appearance of Na^+ -dependent high-affinity transporters could be a general adaptation model for nutrient uptake

in seagrasses and could be considered as a strategy to inhabit salinized, alkaline and P and N-poorly media such as seawater.

In the case of terrestrial vascular plants, it has been a long-standing hypothesis that the plasma membrane H^+ -ATPase provides the energy for membrane transport and therefore for nutrient uptake [46]. Plasma membrane H^+ -ATPase is an electrogenic pump that exports protons from the cytosol and, in addition, generates a transmembrane electrochemical gradient of H^+ , and all plasma membrane secondary transporters exploit the energy accumulated in the membrane to drive ion movements [47]. Nevertheless this model seems to be different in seagrasses. *P. oceanica* leaf cells, as was previously reported for *Z. marina*, exhibit a quite constant and low cytosolic Na^+ concentration (16 ± 1 mM; [25] and this work). This value is in the range of those measured in terrestrial glycophytes, but much lower than the proposed values for terrestrial halophytes, between 70 and 200 mM Na^+ in plants grown in the presence of 50 and 400 mM Na^+ , respectively [48]. Despite the absence of Na^+ -ATPases, which are not found in vascular plants [6], the maintenance of the low cytosolic Na^+ concentrations found in seagrasses seems to be the resultant of both, the very low plasma membrane Na^+/K^+ permeability ratio (0.003 in *Z. marina*, [17]; 0.005 in *P. oceanica*, our data, unpublished) and the operation of an active Na^+ efflux mechanism at the plasma membrane [24]. In fact, considering the high Na^+ concentration (500 mM) of the seawater and the rather negative membrane potential (-174 mV; [25]) measured in *P. oceanica* leaf cells, the inwardly directed Na^+ electrochemical gradient (in millivolts) results -260 mV, two-fold higher than the H^+ motive force value (-122 mV) calculated in *P. oceanica*, considering the cytosolic pH of mesophyll leaf cells (7.3, [25]) and the seawater pH (8.2).

As shown in Figure 5, comparisons of Na^+ electrochemical gradient against those calculated for NO_3^- and Pi provide an estimation of Na^+ stoichiometry of the transport systems. Considering the low NO_3^- and Pi concentrations of the Mediterranean Sea and cytosolic concentrations of NO_3^- (3 mM) and Pi (5 mM) similar to the values previously discussed for *Z. marina* [5], around 340 mV are needed to energize the transport of NO_3^- and Pi into *P. oceanica* leaf cells. This means that a stoichiometry of $2Na^+/NO_3^-$ and $2Na^+/Pi$ would render enough energy to make the transport thermodynamically feasible (Figure 5). Na^+ -dependence of both high-affinity transport systems is half-saturated at low millimolar Na^+ concentrations (7.2 ± 1.1 mM and 4.3 ± 0.5 mM, for NO_3^- and Pi uptake, respectively). These values are higher than the K_m for Na^+ reported in *Z. marina* for the high-affinity NO_3^- transport (0.72 mM Na^+ ; [4]) and are also higher than the value reported for Na^+/NO_3^- transport in the marine diatom *Phaeodactylum tricornerutum* (2.58 mM Na^+ ; [22]) or the Na^+ semisaturation constant for Na^+ of the high-affinity HCO_3^- transporters found in cyanobacteria (1 and 1.7 mM; [19]). In any case, these low-millimolar K_m values for Na^+ imply that the NO_3^- and Pi high-affinity transporters of *P. oceanica*, as well as those of *Z. marina*, would be functioning at saturating Na^+ concentrations in seawater.




Figure 5. Energization model and stoichiometry for plasma membrane high-affinity NO_3^- and Pi transporters in mesophyll leaf cells of *P. oceanica*. A Na^+/H^+ antiporter is also indicated as a putative mechanism to maintain the sodium gradient. Cytosolic and seawater ion concentrations and the calculations of the electrochemical potential gradients, included on the big arrows ($\Delta\mu/F$; mV), are discussed in the text.

4. Materials and Methods

4.1. Plant Material and Assay Solutions

Plants of *Posidonia oceanica* (L.) Delile plants were collected, at 2 m depth, in Punta de Calaburras, Málaga, southern Spain ($36^\circ 30' 23.4''$ N $4^\circ 38' 37.6''$ W). In each sampling, several leaves (10–15) attached to a piece of the rhizome were transported to the laboratory in a thermos container in less than 30 min. Leaves were then placed in an aquarium filled with continuously aerated natural seawater. Temperature was held at 15 °C. Plants were illuminated at a light intensity of 150 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ with a photoperiod 16 L/8 D. Seawater was renewed every 3 days and leaves were used for experiments within 2 weeks after sampling.

Plants were nitrogen starved for at least 15 days prior to experiments on NO_3^- and amino acids transport in N-free artificial seawater (ASW) containing 0.01 mM NaH_2PO_4 . On the other hand, plants were starved of phosphate for at least 5 d in P-free ASW containing 0.01 mM NaNO_3 before assays on Pi transport were carried out. The composition of ASW was: 500 mM NaCl, 55 mM MgCl_2 , 12 mM CaCl_2 , 10 mM KCl, 10 mM NaHCO_3 , and pH was adjusted to pH 8.2 with 1 N NaOH.

For depletion experiments and membrane potential measurements ASW was buffered with 10 mM MOPS-Bis tris Propane, pH 8.2. In Na^+ -free ASW, NaCl was replaced by 800 mM Sorbitol or 500 mM Cl-choline, keeping a similar osmolality ($1.09 \text{ osmol Kg}^{-1}$; cryoscopic osmometer, Osmomat, model 030, Gonotec GmbH, Berlin, Germany) and HCO_3^- was added as HCO_3K . NO_3^- and H_2PO_4^- were added as sodium salts, or as potassium salts when using Na^+ -free media. All chemicals were purchased from Sigma-Aldrich, Darmstadt, Germany.

4.2. Depletion Experiments

Excised leaves (0.3–0.6 g fresh weight) were placed separately in 250 mL flasks and incubated in 50 mL ASW containing 500 mM NaCl or in the absence of Na^+ (Na^+ -free ASW). The assay was carried

out at 25 °C with gentle and constant agitation. At the beginning of the experiment, 100 μM NO_3K or 10 μM KH_2PO_4 were added to the assay medium and samples were taken at 0, 5, 10, 15, 30 min, 1, 2, 4, 8, 12 and 24 h. NO_3^- or Pi concentrations were analyzed colorimetrically in each sample [49,50]. Net uptake rates were estimated as the slope of the linear phase of NO_3^- or Pi depletion curves. Three replicates were conducted for each assay.

4.3. Electrophysiology

Membrane potential (E_m) was measured using the standard glass microelectrode technique described by Felle [51]. As previously reported for *Zostera marina* [4,5,17,24,52] or *Posidonia oceanica* [25] leaf pieces (1.5 cm length), in which the epidermis had been partially removed, were mounted in plexiglass chambers (volume 1.1 mL) and connected to a continuous gravity-based flow through system of the assay medium, maintained at a constant flux rate (10 mL min^{-1}) during the measurements.

For membrane potential measurement, mesophyll leaf cells were impaled with single-barreled microelectrodes. Microelectrodes were backfilled with 500 mM KCl and fixed to electrode holders, containing an Ag/AgCl pellet, that were connected to a high-impedance differential amplifier (FD-223a, World Precision Instruments, Sarasota, FL, USA).

For simultaneous measurements of membrane potential and cytosolic sodium, Na^+ -selective microelectrodes based on double-barreled capillaries were used. Details of pulling and backfilling can be found in our previous works [5,24,25]. In summary, ionophore ETH227 (Fluka 71176), dissolved in PVC/THF (4% *w/v*) was used as Na^+ sensor. The backfilling solution was 0.5 M NaCl and the difference signal was calibrated against 1–500 mM NaCl solutions, prepared in 96 mM KCl to minimize K^+ interference. Calibration slopes were close to 45 mV p Na^+ .

4.4. Data Presentation and Analysis

Time-course measurements are shown as single traces, representative of a number of equivalent experiments carried out under the same conditions, as stated in the figure legends. Data are given as mean \pm SD, and the number of repetitions (n) is indicated in every experiment. Membrane depolarization data were fitted to the Michaelis-Menten equation using a non-linear regression computer program (KaleidaGraph, Synergy Software, Reading, PA, USA). Data were analyzed using SPSS Statistics (Armonk, NY, USA), version 21. The significance level was set at $p < 0.05$.

5. Conclusions

Classical depletion experiments, membrane potential and cytosolic Na^+ measurements demonstrated that high-affinity NO_3^- , amino acid and Pi uptake are mediated by Na^+ -dependent transport mechanisms operating in the plasma membrane of mesophyll leaf cells of *P. oceanica*. Thus, the existence of Na^+ -coupled high-affinity uptake systems could be a general model for the mineral nutrition of seagrasses and could be considered a strategy to inhabit salinized, alkaline and low-nutrient environment. However, further investigation is needed to identify the molecular identity of these transporters, transport mechanisms and their occurrence in other seagrass species.

Author Contributions: J.A.F. conceived and designed the experiments. D.G.-P., L.R. and J.A.F. performed the experiments. All co-authors contributed to the ms outline. L.R. and J.A.F. wrote the article with inputs from M.J.G.-S. Figures were designed and drawn by L.R. and J.A.F.

Acknowledgments: The Spanish MINECO, project BFU2017-85117-R participated by JAF and LR, supports open access costs of this publication and both participate in BIO2016-81957-REDT. Project PCIN-2016-090 finances work by MJGS. DG was funded by BES-2012-052685. This work has been done in the framework of Campus de Excelencia Internacional del Mar, CEIMAR (Andalucia Tech).

Conflicts of Interest: The authors declare no conflict of interest. The funding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

Abbreviations

ASW	Artificial Sea Water
D_{max}	Maximum Depolarization
E_m	Membrane Potential

References

- Bethoux, J.P.; El Boukhary, M.S.; Ruiz-Pino, D.; Porin, P.; Copin-Montégut, C. Nutrient, oxygen and carbon ratios, CO₂ sequestration and anthropogenic forcing in the Mediterranean Sea. In *The Mediterranean Sea. The Handbook of Environmental Chemistry*; Saliot, A., Ed.; Springer: Berlin/Heidelberg, Germany, 2005; pp. 67–87, ISBN-13: 978-3-540-25018-0.
- Alcoverro, T.; Romero, J.; Duarte, C.M.; López, N.I. Special and temporal variations in limitation of seagrass *Posidonia oceanica* growth in the NW Mediterranean. *Mar. Ecol. Prog. Ser.* **1997**, *146*, 155–161. [[CrossRef](#)]
- Pérez, M.; Romero, J.; Duarte, C.M.; Sand-Jensen, K. Phosphorus limitation of *Cymodocea nodosa* growth. *Mar. Biol.* **1991**, *109*, 129–133. [[CrossRef](#)]
- García-Sánchez, M.J.; Jaime, M.P.; Ramos, A.; Sanders, D.; Fernández, J.A. Sodium-dependent nitrate transport at the plasma membrane of leaf cells of the marine higher plant *Zostera marina* L. *Plant Physiol.* **2000**, *122*, 879–885. [[CrossRef](#)] [[PubMed](#)]
- Rubio, L.; Linares-Rueda, A.; García-Sánchez, M.J.; Fernández, J.A. Physiological evidence for a sodium-dependent high-affinity phosphate and nitrate transport at the plasma membrane of leaf and root cells of *Zostera marina* L. *J. Exp. Bot.* **2005**, *56*, 613–622. [[CrossRef](#)] [[PubMed](#)]
- Pedersen, C.N.S.; Axelsen, K.B.; Harper, J.F.; Palmgren, M.G. Evolution of plant P-Type ATPases. *Front. Plant Sci.* **2012**, *3*, 1–19. [[CrossRef](#)] [[PubMed](#)]
- Miller, A.J.; Fan, X.; Orsel, M.; Smith, S.J.; Wells, D.M. Nitrate transport and signalling. *J. Exp. Bot.* **2007**, *58*, 2297–2306. [[CrossRef](#)] [[PubMed](#)]
- Crawford, N.M.; Glass, D.M. Molecular and physiological aspects of nitrate uptake in plants. *Trends Plant. Sci.* **1998**, *3*, 389–395. [[CrossRef](#)]
- Miller, A.J.; Smith, S.J. Nitrate transport and compartmentation in cereal root cells. *J. Exp. Bot.* **1996**, *47*, 843–854. [[CrossRef](#)]
- Lupini, A.; Mercati, F.; Araniti, F.; Miller, A.J.; Sunseri, F.; Abenavoli, M.R. NAR2.1/NRT2.1 functional interaction with NO₃⁻ and H⁺ fluxes in high-affinity nitrate transport in maize root regions. *Plant Physiol. Biochem.* **2016**, *102*, 107–114. [[CrossRef](#)] [[PubMed](#)]
- Amtmann, A.; Hammond, J.P.; Armengaud, P.; White, P. Nutrient sensing and signalling in plants: Potassium and phosphorus. *Adv. Bot. Res.* **2006**, *43*, 209–257. [[CrossRef](#)]
- Péret, B.; Clément, M.; Nussaume, L.; Desnos, T. Root developmental adaptation to phosphate starvation: Better safe than sorry. *Trends Plant. Sci.* **2011**, *16*, 442–450. [[CrossRef](#)] [[PubMed](#)]
- Péret, B.; Desnos, T.; Jost, R.; Kanno, S.; Berkowitz, O.; Nussaume, L. Root Architecture Responses: In Search of Phosphate. *Plant Phys.* **2014**, *166*, 1713–1723. [[CrossRef](#)] [[PubMed](#)]
- Ullrich-Eberius, C.I.; Novacky, A.; Fischer, E.; Lüttge, U. Relationship between energy-dependent phosphate uptake and the electrical membrane potential in *Lemma gibba* G1. *Plant Physiol.* **1981**, *67*, 797–801. [[CrossRef](#)] [[PubMed](#)]
- Rausch, C.; Bucher, M. Molecular mechanisms of phosphate transport in plants. *Planta* **2002**, *21*, 23–37. [[CrossRef](#)] [[PubMed](#)]
- Burkholder, J.M.; Tomasko, D.A.; Touchette, B.W. Seagrasses and eutrophication. *J. Exp. Mar. Biol. Ecol.* **2007**, *350*, 46–72. [[CrossRef](#)]
- Lara, C.; Rodríguez, R.; Guerrero, M.G. Sodium-dependent nitrate transport and energetics in cyanobacteria. *J. Phycol.* **1993**, *29*, 389–395. [[CrossRef](#)]
- Shibata, M.; Katoh, H.; Sonoda, M.; Ohkawa, H.; Shimoyama, M.; Fukuzawa, H.; Kaplan, A.; Ogawa, T. Genes essential to sodium-dependent bicarbonate transport in cyanobacteria: Function and phylogenetic analysis. *J. Biol. Chem.* **2002**, *277*, 18658–18664. [[CrossRef](#)] [[PubMed](#)]
- Price, G.D.; Woodger, F.J.; Badger, M.R.; Howitt, S.M.; Tucker, L. Identification of a SulP-type bicarbonate transporter in marine cyanobacteria. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 18228–18233. [[CrossRef](#)] [[PubMed](#)]

20. Hellebust, J.A. Uptake of organic substrates by *Cyclotella cryptica* (Bacillariophyceae): Effects of ions, ionophores and metabolic and transport inhibitors. *J. Phycol.* **1978**, *14*, 79–83. [[CrossRef](#)]
21. Rees, T.A.; Cresswell, R.C.; Syrett, P.J. Sodium-dependent uptake of nitrate and urea by a marine diatom. *Biochim. Biophys. Acta* **1980**, *596*, 141–144. [[CrossRef](#)]
22. Raven, J.A. *Energetics and Transport in Aquatic Plants*; MBL Lectures in Biology; Alan, R. Liss: New York, NY, USA, 1984; Volume 4, ISBN 0-8451-2203-7.
23. Fernández, J.A.; García-Sánchez, M.J.; Felle, H. Physiological evidence for a proton pump at the plasma membrane of the marine angiosperm *Zostera marina* L. *J. Exp. Bot.* **1999**, *50*, 1763–1768.
24. Rubio, L.; Belver, A.; Venema, K.; García-Sánchez, M.J.; Fernández, J.A. Evidence for a sodium efflux mechanism in the leaf cells of the seagrass *Zostera marina* L. *J. Exp. Mar. Biol. Ecol.* **2011**, *402*, 56–64. [[CrossRef](#)]
25. Rubio, L.; García, D.; García-Sánchez, M.J.; Niell, F.X.; Felle, H.H.; Fernández, J.A. Direct uptake of HCO_3^- in the marine angiosperm *Posidonia oceanica* (L.) Delile driven by a plasma membrane H^+ economy. *Plant Cell Environ.* **2017**, *40*, 2820–2830. [[CrossRef](#)] [[PubMed](#)]
26. Romero, J.; Lee, K.-S.; Pérez, M.A.; Alcoverro, T. Nutrients dynamics. In *Seagrasses: Biology, Ecology and Conservation*; Larkum, A.W.D., Orth, R.J., Duarte, C.M., Eds.; Springer: Dordrecht, The Netherlands, 2006; pp. 227–254. ISBN 1-4020-2942-X.
27. Sanders, D.; Hansen, U.-P.; Gradmann, D.; Slayman, C.L. Generalized kinetic analysis of ion-dependent cotransport systems: A unified interpretation of selective ionic effects on Michaelis parameters. *J. Membr. Biol.* **1984**, *77*, 123–152. [[CrossRef](#)] [[PubMed](#)]
28. Touchette, B.W.; Burkholder, J.M. Review of nitrogen and phosphorus metabolism in seagrasses. *J. Exp. Mar. Biol. Ecol.* **2000**, *250*, 133–167. [[CrossRef](#)]
29. Glass, A.D.; Shaff, J.E.; Kochian, L.V. Studies of the uptake of nitrate in barley, IV. Electrophysiology. *Plant Physiol.* **1992**, *99*, 456–463. [[CrossRef](#)] [[PubMed](#)]
30. Okamoto, M.; Kumar, A.; Li, W.; Wang, Y.; Siddiqi, M.Y.; Crawford, N.M.; Glass, A.D.M. High-affinity nitrate transport in roots of *Arabidopsis* depends on expression of the *NAR2*-like gene *AtNRT3.1*. *Plant Phys.* **2006**, *140*, 1036–1046. [[CrossRef](#)] [[PubMed](#)]
31. Wang, Y.Y.; Hsu, P.K.; Tsay, Y.F. Uptake, allocation and signaling of nitrate. *Trends Plant. Sci.* **2012**, *17*, 458–467. [[CrossRef](#)] [[PubMed](#)]
32. Van Engeland, T.; Bouma, T.J.; Morris, E.P.; Brun, F.G.; Peralta, G.; Lara, M.; Hendriks, I.E.; Soetaert, K.; Middelburg, J.J. Potential uptake of dissolved organic matter by seagrasses and macroalgae. *Mar. Ecol. Prog. Ser.* **2011**, *427*, 71–81. [[CrossRef](#)]
33. Nafie, Y.A.; Van Engeland, T.; Van Katwijk, M.M.; Bourma, T. Uptake of nitrogen from compound pools by the seagrass *Zostera noltii*. *J. Exp. Mar. Biol. Ecol.* **2014**, *460*, 47–52. [[CrossRef](#)]
34. Vonk, J.A.; Middelburg, J.J.; Stapel, J.; Bouma, T.J. Dissolved organic nitrogen uptake by seagrasses. *Limnol. Oceanogr.* **2008**, *53*, 542–548. [[CrossRef](#)]
35. Lee, Y.H.; Foster, J.; Chen, J.; Voll, L.; Weber, A.; Tegeder, M. AAP1 transports uncharged amino acids into roots of *Arabidopsis*. *Plant J.* **2007**, *50*, 305–316. [[CrossRef](#)] [[PubMed](#)]
36. Svennerstam, H.; Ganeteg, U.; Näsholm, T. Root uptake of cationic amino acids by *Arabidopsis* depends on functional expression of amino acid permease 5. *New Phytol.* **2008**, *180*, 620–630. [[CrossRef](#)] [[PubMed](#)]
37. Svennerstam, H.; Jämtgård, S.; Ahmad, I.; Huss-Danell, K.; Näsholm, T.; Ganeteg, U. Transporters in *Arabidopsis* roots mediating uptake of amino acids at naturally occurring concentrations. *New Phytol.* **2011**, *191*, 459–467. [[CrossRef](#)] [[PubMed](#)]
38. Hirner, A.; Ladwig, F.; Stransky, H.; Okumoto, S.; Keinath, M.; Harms, A.; Frommer, W.; Koch, W. *Arabidopsis* LHT1 is a high-affinity transporter for cellular amino acid uptake in both root epidermis and leaf mesophyll. *Plant Cell* **2006**, *18*, 1931–1946. [[CrossRef](#)] [[PubMed](#)]
39. Svennerstam, H.; Ganeteg, U.; Bellini, C.; Näsholm, T. Comprehensive screening of *Arabidopsis* mutants suggests the lysine histidine transporter 1 to be involved in plant uptake of amino acids. *Plant Physiol.* **2007**, *143*, 1853–1860. [[CrossRef](#)] [[PubMed](#)]
40. Lehmann, S.; Gumy, C.; Blatter, E.; Boeffel, S.; Fricke, W.; Rentsch, D. In planta function of compatible solute transporters of the *AtProT* family. *J. Exp. Bot.* **2011**, *62*, 787–796. [[CrossRef](#)] [[PubMed](#)]
41. Tegeder, M. Transporters involved in source to sink partitioning of amino acids and ureides: Opportunities for crop improvement. *J. Exp. Bot.* **2014**, *65*, 1865–1878. [[CrossRef](#)] [[PubMed](#)]

42. Pérez-Lloréns, J.L.; Niell, F.X. Short-term phosphate uptake kinetics in *Zostera noltii* Hornem: A comparison between excised leaves and sediment-rooted plants. *Hydrobiologia* **1995**, *297*, 17–27. [[CrossRef](#)]
43. Stapel, J.; Aarts, T.L.; van Duynhoven, B.H.M.; de Groot, J.D.; van den Hoogen, P.H.W.; Hemminga, M.A. Nutrient uptake by leaves and roots of the seagrass *Thalassia hemprichii* in the Spermonde Archipelago, Indonesia. *Mar. Ecol. Prog. Ser.* **1996**, *134*, 195–206. [[CrossRef](#)]
44. Gras, A.F.; Marguerite, S.K.; Madden, C.J. Phosphorus uptake kinetics of a dominant tropical seagrass *Thalassia testudinum*. *Aquat. Bot.* **2003**, *76*, 299–315. [[CrossRef](#)]
45. Rae, A.L.; Cybinski, D.H.; Jarmey, J.M.; Smith, F.W. Characterization of two phosphate transporters from barley: Evidence for diverse function and kinetic properties among members of the *Pht1* family. *Plant Mol. Biol.* **2003**, *53*, 27–36. [[CrossRef](#)] [[PubMed](#)]
46. Palmgren, M.G. Plant plasma membrane H⁺-ATPases: Powerhouses for nutrient uptake. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **2001**, *52*, 817–845. [[CrossRef](#)] [[PubMed](#)]
47. Falhof, J.; Pedersen, J.T.; Fuglsang, A.T.; Palmgren, M. Plasma Membrane H⁺-ATPase Regulation in the Center of Plant Physiology. *Mol. Plant* **2016**, *9*, 323–337. [[CrossRef](#)] [[PubMed](#)]
48. Flowers, T.J.; Munns, R.; Colmer, T.D. Sodium chloride toxicity and the cellular basis of salt tolerance in halophytes. *Ann. Bot.* **2015**, *115*, 419–431. [[CrossRef](#)] [[PubMed](#)]
49. García-Robledo, E.; Corzo, E.; Paspasyrou, S. A fast and direct spectrophotometric method for the sequential determination of nitrate and nitrite at low concentrations in small volumes. *Mar. Chem.* **2014**, *162*, 30–36. [[CrossRef](#)]
50. Fernández, J.A.; Niell, F.X.; Lucena, J. A rapid and sensitive automated determination of phosphate in natural waters. *Limnol. Oceanogr.* **1985**, *30*, 227–230. [[CrossRef](#)]
51. Felle, H. A study of the current-voltage relationships of electrogenic active and passive membrane elements in *Riccia fluitans*. *Biochim. Biophys. Acta* **1981**, *646*, 151–160. [[CrossRef](#)]
52. Rubio, L.; Linares-Rueda, A.; García-Sánchez, M.J.; Fernández, J.A. Ammonium uptake kinetics in root and leaf cells of *Zostera marina* L. *J. Exp. Mar. Biol. Ecol.* **2007**, *352*, 271–279. [[CrossRef](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).