



Review

A Narrative Role of Vitamin D and Its Receptor: With Current Evidence on the Gastric Tissues

Shaima Sirajudeen ¹, Iltaf Shah ² and Asma Al Menhali ^{1,*}

¹ Department of Biology, College of Science, United Arab Emirates University (UAEU),
P.O. Box 15551, Al Ain, United Arab Emirates

² Department of Chemistry, College of Science, United Arab Emirates University (UAEU),
P.O. Box 15551, Al Ain, United Arab Emirates

* Correspondence: asmaa@uaeu.ac.ae; Tel.: +971-3-7136732

Received: 1 July 2019; Accepted: 1 August 2019; Published: 5 August 2019

Abstract: Vitamin D is a major steroid hormone that is gaining attention as a therapeutic molecule. Due to the general awareness of its importance for the overall well-being, vitamin D deficiency (VDD) is now recognized as a major health issue. The main reason for VDD is minimal exposure to sunlight. The vitamin D receptor (VDR) is a member of the steroid hormone receptors that induces a cascade of cell signaling to maintain healthy Ca²⁺ levels that serve to regulate several biological functions. However, the roles of vitamin D and its metabolism in maintaining gastric homeostasis have not yet been completely elucidated. Currently, there is a need to increase the vitamin D status in individuals worldwide as it has been shown to improve musculoskeletal health and reduce the risk of chronic illnesses, including some cancers, autoimmune and infectious diseases, type 2 diabetes mellitus, neurocognitive disorders, and general mortality. The role of vitamin D in gastric homeostasis is crucial and unexplored. This review attempts to elucidate the central role of vitamin D in preserving and maintaining the overall health and homeostasis of the stomach tissue.

Keywords: 1 α ,25(OH)₂D; vitamin D deficiency; vitamin D epimers; cytochrome P450; 1,25-MARRS; stomach

1. Introduction

Vitamin D deficiency (VDD) is a global health issue. More than 1 billion people suffer from vitamin D deficiency or insufficiency [1]. *Endocrine Connections* reported a definite increase in the number of individuals affected with VDD in recent decades [2]. The susceptible risk groups include people who spend more time indoors [3], like the elderly and those who are homebound [4], people with pigmented skins, vegans [5], and pregnant women and their developing children [6,7].

The regions most affected by this disease comprise China, India, Mongolia, and the middle eastern countries including the UAE, Saudi Arabia, Jordan, and Qatar [8]. Despite receiving plentiful sunshine, Vitamin D status is markedly low in the Middle East, for example, over 90% of the UAE population is affected [9]. The deficiency is more pronounced in women than in men in the Middle East, chiefly due to their traditional clothing style of covering the skin completely [10]. The intense heat also contributes to VDD in people staying indoors and in excessive users of sunscreens [11].

The discovery of Vitamin D and its role in skeletal health can be traced back to the middle of the 17th century, to the English physician, Francis Glisson. He published a study titled “A treatise of rickets, as a disease common to children”, wherein he described rickets as an “absolutely new disease that has never been described in any of the ancient or modern books” [12]. The role of diet in this disease was not known at that time [13]. Several years later, Sir Edward Mellanby (1921) ascribed the cause of rickets to the deficiency of a fat-soluble vitamin [14]. However, the breakthrough discovery

was when McCollum and his associates found that this fat-soluble entity was a dietary supplement and named it Vitamin D [15].

The detection and role of the vitamin D receptor (VDR) were followed thereafter in various human and animal tissues. This led to determining the actual role of vitamin D in health and disease. This query led to understanding the role of vitamin D in the development, metabolism, and well-being of the organism, particularly in calcium homeostasis and bone maintenance [16,17]. The non-classical functions of vitamin D and its importance in human physiology became evident over the past few decades [16–18]. Vitamin D is synthesized by cells of the immune system and controls the immune function along with anti-bacterial activities [18,19]. It also plays a role in anti-proliferative and anti-tumorigenic activities in the cancer cells of the breast, colon, skin, stomach, and prostate [20,21]. Low levels of vitamin D have been coupled to various epidemiological states such as inflammatory bowel disease and autoimmune disorders including multiple sclerosis, inflammation [22], chronic kidney disease [23], various types of cancer [20], cardiovascular disease [24], obesity [25], mortality [26], type 1 diabetes mellitus [27], and hypertension [28].

Vitamin D status is assessed by determining the amount of total 25-hydroxyvitamin D (25(OH)D) in circulation. The adequate levels of total 25(OH)D are between 50 and 70 ng/mL. In VDD, the range is less than 30 ng/mL (Table 1) [27,29].

Table 1. The concentrations of circulating 25(OH)D in serum found in various biological conditions.

25(OH)D Levels	Condition
<30 ng/mL	Severe deficiency
30–50 ng/mL	Minor deficiency
50–70 ng/mL	Adequate levels
>80 ng/mL	Excess

There have been a few methods used for the clinical detection of total 25(OH)D. Most of these methods measure total 25(OH)D along with other hydroxylated vitamin D metabolites in human serum. Vitamin D metabolites in circulation at a given time account for 2–20% of the total 25(OH)D present. Therefore, the active levels of vitamin D could be misinterpreted.

The common detection methods of vitamin D in clinical laboratories include high-performance liquid chromatography (HPLC), radioimmunoassay (RIA), enzyme-linked immunoassay (ELISA), and quantitative chemiluminescent immunoassay (CLIA). A kit-based method of RIA developed by DiaSorin S.p.A (Saluggia, Italy) is the gold standard for total 25(OH)D measurements and is used to establish the reference range of serum vitamin D [30,31]. Lately, liquid chromatography-tandem mass spectrometry (LC-MS/MS) is gaining attention due to its application in measuring the various forms of vitamin D and the D3 epimers in the serum of people of all ages [32]. This method is labor-intensive and technically strenuous, but is earning extensive credence. Many publications evaluating the different methods to determine serum vitamin D levels authenticate the fact that this method is cost-effective and highly precise [30,33].

People are given vitamin D supplements to overcome VDD for specific periods, depending on the nature and severity of the scarcity [34]. Global awareness of the health benefits of vitamin D are alarmingly low. For example, there are a few studies investigating the role of vitamin D or its receptor in the gastrointestinal tract (GIT). This review discusses the general characteristics, function, and importance of vitamin D, its metabolism and the two different forms of the VDRs. The second half of the review highlights the role of vitamin D in the stomach tissue and its function in maintaining the homeostasis of the gastric environment and the preservation of gastric health.

2. Vitamin D Ligands

Vitamin D is a hormone. Its active form is commonly known as 1 α ,25-dihydroxyvitamin D (1 α ,25(OH)₂D). The latest research reveals a dual role for 1 α ,25(OH)₂D, which is generated as an activator or inhibitor, depending on the location and function of the target cells [35]. Vitamin D is a

fat-soluble vitamin existing mainly in two forms, ergocalciferol (Vitamin D₂, 1 α ,25(OH)₂D₂), which is of plant origin, and cholecalciferol, (Vitamin D₃, 1 α ,25(OH)₂D₃), which is of animal origin (Figure 1) [36]. The presence of 1 α ,25(OH)₂D₂ has been confirmed in several members of the plant kingdom, including algae, fungi, and in plants contaminated with certain types of fungi. 1 α ,25(OH)₂D₃ on the other hand, is found mostly in dairy products, fatty fish, beef liver, and egg yolks [35].




Figure 1. The structure of vitamin D₂, vitamin D₃, and their precursors. The structural difference between vitamin D₂ and D₃ is present in their side chains. The side chain of vitamin D₃ has a broken ring, while D₂ contains a double bond between carbons, 22 and 23, and a methyl group on carbon 24 on the broken ring.

Askew et al. identified the structure of 1 α ,25(OH)₂D₂ in 1931. Later, the structure of 1 α ,25(OH)₂D₃ was determined through synthetic means [37]. The findings from the second half of the 20th century showed that vitamin D is a prohormone [38]. Prohormones are inactive precursors of hormones that are converted to active anabolic forms with the help of enzymatic activities in the body [39]. Vitamin D is photochemically produced in the skin from 7-dehydrocholesterol [40]. Subsequent chemical reactions in the liver and kidneys convert 7-dehydrocholesterol into the active form that exerts biological functions by binding to the VDRs.

Maestro et al., (2016) reported that VDR is found in most human tissues and has more than 1000 target genes [41]. 1 α ,25(OH)₂D exerts its biological action by binding to VDR and generates biological effects through genomic and rapid non-genomic responses. The details of the two types of VDRs are discussed in Sections 3.1 and 3.2.

2.1. Vitamin D Structure and Biosynthesis

1 α ,25(OH)₂D is a secosteroid with a broken 9,10 carbon-carbon bond in the B ring of the cyclopentanoperhydrophenanthrene structure, which is a carbon skeleton found in steroids. The main structural difference between 1 α ,25(OH)₂D₂ and 1 α ,25(OH)₂D₃ is the presence of a double bond and a methyl group on the side chain of vitamin D₂ (Figure 1) [20,42]. Norman et al. (1982) reported that the structurally flexible nature of 1 α ,25(OH)₂D molecule helps it to successfully fit into the ligand binding domains (LBD) of the vitamin D binding protein (DBP), nuclear VDR (nVDR), and membrane VDR (mVDR), mainly depending on the function of 1 α ,25(OH)₂D [43]. 1 α ,25(OH)₂D can

exist in both bowl-like and planar-like shapes, which support a mechanism by which it can induce many cellular responses [44].

$1\alpha,25(\text{OH})_2\text{D}_3$ is synthesized through the action of sunlight along with subsequent enzymatic reactions. $1\alpha,25(\text{OH})_2\text{D}_2$ is formed by ultraviolet B (UVB) irradiation of the ergosterol in plants and fungi (Figure 2).




Figure 2. Photobiosynthesis and activation of Vitamin D. 7-dehydrocholesterol in the skin is converted to pre-vitamin D3, upon exposure to sunlight, which contains UVB radiation. Pre-vitamin D3 is converted to vitamin D3 in a heat-dependent process. Vitamin D2 and D3 from the diet are incorporated into chylomicrons and introduced into the circulation. Vitamin D formed in the skin or ingested through diet is stored and released from fat cells. In the serum, vitamin D is circulated while being bound to the vitamin D binding protein, DBP. DBP transports it to the liver where it is converted to 25(OH)D (circulating form) by vitamin D-25-hydroxylase. This form of vitamin D is biologically inactive and is converted in the kidneys to the biologically active form, $1\alpha,25(\text{OH})_2\text{D}$ by 25-hydroxyvitamin D-1 α -hydroxylase (1 α -OHase). The $1\alpha,25(\text{OH})_2\text{D}$ binds to the membrane vitamin D receptor (mVDR) or the nuclear vitamin D receptor (nVDR) and elicits specific biological responses.

The skin synthesizes $1\alpha,25(\text{OH})_2\text{D}_3$ from 7-dehydrocholesterol, by the photochemical conversion of pre-vitamin D3 (pre-D). The continued UV-irradiation causes isomerization of pre-D3 to D3, which is taken up by DBP. The presence of methyl group at C24 side chain in vitamin-D2 reduces its affinity for DBP (this is discussed in Section 2.3), resulting in faster clearance from the circulation. This factor limits the conversion of $1\alpha,25(\text{OH})_2\text{D}_2$ to 25OHD2 by the CYP hydroxylases present in the system [45].

The subsequent steps in the biosynthesis, function, and regulation of $1\alpha,25(\text{OH})_2\text{D}$ implicate a major group of enzymes, widely known as cytochrome P450 (CYP) enzymes or vitamin D hydroxylases. The key members of the CYP enzyme family, include CYP2R1, CYP27A1, CYP24A1, CYP2J3, and CYP3A4. In eukaryotes, they are membrane-bound and take part in biosynthetic reactions of steroid hormones and bile acid [46]. The initial step (Figure 2) occurs in the liver and also in some extrahepatic tissues such as the kidney. Vitamin D2 and D3 ingested orally or from the skin are metabolized to form 25-hydroxy vitamin-D2 and D3 ($25(\text{OH})\text{D}_2$ and $25(\text{OH})\text{D}_3$), which is the circulatory form [47]. The multistep reaction is catalyzed by different members of CYP enzymes such as CYP2J3 and CYP3A4, in addition to CYP27A1 (a mitochondrial 25-hydroxylase) and CYP2R1. Both of the former enzymes are expressed in the liver and conserved among species and cause the generation of $25(\text{OH})\text{D}$ by the addition of a hydroxyl group.

Consequently, a dominant target gene of $1\alpha,25(\text{OH})_2\text{D}$ codes for CYP27B1 or 25-hydroxyvitamin D3 1α -hydroxylase, an enzyme found principally in the kidney, which metabolizes $25(\text{OH})\text{D}$ to $1\alpha,25(\text{OH})_2\text{D}$ [48,49]. $1\alpha,25(\text{OH})_2\text{D}$ is the hormonally active form of vitamin D and is responsible for most of its biological functions. Various factors regulate the formation and inhibition of $1\alpha,25(\text{OH})_2\text{D}$. The appropriate signals such as the action of parathyroid hormone (PTH) promote the synthesis of $1\alpha,25(\text{OH})_2\text{D}$, while calcium, fibroblast growth factor-23 (FGF23), and phosphate, inhibit $1\alpha,25(\text{OH})_2\text{D}$ [50]. CYP24A1 is induced by $1\alpha,25(\text{OH})_2\text{D}$ and functions through a feedback mechanism to prevent vitamin D toxicity (VDT).

2.2. Vitamin D Metabolites

Clinicians find it difficult to quantify major metabolites of $1\alpha,25(\text{OH})_2\text{D}$ found in circulation. Most of the vitamin-D intermediate metabolites can be epimerized with the help of epimerization enzymes like 3-epimerase [51]. These epimerization enzymes belong to the CYP group of enzymes and have the ability to further metabolize these epimers as in the common pathway [36]. Epimerization is a mechanism involved in vitamin D metabolism and occurs in extrarenal tissues [52]. Some of the common metabolites of vitamin D that are found in circulation along with their major functions have been described.

2.2.1. 25-hydroxyvitamin-D2 ($25(\text{OH})\text{D}_2$)

The levels of $25(\text{OH})_2\text{D}_2$ (also known as 25-Hydroxyergocalciferol) produced by hydroxylation in the liver reflect an individual's overall vitamin D status. Plants and mushrooms are important sources. Its deficiency has been linked to hypertension, autoimmune diseases, and cancer. It is a bone density conservation agent. Thus, it can inhibit bone resorption and aid in bone mineralization and regeneration. It is used to heal bone fractures and to treat bone diseases, such as osteopenia and osteoporosis [39].

2.2.2. 25-hydroxyvitamin-D3 ($25(\text{OH})\text{D}_3$)

Obtained from the animal diet, 25OHD_3 is transported to the kidney where it is hydroxylated in the mitochondria by 1-hydroxylase to form $1\alpha,25(\text{OH})_2\text{D}_3$ and $24,25(\text{OH})_2\text{D}_3$. The deficiency of this metabolite is linked to hypertension, autoimmune diseases, and cancer [45].

2.2.3. 24, 25-dihydroxyvitamin-D ($24,25(\text{OH})_2\text{D}$)

Vitamin D obtained from the diet goes through a series of biosynthetic reactions in the liver and kidney where it gets converted to either 24, 25-dihydroxyvitamin-D ($24,25(\text{OH})_2\text{D}$) or 1,25-dihydroxyvitamin D ($1\alpha,25(\text{OH})_2\text{D}$) [53]. It has been reported that $24,25(\text{OH})_2\text{D}$ may have a role in bone mineralization [54]. Competition studies conducted by Sömjen et al. in (1982) demonstrated the existence of specific cytoplasmic and nuclear receptors (NRs) for $24,25(\text{OH})_2\text{D}$ D3 [55].

2.2.4. 1- α , 25-dihydroxyvitamin-D3 (1 α ,25(OH)₂D3 or 1,25D3)

1,25D3 is biologically active; has roles in calcium absorption and deposition; affects cellular differentiation and proliferation and central nervous system (CNS); and modulates immune responsiveness. Recent reports advocate it as a major chemopreventive agent against various malignancies including cancers of the stomach, prostate, and colon [56].

2.2.5. 1- α , 25-dihydroxy vitamin-D2 (1 α ,25(OH)₂D2 or 1,25D2)

Activated vitamin-D2 has been used in a broad range of studies to assess the immune function and calcium homeostasis. The ingestion of vitamin D₂ results in an increase in serum concentrations of 1 α ,25(OH)₂D2. Knutson et al., reported that an increase in 1 α ,25(OH)₂D2 is accompanied by a comparable decrease in serum concentrations of 1 α ,25(OH)₂D3 in rats and monkeys [56]

2.2.6. 3-epi-vitamin D

3-epi-vitamin D denotes the following metabolites: 3-epi-25-hydroxyvitamin-D3 (3-epi-25(OH)D3); 3-epi-24, 25-dihydroxy-vitamin-D3 (3-epi-24,25(OH)₂D3); 3-epi-1 α ,25-hydroxyvitamin-D3 (3-epi-1 α ,25(OH)₂D3) [51]. Reddy and colleagues (2001) described an alternate metabolic pathway where C3 epimerization causes a change in the orientation of the hydroxyl group at position C3 of the A ring. 3-epi-vitamin D metabolites have less affinity to DBP and the VDR than the respective 25(OH)D3 and 1 α ,25(OH)₂D3 forms [57]. 3-epi-1 α ,25(OH)₂D3 possesses some anti-proliferative and differentiation activities and suppresses PTH secretion but only at lower levels compared to the non-epimeric compounds [58]. Recent publications have shown the presence of 3-epi-vitamin D in infant, pediatric, and adult populations [37].

Further studies are required to ascertain the source of epimers, whether they are attained exogenously or formed endogenously and to determine their specific roles. Since 1 α ,25(OH)₂D3 extracts are sensitive to photodegradation, the quantification of epimers present in vitamin D supplements is difficult.

2.3. Vitamin D Transport

A minor portion of the free or unbound 1 α ,25(OH)₂D enters the cells through simple diffusion. The majority of circulating vitamin D and its metabolites are attached to special carrier proteins called DBP or albumin. 1 α ,25(OH)₂D-DBP complex enters the cell with the help of membrane proteins like caveolin, cubilin, and megalin or through simple diffusion (Figure 3) [59]. DBP, which is synthesized in the liver is a steroid binding protein and acts as a carrier molecule of 25(OH)D [20,60,61]. It has a higher affinity to 25(OH)D and 24,25(OH)₂D than vitamin D or 1 α ,25(OH)₂D. DBP acts as a storage protein that accumulates 25-OHD under conditions of accessibility to 1 α ,25(OH)₂D and usually releases it for biological requirements [62]. In patients suffering from liver diseases, like specific protein-losing enteropathies or the nephrotic syndrome, the synthesis of normal levels of DBP and albumin may be affected. These individuals may have low total levels of DBP, resulting in the presence of unbound vitamin D metabolites in the serum [63].

Vitamin-D2 and D3-DBP complexes are internalized by the liver. The hepatocytes convert the vitamin D into 25(OH)D. DBP-25(OH)D is internalized by a process called receptor-mediated endocytosis in the renal tubules of the kidney. Megalin and cubilin proteins help in receptor-mediated endocytosis and occur in the proximal convoluted tubules of the kidney and in extra-renal tissues, like mammary cells [64]. Figure 3 shows the critical steps in vitamin D activation taking place in the renal cell. 25(OH)D dissociates from DBP and gets hydroxylated to 1 α ,25-(OH)₂D by CYP27B1. Subsequently, CYP24A1 (24-hydroxylase) can act on two substrates, 1 α ,25(OH)₂D and 25(OH)D through negative feedback mechanisms and convert them to inactive end-products like 1,24,25-trihydroxyvitamin D (1,24,25-(OH)₃D) and 24,25-dihydroxyvitamin D (24,25-(OH)₂D), the roles of which are discussed in Section 5 [65].




Figure 3. The roles of endocytic proteins in the delivery of $1\alpha,25(\text{OH})_2\text{D}$ in the renal cells. The majority of the circulating 25-hydroxyvitamin D is bound to DBP, which is endocytosed via megalin and cubilin-mediated endocytosis. DBP is degraded, and 25(OH)D is either converted to $1\alpha,25(\text{OH})_2\text{D}$ in the mitochondria for CYP27B1-mediated bioactivation or is secreted into circulation where it binds to DBP by CYP24A1-mediated inactivation. Cubilin and megalin then return back to the cell surface and the process gets repeated.

3. Vitamin D Receptors (VDRs)

Maestro et al., (2016) reported that the vitamin D receptor (VDR) is found in most human tissues and has more than 1000 target genes [41]. $1\alpha,25(\text{OH})_2\text{D}_3$ exerts its effect with the help of high-affinity VDR through a series of cell-signaling reactions or as a ligand-activated transcription factor [66,67]. The mode of action may be genomic or non-genomic and can be executed using nVDR or mVDR, respectively [68,69]. In a cell, $1\alpha,25(\text{OH})_2\text{D}_3$ exerts its non-genomic effects by initiating multiple signaling pathways via binding to a membrane receptor. These result in immediate responses in the target cells. However, $1\alpha,25(\text{OH})_2\text{D}_3$ exerts its genomic effects through the NR, which binds to target genes and leads to gene expression [70]. Detailed discussions on the distinguishing features of the two types of receptors, the nature of their diversity, and functions are provided in the upcoming sections.

3.1. Nuclear VDR (nVDR)

The VDR is widely distributed and not limited to the classic target tissues of vitamin D. The VDR belongs to the superfamily of NRs. It acts as a transactivating transcription factor by regulating the expression of genes that mediate its biological activity [71]. The 1,25D-dependent genomic responses in the nucleus materialize as a result of the ligand-activated transcription factors and are relatively slow (may take from several hours to days) [72].

In humans, the VDR is encoded by a large gene (>100 kb), which is present on the long arm of chromosome 12q12. It possesses a complex intron/exon structure and is composed of nine exons (Figure 4) [66]. Exons 1a to 1f encode the 5' untranslated region, UTR of the VDR mRNA and are alternatively spliced. Exons 2 and 3 encode the translation start site, and consists of the DNA-binding domain. This region comprises two highly conserved zinc-finger motifs, one in each domain and is a distinguishing feature of the NRs. Exons 4 to 9 encode the overlapping ligand-binding domain, LBD as well as strong heterodimerization domains. Exon 9 localizes at the 3' UTR region [71]. The *Vdr* gene of a mouse is located on chromosome 15 [72].




Figure 4. A map of the *nVDR* gene on chromosome 12q12. Blue boxes: Exon 1 (a to f), maroon boxes: Exons 2 to 9, yellow box: 3' UTR.

The DNA binding domain of the VDR contains a zinc-finger motif which is the core of the VDR protein, involved in the activation of the target genes. Upon binding of $1\alpha,25(\text{OH})_2\text{D}_3$ to the receptor, two independent protein interaction surfaces form on the VDR protein, one of which is needed for specific DNA binding, while the other recruits the large coregulatory complexes and is necessary for gene modulation [58].

Subsequently, the VDR forms a heterodimer with the retinoid X receptors, RXR (Figure 5). This heterodimer recognizes the vitamin D response element (VDRE) located in the promoter of vitamin D target genes. The 12-helical LBD is located in the C-terminal region of the receptor and binds to $1\alpha,25(\text{OH})_2\text{D}_3$ [73]. The coregulator proteins complex with $1\alpha,25(\text{OH})_2\text{D}_3$ to activate (coactivators) or inhibit (corepressors) the target gene transcriptional activity. The coactivators mainly include the following two types: (i) Proteins like the vitamin D receptor interacting protein complex, DRIP, also known as a mediator, form a bridge that connects VDR binding to VDRE at the transcription start site (TSS) and (ii) Proteins such as the steroid receptor coactivator family 1-3 (SRC 1-3) help in transactivation [17]. VDR trans-repression involves diverse sets of mechanisms involving the suppression of gene expression by various transcription factors [74,75]. Genes get transcribed, leading to the formation of respective proteins like SLUG, which may downregulate genetic functions by the VDR [76].

The target genes of vitamin D respond in a cell-specific manner. It has been discovered that the total number of vitamin D targets in the human genome is more than one thousand. This is further supported in cells such as human lymphoblastoid and THP-1 human monocytic leukemia cells used for chromatin immunoprecipitation combined with high throughput sequencing (ChIP-seq) [77,78]. Table 2 shows some of the target genes of $1\alpha,25(\text{OH})_2\text{D}_3$, along with their specific roles and functions [42,79–81].

Table 2. Direct target genes of $1\alpha,25(\text{OH})_2\text{D}_3$, their location, and roles.

Target Gene	Cell Type	Function
<i>ASAP2</i> (<i>ArfGAP with SH3 domain, ankyrin repeat and PH domain 2</i>)	Human THP-1 cells (monocytes)	Regulates autophagy, cellular migration and vesicular transport [44]
<i>CYP24A1</i> (<i>Cytochrome P450 24A1</i>)	Kidney	Vitamin-D3 catabolizing enzyme [82]
<i>PTH1H</i> (<i>Parathyroid hormone-like hormone</i>)	Cytoplasm, golgi complex and nucleus of most cells	Activates PLC signaling pathways, proliferation of chondrocytes, regulation of bone formation by promoting recruitment

		and survival of osteoblasts, and plays a role in the physiological regulation of bone resorption [83]
<i>CAMP (Cathelicidin antimicrobial protein)</i>	Primary keratinocytes, monocytes, phagocytes, B cells and neutrophils	Antibacterial, antifungal and antiviral activities, the encoded protein functions in cell chemotaxis, immune mediator induction and inflammatory response regulation [84]
<i>Trpv6 (Transient Receptor Potential Vanilloid 6)</i>	Brush border membranes of the intestinal epithelia	Potential mediator of calcium uptake into the enterocyte [85]
<i>CYP27B1 (Cytochrome P450 27B1)</i>	Kidneys, epithelial cells, lungs, breast, intestine, stomach, endocrine glands, cells of the immune system, osteoblasts chondrocytes	Expression of 1 α -hydroxylase [44]
<i>CALB1</i>	Avian intestine and kidney and mammalian intestine, respectively	Codes for Calbindin-D28K and Calbindin-D9K proteins, upregulated by 1 α ,25(OH) $_2$ D3 [86,87]
<i>Osteocalcin</i>	Osteocytes, cartilages	Mineral deposition, bone resorption [86]
<i>FOXP3 (fork head box P3)</i>	Immune system	Maturity and performance of T regulatory cells [88]
<i>CD14 (Cluster of differentiation)</i>	Monocytes and most tissue macrophages and to a minor extent in monoblasts and promonocytes	Interacts with soluble lipopolysaccharide (LPS) released from gram-negative bacteria in combination with a plasma protein, LPS-binding protein [44]
<i>NINJ1 (Ninjurin 1)</i>	Hepatic stellate cells	Promotes axonal growth, may play a role in nerve regeneration and in the formation and function of other tissues [89]

X-ray crystallographic studies have shown that the nVDR-1 α ,25-(OH) $_2$ D complex has a classic genomic ligand binding pocket called the VDR-genomic pocket, GP and an alternative ligand binding pocket, VDR-alternative pocket, AP, that partially overlaps the VDR-GP [81]. Rochel et al. (2000) demonstrated that the VDR-GP favors a bowl-like molecular geometry with 1 α ,25(OH) $_2$ D3 by X-ray and computational studies [82]. This shape initiates genomic responses, whereas the VDR-AP accepts a ligand with a more planar-like, extended molecular geometry and initiates a rapid response [82].

The nVDR gene is frequently susceptible to polymorphisms. Polymorphism can be defined as subtle sequence variations transpiring in a population and are detected using techniques like restriction fragment length polymorphisms (RFLPs). It is a technique that exploits variations in homologous DNA sequences. Polymorphisms alter gene expression, thereby affecting protein levels of the VDR gene, leading to functional changes [83,84]. The research data indicate that VDR gene polymorphisms are associated with diabetes; cancers of the colon and prostate; autoimmune disorders like Crohn's disease, Grave's disease, and multiple sclerosis; kidney diseases; and various neurodegenerative diseases like schizophrenia [85]. The actions of 1 α ,25-(OH) $_2$ D3 are most likely due to the combined effects of both genomic and non-genomic responses [86].

3.2. Membrane VDR (mVDR)

The rapid, non-genomic activity is affected by the interaction of $1\alpha,25(\text{OH})_2\text{D}_3$ ligand with specific binding sites on the plasma membrane and vesicle membranes of target cells referred to as membrane VDR (mVDR). mVDRs are also known as 1,25 membrane-associated rapid response steroid-binding proteins (1,25-MARRS), protein disulfide isomerase family A member 3 (Pdia3), endoplasmic reticulum proteins 57 and 60 (ERp57, ERp60), and glucose-regulated protein 58 (GRP58) [87,88]. These complexes are thought to be modified VDRs. They are chaperone proteins and coupled to signal transduction systems [87]. In this review, the mVDRs are addressed by the name 1,25-MARRS.

Chick intestinal cells; adult rat and mouse cardiomyocytes, epithelial cells, myocytes, osteoblasts, chondrocytes, and cancer; and in humans, bone, teeth, brain, parathyroid, muscle, kidney, and cancer cells are 1,25D-responsive cells that function via the rapid response mechanism [88,89]. Some of the widely studied responses include calcium uptake from the intestine, insulin secretion by β -cells of the islets of pancreas, the growth and differentiation of smooth muscle cells, deviations in chondrocyte and keratinocyte growth, the changes in intracellular second messengers including Mitogen-Activated Protein Kinase (MAPK), phosphoinositides and cyclic AMP (cAMP), as well as the opening of the voltage-gated ion channels and the activation of cytosolic kinases [47,90]. Nevertheless, the rapid genomic actions are less studied, and the downstream responses following ligand-receptor binding are obscure.

1,25-MARRS are associated with membrane scaffold proteins like caveolin that are associated with lipid-rich craters of the plasma membrane and are responsible for the formation of caveolae in multiple cell types [91]. They are associated directly with the signaling pathways of the cells, using a rapid response that is elicited in the cell by the activation of the receptor complex upon ligand binding (Figure 4) [92]. Garbi et al. (2006) reported that ubiquitous ERp57 deletion in mice is embryonically lethal [93]. Wang et al. (2010) developed *Pdia3*-deficient mice to study the role of skeletal development [94]. It was revealed that no homozygous mice were observed at birth indicating that the *Pdia3* gene knockdown resulted in lethality of the homozygous mice at the embryonic stage. There were abnormalities in the skeletal tissues of *Pdia3*^{-/-} mice compared to the *Pdia3*^{+/+} mice [94]. In chick intestinal epithelial cells, targeted disruption of the 1,25-MARRS receptor gene was observed to eliminate the rapid response to $1\alpha,25(\text{OH})_2\text{D}_3$ in terms of rapid calcium uptake and Protein kinase A (PKA) signaling [95]. An association between the nVDR and 1,25-MARRS was observed in $1\alpha,25(\text{OH})_2\text{D}_3$ -mediated growth regulation of rat growth plate chondrocytes. It was also proven that chondrocytes from *nVdr*^{-/-} mice have Pdia3 and exhibit a rapid escalation in Protein kinase C (PKC) levels in response to $1\alpha,25(\text{OH})_2\text{D}_3$ [96]. These findings suggest that 1,25-MARRS has other functions in addition to acting as a receptor for $1\alpha,25(\text{OH})_2\text{D}_3$.

Caveolae are flask-shaped invaginations found on the plasma membrane and other membranes of cellular organelles, the main components of which are cholesterol, sphingolipids, and the protein, caveolin [97]. The structural components of caveolae are caveolin and exist as three isoforms, caveolin-1, -2, and -3, according to their location and function in different cells. Caveolin-1 and -2 are the most extensively studied [98]. The primary function of caveolae was thought to be vesicle trafficking, but subsequent studies have shown that they are significant partakers in cholesterol biosynthesis, signal transduction, tumor suppression, and anti-proliferative action. Caveolin-1 is found to interact with and regulate the receptor protein signaling and cellular docking mechanisms. Upon activation, they get endocytosed and translocated to the nucleus where they exert their genomic function. Recently, a loss of caveolin-1 was identified as a marker of the cancer-associated fibroblast phenotype. Caveolae are found to contain mVDR, which indicates the vital role of caveolae and caveolin proteins in VDR signaling on cell membranes [99].

Studies have linked components of cell-signaling pathways like diacylglycerol (DAG), phospholipase A2 (PLA2), and phospholipase A2-activating protein (PLAA) to mVDR, which may point to the role of caveolae in the signaling axis of $1\alpha,25(\text{OH})_2\text{D}_3$ and 1,25-MARRS by interacting with G proteins [87].

4. Signaling Pathways of Vitamin D

As mentioned earlier, recent studies have established that vitamin D and associated metabolites can evoke a rapid, non-genomic response in the cell by binding to 1,25-MARRS [87,90,91,99,100]. Numerous studies on 1,25-MARRS activated signaling pathways over the past three decades indicate that the major pathway was related to calcium influx, and is termed transcaltachia [101]. This review is focusing on a few of the most common and widely discussed signaling pathways like regulation of PKC-mediated calcium and phosphorus uptake [102], phosphatidylinositol (PI) and PLA2 activity [103], and MAPK pathway [104] and how they are involved in genomic cross-talk (Figure 5). However, there are several other signaling pathways like PKA/cAMP, protein kinase B [104], CAMK, and multiple MAPK pathways [105] that are activated by 1,25-MARRS. Detailed reports on how the various signaling pathways are affected by 1,25-MARRS can be found elsewhere [106].




Figure 5. The mechanism of action of membrane VDR and nuclear VDR. On binding of the appropriate ligand to mVDR, cellular signal transduction systems that are linked to the membrane receptor get activated, which in turn, trigger the second messengers, resulting in a rapid response. The $1\alpha,25(OH)_2D$ binds to the membrane-associated VDR and activates signaling pathways such as PKA and PKC, following which, polyisoprenyl phosphate (PIPP) levels are elevated, thereby triggering the formation of inositol triphosphate (IP3). These signaling pathways help the entry of extracellular calcium into the cells or prompt the release of calcium from intracellular stores in the endoplasmic reticulum (ER). However, binding of $1\alpha,25(OH)_2D$ to the canonical nVDR causes a genomic response by initiating the transcription of targeted genes. Nemere et al. (2004), reported that 1,25-MARRS have a similar affinity for the ligand as that of the nVDR, but, the membrane-associated

protein is 6–10 times more abundant in the cells than the nuclear receptor [107]. 1,25-MARRS is usually found associated with caveolin proteins. The intracellular Ca^{2+} levels are enhanced on binding of $1\alpha,25(\text{OH})_2\text{D}$ to 1,25-MARRS. A study in keratinocytes showed that binding of $1\alpha,25(\text{OH})_2\text{D}_3$ to the membrane receptor resulted in elevated metabolism of phosphatidylinositol (PI) to phosphatidylinositol triphosphate (PIP₃), resulting in increased levels of IP₃ in the cells [108,109]. The rise in IP₃ were in accordance with a rise in calcium levels, eliciting a rapid response within 2–5 min [50]. Calcium is released from ER storage pools or through the transmembrane trafficking of calcium through the membrane calcium channels [110].

PKC activation is observed in chondrocytes by binding of $1\alpha,25(\text{OH})_2\text{D}_3$ to 1,25-MARRS [111]. The receptor-ligand binding results in a rapid increase in the amount of PLA₂ activity. PLA₂ activates PKC with the help of several intermediate products, the key one being arachidonic acid. Arachidonic acid can directly activate PKC, while serving as a substrate for several other enzymes. Lysophospholipids are intermediates of arachidonic acid release. They are capable of activating phospholipase C, which hydrolyzes PI to DAG and IP₃. IP₃ binds to receptors that are located on the smooth endoplasmic reticulum, causing Ca^{2+} release, which acts as a co-factor for PKC α . DAG binds to PKC α , activating the enzyme and translocating it to the plasma membrane [112]. Richard et al. first demonstrated the involvement of 1,25-MARRS in anti-cancer activity in breast cancer cells [101]. $1\alpha,25(\text{OH})_2\text{D}_3$ is thought to inhibit the phosphorylation and expression of Akt, a kinase responsible for regulating cell survival and proliferation [113]. $1\alpha,25(\text{OH})_2\text{D}_3$ promotes angiogenic factor interleukin-8 (IL-8) in squamous carcinoma cells, but calcitriol interrupts IL-8 signaling in prostate cancer cells and inhibits the migration and tube formation of endothelial cells [114,115]. Hilliard and coworkers in 1994 reported that treatment of COS-1 cells with $1\alpha,25(\text{OH})_2\text{D}_3$ caused an increase in the phosphorylation of serine residue 208 (S208) located in the VDR, thus recruiting the coactivator to form the nVDR-RXR heterodimer [116]. On the other hand, antagonistic functions of genomic and rapid response pathways have been observed in breast cancer cells [117].

Cross-talk exists between the genetic and rapid signaling pathways. This is brought about by phosphorylation of the chief proteins in the VDR-transcriptional complex and is often observed in cell-cycle regulation. The different modes of signaling actions and cross-talk operated by $1\alpha,25(\text{OH})_2\text{D}_3$ -1,25-MARRS association lead to transcriptional modulation of several genes, depending on the cell- or tissue-type considered. Barletta and coworkers (2002) reported that $1\alpha,25(\text{OH})_2\text{D}_3$ -1,25-MARRS mediated rapid activation of cytosolic kinases and may be responsible in phosphorylating key coactivators involved in diverse pathobiological and physiological processes [118]. The diverse biological effects result from genomic or non-genomic cross-talk to some extent as well as through context-specific functions of the target genes. Certain second messengers may engage in cross-talk with the nucleus to regulate gene expression, particularly the Rapidly Accelerated Fibrosarcoma / Mitogen-Activated Protein Kinase (RAF/MAPK) pathway [119]. Deeb et al. (2007) reported the activation of MAPK–ERK (extracellular signal-regulated kinase) 1 and 2 cascades through the phosphorylation and activation of RAF by PKC, partly induced by the changes in intracellular Ca^{2+} concentration [120].

5. Regulation of Calcium and Phosphate Homeostasis by $1\alpha,25(\text{OH})_2\text{D}_3$ and PTH

An essential function of $1\alpha,25(\text{OH})_2\text{D}_3$ is the maintenance of serum calcium and phosphate levels. Normal serum Ca^{2+} and PO_4 levels serve as the cornerstones for several biological processes. $1\alpha,25(\text{OH})_2\text{D}_3$ helps in effective utilization of calcium by enhancing intestinal Ca^{2+} absorption or Ca^{2+} mobilization from bones in the absence of calcium in the diet. It acts together with the parathyroid hormone (PTH) to help maintain mineral homeostasis between the blood and bones, the neuromuscular and immune function, and cardiovascular health, and is essential in the suppression of malignant changes [121].

Plasma Ca^{2+} and PO_4 concentrations are maintained at constant levels of 2.2–2.6 mMol/L and 0.8–1.45 mMol/L, respectively [121]. A drop in serum Ca^{2+} levels stirs a series of quick reactions in the biological system, illustrated in Figure 6.




Figure 6. The regulation of mineral homeostasis by parathyroid hormone (PTH) and $1\alpha,25(\text{OH})_2\text{D}_3$. The physiological functions of PTH and $1\alpha,25(\text{OH})_2\text{D}_3$ are activated when serum calcium levels drop. The hormones act in conjunction with each other and exert coordinated effects on the kidneys, bones, and intestine to increase Ca^{2+} levels to normal. There is bone resorption, increased calcitriol formation by the kidneys and decreased calcium excretion from urine, and increased Ca^{2+} absorption by the intestine. Upon achievement of homeostasis, the process is shut down by a negative feedback loop, which is initiated by calcitonin secreted by the thyroid gland. Thus, the combined effect of PTH and $1\alpha,25(\text{OH})_2\text{D}_3$ is necessary to maintain mineral homeostasis.

The parathyroid glands possess transmembrane calcium-sensing proteins that are coupled to a G protein system and detect the levels of Ca^{2+} in the plasma [122]. This stimulates the secretion of PTH [123]. PTH advances to the osteoblasts in bones and the proximal convoluted tubule cells in the kidney within seconds. PTH stimulates the mobilization of Ca^{2+} from the bones and reduces its excretion by the kidney. These brisk effects increase the concentration of Ca^{2+} in the blood. In conjunction with these changes, PTH promotes the activity of the CYP27B1 gene in the proximal convoluted tubule cells of the kidney [124]. Therefore, the proximal convoluted tubule cells of the kidney act as the endocrine gland of the vitamin D hormone [125]. $1\alpha,25(\text{OH})_2\text{D}_3$ has a quintessential role in Ca^{2+} level homeostasis. $1\alpha,25(\text{OH})_2\text{D}_3$ signals the intestine to augment Ca^{2+} absorption and in association with PTH, promotes mobilization of bone Ca^{2+} and PO_4 and renal reabsorption of minerals. Consequently, this leads to the deposition of more Ca^{2+} and PO_4 in the system, which boosts both mineral levels in the serum [122]. When the levels of serum Ca^{2+} exceed the set point of the

calcium-sensing system, the parathyroid gland-induced cascade of events is shut down. The C-cells present in the thyroid gland secrete calcitonin in response to the high serum Ca^{2+} levels. Calcitonin is a 32-amino acid peptide that blocks Ca^{2+} mobilization from the bone [126]. Calcitonin also stimulates the renal CYP27B1 to provide $1\alpha,25(\text{OH})_2\text{D}_3$ for non-calcemic needs under normal Ca^{2+} conditions [127].

When the Ca^{2+} levels return to normal, the 24-hydroxylase enzyme in the kidney is stimulated, which breaks down the active $1\alpha,25(\text{OH})_2\text{D}_3$ to its inactive form. The enzyme also breaks down $25(\text{OH})\text{D}_3$, which is the form of vitamin D that is stored in the body.

The increase in the synthesis of $1\alpha,25(\text{OH})_2\text{D}_3$ in response to hypocalcemia (via the secretion of parathyroid hormone) occurs within a few hours, whereas the parathyroid glands respond to hypocalcemia within a few minutes. Under hypercalcemic conditions, the secretion of PTH is suppressed, and the synthesis of $1\alpha,25(\text{OH})_2\text{D}_3$ is significantly reduced, whereas the synthesis of another inactive metabolite, 24R, 25-dihydroxyvitamin D3 ($24(\text{R}),25-(\text{OH})_2\text{D}_3$), destined for excretion is increased [128].

6. Vitamin D and the Gastric Tissue

Notably, in the last four decades, non-calcemic effects of $1\alpha,25(\text{OH})_2\text{D}_3$ have been explored and investigated. Studies have revealed that $1\alpha,25(\text{OH})_2\text{D}_3$ as well as its metabolites play a key role in controlling and regulating genes responsible for preserving the integrity of the epithelial barrier including the GI tract, in addition to immunoregulatory and inflammatory responses [18]. One of the best studied non-traditional functions of $1\alpha,25(\text{OH})_2\text{D}_3$ is the cellular mechanisms involved in tumor control and repression of proliferating epithelial cells of the skin [129], breast [130,131], colon [132], and prostate [133]. Therefore, the majority of the studies have focused on tumor cells and their progenitors along with the tumor microenvironment (TME) [134].

There is a need for further research to study the roles of $1\alpha,25(\text{OH})_2\text{D}_3$ and its metabolites in maintaining the equilibrium and stability of normal tissue. Despite the increase in the number of studies, publications, and summits centering on the various utilities of $1\alpha,25(\text{OH})_2\text{D}_3$, there is little awareness regarding the metabolites and precise functions of vitamin D in the gastric environment [135]. Scientists, clinicians, and dieticians have started to explore the therapeutic advantages of this steroid hormone in gastric cancer (GC) coupled with its role in the TME.

This review is an attempt to assemble a few of the important articles that presents information regarding the regular expression and function of vitamin D and the VDR in the gastric tissues.

Furthermore, the authors were interested in studying the role of vitamin D and the VDR in the normal proliferation and differentiation of the gastric cells as well as in maintaining the cellular and physiological homeostasis of the gastric tissues. A systematic literature review was conducted using the electronic search engines and online databases like Cochrane Library, Google Scholar, PubMed Central, Semantic Scholar, Scopus, ScienceDirect, SciFinder, and RefSeek. The databases were explored for articles published in the English language between the years 1927 and 2019, using the keywords, vitamin D, calcitriol, stomach and gastric environment and gastric tissue. However, the articles focusing solely on the role of vitamin D and its analogues in gastric progenitor cell proliferation and the differentiation and formation of gastric epithelium were challenging to acquire. Nevertheless, a few articles that are related to this area to some extent are included in the study. This endeavor warrants the scrutinization of the central role of the VDR in $1\alpha,25(\text{OH})_2\text{D}_3$ action and the therapeutic roles of $1\alpha,25(\text{OH})_2\text{D}_3$ and its analogs in maintaining gastric tissue homeostasis and controlling tumor proliferation, and thereby in the betterment of human health. Table 3a,b indicate the major available studies that emphasize critical findings in the human and animal stomachs. Table 3a,b indicate the major available studies that emphasize key findings in the human and animal stomachs.

Table 3. A list of publications in humans and animals between the years, 1927 and 2019. (a) Major findings from animal studies; (b) Major findings from human studies.

(a)

No.	References	Organism Studied	Form of vitamin D Used in the Study	Key Findings	Ref. No.
1	Sato K. (1993)	Rats	$1\alpha,25(\text{OH})_2\text{D}_3$	Elevated levels of $1\alpha,25(\text{OH})_2\text{D}_3$ when administered to hypercalcemic patient with PTHrP-producing gastric carcinoma, aggravated the severity of malignancy-associated hypercalcemia (MAH).	[110]
2	Leigh-Clare J. (1927)	Australasian Petrel	Vitamin D	One of the first articles to describe the presence of Vitamin D in stomach oil. The study was an attempt to elucidate the source of vitamin D in the oil.	[136]
3	Selye H. and Bois (1957)	Sprague-Dawley rats	Ergocalciferol	VDT lead to calcium deposition in the muscularis of rat stomachs	[137]
4	Stumpf W. (1979)	Rats	$1\alpha,25(\text{OH})_2\text{D}_3$ and its metabolites	Vitamin D receptors for $1\alpha,25(\text{OH})_2\text{D}_3$ or its metabolites target tissues of the GIT including the nuclei of some of the cells of the stomach	[138]
5	Kirui N. (1981)	Rats	$1\alpha,25(\text{OH})_2\text{D}_3$	$1\alpha,25(\text{OH})_2\text{D}_3$ has a direct role on hypercalcemia and STM with lesions forming in most of the soft tissues, including the gastric glandular mucosa and muscularis	[139]
6	Selking O. (1982)	Rats	Ergocalciferol	Vitamin D induced hypercalcemia in parathyroidectomized rats is associated with a thickened gastric mucosa, but the serum gastrin, number of gastrin (G) cells or antral gastrin remained unchanged	[140]
7	Kurose, T. (1988)	Rats	$1\alpha,25(\text{OH})_2\text{D}_3$	Calcium and $1\alpha,25(\text{OH})_2\text{D}_3$ deficiency impair gastrin and somatostatin secretion in the perfused rat stomach	[141]
8	Axelsson J. (1991)	Rats	$1\alpha,25(\text{OH})_2\text{D}_3$	Gastrectomy may lead to an increase in the levels of serum $1\alpha,25(\text{OH})_2\text{D}_3$, resulting in enhanced absorption of calcium by the small intestine. This may be the cause for diseases like osteomalacia in the gastrectomized patients.	[142]
9	Holick M. (1995)	Rats, mice, humans	$1\alpha,25(\text{OH})_2\text{D}_3$	Non-calcemic tissues including the gonads, pituitary gland, thymus, pancreas, stomach, breast, and skin possess the nuclear receptors for $1\alpha,25(\text{OH})_2\text{D}_3$, targeting these receptors with analogs of $1\alpha,25(\text{OH})_2\text{D}_3$ provides treatment against different diseases.	[143]
10	Stumpf W. (1995)	Two-month old mice	$1\alpha,25(\text{OH})_2\text{D}_3$ and its analogue 22-Oxacalcitriol (OCT)	Autoradiographic studies showed nuclear concentration and retention of $1\alpha,25(\text{OH})_2\text{D}_3$ and its analogue 22-Oxacalcitriol (OCT) in neck mucus cells of gastric and pyloric glands and	[144]

				in dispersed endocrine cells in the antrum	
11	Ikezaki S. (1996)	Male Wistar rats	24R,25(OH)2D3	The development of atypical hyperplasias and adenocarcinomas in the glandular stomachs was decreased by exposure to 24R,25(OH)2D3, which shows that 24R,25(OH)2D3 has chemopreventive effects.	[145]
12	Gagnemo-Persson R. (1999)	Male Sprague-Dawley rats	Ergocalciferol / Vitamin D2	Gastrin-ECL-cell axis can be suppressed by vitamin D or by vitamin D-dependent mechanisms. Also, vitamin D receptor gene expression was seen in the rat oxyntic mucosa.	[146]
13	Stumpf W. (2008)	Rats, mice, hamsters and zebra finch	1 α ,25(OH) ₂ D3	Autoradiography studies confirmed the binding of 1 α ,25(OH) ₂ D3 and its oxygen analog OCT in numerous regions of the digestive tract	[147]
14	Häkkinen I. and Lindgren I. (2009)	Albino rats	1 α ,25(OH) ₂ D3	Excess of 1 α ,25(OH) ₂ D3 leads to calcification of gastric tissues	[148]
15	Sahin H. (2018)	Rats	1 α ,25(OH) ₂ D3	1 α ,25(OH) ₂ D3 protects the gastric mucosa via attenuation of inflammatory reaction, oxidative stress and apoptosis.	[149]

(b)

No.	First Author, Year	Form of Vitamin D Used in the Study	Key Findings	Ref. No.
1	Paterson C. (1965)	1 α ,25(OH) ₂ D3	Gastrectomy and other surgeries of the stomach can trigger osteomalacia due to the increased levels of serum vitamin D which results in increased absorption of calcium by the small intestine.	[150]
2	Kimura K. (1967)	1 α ,25(OH) ₂ D3	Calcium deposits were found in the soft tissues of a patient who died of uremia after administration of toxic levels of Vitamin D for 6 months	[151]
3	Lawrence, W. (1977)	1 α ,25(OH) ₂ D3	Inadequate absorption of Vitamin D in patients who underwent gastrectomy is due to steatorrhea which may contribute to osteomalacia later on.	[152]
4	Giovannucci E. (2006)	1 α ,25(OH) ₂ D3	Black men with poor vitamin D status are highly susceptible for digestive system cancer and have a higher mortality rate.	[153]
5	Chen W. (2007)	25(OH)D	A direct association between higher serum 25(OH)D concentration and increased risk of oesophageal squamous cell carcinomas (ESCC) in men but not women but no association with risk of gastric cardia or noncardia adenocarcinoma in either sex.	[154]
6	Pan L. (2010)	1 α ,25(OH) ₂ D3	Studies in HGC-27 adenocarcinoma cells showed that Vitamin D can be used in gastric cancer therapies in association with trichostatin A/sodium butyrate and 5-aza-2-deoxycytidine	[155]
7	Antico A. (2012)	1,25(OH) ₂ D	Low vitamin D concentration in <i>H. pylori</i> gastritis patients might lead to a more severe Th1-type aggression to the stomach epithelium	[156]

8	Park M.R. (2012)	Paricalcitol (19-nor-1,25-(OH) ₂ D ₂)	Paricalcitol, has anticancer activity on GC cells by regulating cell cycle, apoptosis and inflammation	[157]
9	Kopic S. (2013)	1 α ,25(OH) ₂ D ₃	Gastric acid, gastrin secretion is increased in humans under conditions of hyperparathyroidism with the combined actions of both PTH and vitamin D	[158]
10	Guo L. (2014)	1 α ,25(OH) ₂ D ₃	VDR and CAMP expression is up-regulated in the gastric epithelium during <i>H. pylori</i> infection; thus, VDR is important in maintaining the homeostasis of gastric mucosa and protection from <i>H. pylori</i> infection.	[159]
11	Wen Y. (2015)	1,25(OH) ₂ D ₃	VDR expression was seen to decline from the premalignant stage, to low expression in gastric cancer tissues. VDR could be a potential prognostic factor for patients with gastric cancer.	[160]
12	Bashir M. (2016)	1 α ,25(OH) ₂ D ₃	Vitamin D influences the composition of gut microbiome.	[161]
13	Vyas N. (2016)	25(OH)D	There is a positive relationship between VDD and gastric adenocarcinoma	[162]
14	Du C. (2017)	1 α ,25(OH) ₂ D ₃	Vitamin D supplement might be a safe and economical way to prevent or treat gastric cancer	[163]
15	Yildirim O. (2017)	25(OH)D	Deficiency of 25(OH)D may be a risk factor related to eradication failure of <i>H. pylori</i> , and supplementation of 25(OH)D before eradication of <i>H. pylori</i> may provide better results.	[164]
16	El Shahawy M.S. (2018)	25(OH)D ₃	25(OH)D deficiency can be a risk factor in eradication failure of <i>H. pylori</i> infection.	[165]
17	Fletcher J. (2019)	1 α ,25(OH) ₂ D	1 α ,25(OH) ₂ D protects the gastrointestinal barrier and evokes immune responses	[166]

One of the first studies regarding 1 α ,25(OH)₂D in the stomach was conducted by Leigh Claire in 1927. The study attempted to explicate the source of 1 α ,25(OH)₂D in the oil present in the stomach of fulmar petrel (*Australata lessoni*), also known as muttonbird. The main diet of fulmar petrel and Australasian cod includes whale feed which consists principally of a bright red crustacean. It was earlier thought that the stomach oil was a residue from the digestion of whale feed, but an analysis of the stomach oil showed an abundance in cetyl esters, which are relatively absent in whale feed. The study showed that rats fed with muttonbird stomach oil were cured of rickets. This confirmed the presence of 1 α ,25(OH)₂D in the oil. However, the source of 1 α ,25(OH)₂D in the stomach oil remained unknown, even though it was suggested that whale feed is the common source of 1 α ,25(OH)₂D in stomach oil and cod-liver oil [136].

The field received a spur when the toxic effects of 1 α ,25(OH)₂D₃ were described in various publications as early as 1928 [138]. Hypervitaminosis of 1 α ,25(OH)₂D₃ causes pathological modifications such as calcification in the skeletal and soft tissues like the stomach and kidneys [167]. One such study published by Dewind in 1961 stated that excessive ingestion of 1 α ,25(OH)₂D₃ might contribute to osteosclerosis [168]. The pathological effects of ergocalciferol on gastric tissues were reported by Selye and Bois (1957). They demonstrated that a subcutaneous injection of ergocalciferol simultaneously with cortisol acetate in female Sprague-Dawley rats caused calcification of the tissues of the gastric mucosa, but steroids like deoxycorticosterone acetate inhibited this effect [137]. Another study in albino rats treated with 1 α ,25(OH)₂D₃, dihydrotachysterol (AT10) and parathyroid extract showed that overdosage of these substances could lead to metastatic calcification of the stomach tissues as well as the gastric mucosa [148].

Kimura et al. (1967) reported an autopsy case of hypervitaminosis D where the patient died due to calcium deposition in bones and soft tissues including the stomach and esophagus, subsequently leading to uremia [151]. Subsequently, several researchers have studied the toxic effects of hypervitaminosis D and hypercalcemia in soft-tissue mineralization (STM), which induces lesions in

the proximal and cortico-medullary regions of the renal tubules, aortic media and intima, myocardial arterioles and myofibers, and gastric glandular mucosa and muscularis [139]. The 1960s witnessed fundamental research on the synthesis, metabolism, and clinical relevance of $1\alpha,25(\text{OH})_2\text{D}_3$. It was found that a deficiency of $1\alpha,25(\text{OH})_2\text{D}_3$ can lead to hepatic and gastrointestinal diseases that ultimately affect the bone [84].

The VDR was discovered in 1969, and it was established that the VDR was a cell nucleus-localized receptor [169]. It was found that the VDR is present in over 30 target tissues in humans comprising the bone, cartilage, intestine and kidney, as well as in pancreatic β -cells [170], hair follicles [43], various cancer cells [119], and B and T lymphocytes [171]. With the advent of new techniques, Stumpf and coworkers (1979 and 2008) and Holick (1995) established that rat duodenum, jejunum, ileum, colon, stomach [143], pituitary, parathyroid, kidney [172], and skin are target tissues of $1,25(\text{OH})_2\text{D}_3$ [147]. Autoradiographic studies by Stumpf et al. (1995) indicated the nuclear binding of radioactively labeled $1,25(\text{OH})_2\text{D}_3$, and its analog, 22-Oxacalcitriol, in neck mucus cells of gastric and pyloric glands and dispersed endocrine cells in the antrum [144]. Moreover, the *Vdr* gene expression was found in the oxyntic mucosa of rats [146]. These discoveries assisted in correlating a $1\alpha,25(\text{OH})_2\text{D}_3$ deficiency and fat malabsorption to partial gastrectomy [82,173,174]. Gastric bypass surgery (GBP) or sleeve gastrectomy is a common and widely advised medical treatment (surgery) to overcome morbid obesity by achieving significant weight loss and improving associated medical conditions [175]. Despite the beneficial effects of GBP, the complications such as nutritional deficiencies, metabolic bone disease (MBD), and osteoporosis have been reported. Post-gastrectomy MBD is a distinct condition associated with gastrectomy, due to calcium and $1\alpha,25(\text{OH})_2\text{D}_3$ malabsorption, and includes subclinical osteomalacia symptoms [176,177]. A possible explanation provided by Axelson for this trend is that an escalation of serum $1\alpha,25(\text{OH})_2\text{D}$ levels in such patients may enhance the ability of the small intestine to absorb calcium [142]. Rino et al. points to less food intake and fat malabsorption to be potential causes of the deficiency of fat-soluble vitamins in general post-gastrectomy, but no significant reduction in serum vitamin D levels was seen [178]. Several case studies have reported attempts to examine and investigate the pathophysiology of osteomalacia and osteoporosis following gastrectomy [150,152].

The scientific world witnessed novel endeavors to uncover the numerous mechanisms of $1\alpha,25(\text{OH})_2\text{D}_3$ responsible for maintaining gastric homeostasis. A major objective is to unearth a relationship between calcium or phosphate homeostasis and gastric secretions, like gastric acid and gastrin hormone. Gastric secretions and homeostasis are maintained by highly meticulous and precise mechanisms [179]. Acid-secretion in the stomach is guarded by a well-schemed machinery comprising hormonal, neuronal, and paracrine modules. A remarkable study in this subject was conducted by Selking O. et al. (1982), which indicates that hypercalcemia does not cause an increase in gastrin secretion in rats, and further reinforces previous findings [140,150,152,179]. However, follow-up investigations revealed more than a few unpredictable findings. Kurose, T. et al. (1988) and Gagnemo-Persson (1999) demonstrated that calcium and $1\alpha,25(\text{OH})_2\text{D}_3$ -deficiency impairs gastrin and somatostatin secretion in the perfused rat stomach [141,146,180].

Enterochromaffin-like (ECL) cells constitute one of the largest endocrine cell populations in the mammalian system. They are controlled by the gastrin hormone and are rich in histamine, calcium-binding protein calbindin, and chromogranin A. Gastrin and the ECL cells form a functional unit, known as the gastrin–ECL-cell axis [181]. Gagnemo-Persson (1999) reported that $1\alpha,25(\text{OH})_2\text{D}_3$ or $1\alpha,25(\text{OH})_2\text{D}_3$ -dependent mechanisms play a direct role in suppressing the gastrin–ECL-cell axis [146]. Kopic and Geibel (2013) investigated the effect of calcium on gastric physiology. Serum calcium levels are controlled by the combined actions of PTH and $1\alpha,25(\text{OH})_2\text{D}_3$ [158]. The minute variations in calcium homeostasis can have adverse effects. The homeostasis is maintained by a highly sensitive calcium-sensing receptor (CaSR) present on the parathyroid gland in addition to tissues, like the kidney and gastrointestinal tract (GIT) along with a negative feedback loop by $1,25(\text{OH})_2\text{D}_3$ [158]. In humans, calcium stimulates gastric acid secretion. In diseased states like hyperparathyroidism, PTH influences gastric cells and gastrin release with the aid of calcium [158,182]. The studies revealed controversial results regarding the roles of $1,25(\text{OH})_2\text{D}_3$ and its impact on gastric physiology.

Studies have demonstrated an association between low levels of $1\alpha,25(\text{OH})_2\text{D}$ and related complications in people with highly melanized skin. Harris in 2006 reported that a high percentage of American black people have lower serum $25(\text{OH})\text{D}_3$ than their white counterparts [183]. Giovannucci et al. (2006) conducted a study in a cohort of black and white male health professionals that were relatively homogenous in education and living standards [153]. The results indicated that black people are at a higher risk of digestive system cancers associated with hypovitaminosis D, and thus a higher mortality rate [153]. Several studies have hypothesized that the nutritional status and serum levels of $1\alpha,25(\text{OH})_2\text{D}_3$ might influence the chances of GC. Chen et al., (2007) observed that higher concentrations of serum $25(\text{OH})\text{D}_3$ intensifies the hazard of esophageal squamous cell carcinomas in males but not in females, but no association was found in the case of the gastric cardia or non-cardia adenocarcinoma in either sex [154].

Gastric ulcers are caused by a disruption in the homeostasis between the acidic and protective environment of the gastric mucosa [161,184]. The ulceroprotective and antioxidant effects of $1\alpha,25(\text{OH})_2\text{D}_3$ were reported in rats [149]. *H. pylori* gastritis infection, which is the Th1 type, causes autoimmune gastritis (AIG), and is classified as an autoimmune disease. The low serum $1\alpha,25(\text{OH})_2\text{D}_3$ status in *H. pylori* gastritis patients could possibly act as a predisposing factor for the severe Th1-type AIG, which causes intense damage to the stomach epithelium [156]. The gut microbiome is reactive to $1\alpha,25(\text{OH})_2\text{D}_3$ supplementation and deficiency despite considerable heterogeneity [164,165]. For example, $1\alpha,25(\text{OH})_2\text{D}_3$ could boost the intracellular killing of the replicating *H. pylori* bacteria [159,185]. Fletcher et al., (2019) reported the importance of $1\alpha,25(\text{OH})_2\text{D}_3$ in maintaining the gastrointestinal barrier integrity, surveillance of the gut microbiota, and inflammatory immune responses [166].

Du et al. reported that $1\alpha,25(\text{OH})_2\text{D}_3$ and the VDR interaction can induce a cascade of gene regulation and cell signaling reactions to promote anti-tumor mechanisms [163]. Vitamin D and its analogues play an important role in reducing the risk of gastric adenocarcinomas. Expanding evidences have indicated that prevention of vitamin D deficiency is an economic and promising way to reduce risk of GC [157,162]. In 2015, Wen et al., studied the expression of local VDR in normal, premalignant, and malignant gastric tissues by immunohistochemical techniques [160]. The relationship between the VDR and clinicopathological factors of GC patients was also analyzed. They reported that the VDR was dynamically expressed in various types of gastric tissues. On treatment with $1\alpha,25(\text{OH})_2\text{D}_3$, a decline in the VDR expression was observed from pre-malignant to gastric cancer tissues, and the VDR expression was the least in poorly differentiated tissues [160]. This shows that the VDR can be a prospective prognostic factor for GC. Ikezaki (1996) reported that the development of atypical hyperplasias and adenocarcinomas was significantly diminished in the glandular stomachs of rats by exposure to $24\text{R},25(\text{OH})_2\text{D}_3$, which shows that $24\text{R},25(\text{OH})_2\text{D}_3$ has chemopreventive effects [145].

Epigenetic mechanisms like chromatin remodeling complexes, DNA methylation, miRNA, and histone modifications are associated with GC [186]. Chromatin remodeling is capable of altering the chromatin structure, which disrupts the competence of regulatory proteins from gaining access to the target sites. Histone modifications, like acetylation, are vital in transcription and are synchronized by a balance between the different activities of histone acetyltransferases and histone deacetylases (HDACs) [187]. This process thereby amends the interaction of regulatory proteins with their targets and correlates with the transcriptional status of the genes [188]. Trichostatin A (TSA) is a known potent inhibitor of HDACs. TSA reportedly has therapeutic effects as it inhibits HDACs with zinc-containing catalytic sites, leading to the accumulation of acetylated histones in the nucleus and subsequent activation of target genes [189]. DNA methylation involves the addition of a methyl group to cytosine residues by enzymes called DNA methyltransferases (DNMTs). It is associated with transcriptional silencing. The presence of 5-Aza-2'-deoxycytidine (5azadC), a cytidine analog in DNA, prevents DNA methylation [155,190,191]. In association with TSA/sodium butyrate (TSA/NaBu) and 5-aza-2-deoxycytidine (5-Aza), $1\alpha,25(\text{OH})_2\text{D}_3$ can induce apoptosis in GC cells through the expression of phosphatase and tensin homolog deleted on chromosome 10 (PTEN), which is an

important tumor suppressor gene. This can lead to the potential application of $1\alpha,25(\text{OH})_2\text{D}_3$ as a novel molecular target in GC therapies in association with the usage of TSA/NaBu and 5-Aza [155].

7. Conclusions

Unfortunately, for the gastroenterologist, there are limited data available on the impacts of VDD associated with gastric disorders. Considering the main roles of $1\alpha,25(\text{OH})_2\text{D}_3$ in the immunoregulatory and preservation of the morphology and physiology of tissues, it is worth investigating the importance of $1\alpha,25(\text{OH})_2\text{D}_3$ and its metabolites in maintaining stomach homeostasis. The information from research can be used for public awareness, recommending it as a necessary dietary supplement in daily lives thereby, improving human health. In the past 30–40 years, researchers have revealed an in-depth and detailed mechanism of how this miracle hormone facilitates various physiological processes in many biological systems. However, extensive research, both basic and applied, is required to explain the role of $1\alpha,25(\text{OH})_2\text{D}_3$ and its metabolites in the gastric environment. If scientists, and the agencies that fund them, direct their efforts and expertise to focus on the role of this vitamin, better ways to treat abnormalities arising in the gastric tissues can be anticipated. These attempts can make more efficacious and serious use of $1\alpha,25(\text{OH})_2\text{D}_3$, its metabolites, and synthetic analogs, which can cause a significant paradigm shift in the progress of medical science and the welfare of human and animal populations.

Author Contributions: A.A.M. and I.S. initiated the study. The review first draft was written by S.S. with contributions from A.A.M. and I.S. All three authors contributed significantly to materialize the manuscript, and all authors contributed to the study design, preparation of the manuscript, and have read and approved the final version.

Funding: This research was funded by Zayed bin Sultan Center for Health Sciences, UAE University, grant number 31R138.

Acknowledgments: We would like to express our gratitude to Zayed bin Sultan Center for Health Sciences, UAE University, for funding and supporting the project.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

$1\alpha,25(\text{OH})_2\text{D}_2$	$1\alpha,25$ -dihydroxyvitamin D2
$1\alpha,25(\text{OH})_2\text{D}_3$	$1\alpha,25$ -dihydroxyvitamin D3
<i>1,25-MARRS</i>	1,25 membrane-associated rapid response steroid-binding proteins
$24(\text{R}),25-(\text{OH})_2\text{D}$	24R, 25-dihydroxyvitamin D
$25(\text{OH})\text{D}$	25-hydroxyvitamin D
$25(\text{OH})\text{D}_3$	25-hydroxyvitamin-D3
<i>5azadC</i>	5-Aza-2'-deoxycytidine
<i>ASAP2</i>	ArfGAP with SH3 domain, ankyrin repeat and PH domain 2
<i>cAMP</i>	Cyclic AMP
<i>CaSR</i>	Calcium-sensing receptor
<i>CAMP</i>	Cathelicidin antimicrobial protein
<i>CLIA</i>	Chemiluminescent immunoassay
<i>CYP</i>	Cytochrome P450
<i>DAG</i>	Diacylglycerol
<i>DBP</i>	Vitamin D binding protein
<i>DNMT</i>	DNA methyltransferase
<i>ECL</i>	Enterochromaffin-like cells
<i>ELISA</i>	Enzyme-linked immunoassay

<i>ER</i>	Endoplasmic reticulum
<i>ERp57</i>	Endoplasmic reticulum proteins 57
<i>ERp60</i>	Endoplasmic reticulum proteins 60
<i>FGF23</i>	Fibroblast growth factor-23
<i>FOXP3</i>	Fork head box P3
<i>GBP</i>	Gastric bypass surgery
<i>GC</i>	Gastric cancer
<i>GIT</i>	Gastrointestinal tract
<i>GRP58</i>	Glucose-regulated protein 58
<i>HDAC</i>	Histone deacetylases
<i>HPLC</i>	High-performance liquid chromatography
<i>IL-8</i>	Interleukin-8
<i>IP3</i>	Inositol triphosphate
<i>LC-MS/MS</i>	Liquid Chromatography-Tandem Mass Spectrometry
<i>LBD</i>	Ligand binding domains
<i>mVDR</i>	Membrane VDR
<i>MAH</i>	Malignancy-associated hypercalcemia
<i>MAPK</i>	Mitogen-Activated Protein Kinase
<i>MBD</i>	Metabolic bone disease
<i>NINJ1</i>	Ninjurin 1
<i>nVDR</i>	Nuclear VDR
<i>Pdia3</i>	Protein disulfide isomerase family A member 3
<i>PI</i>	Phosphatidyl inositol
<i>PIP3</i>	Phosphatidylinositol triphosphate
<i>PIPP</i>	Polyisoprenyl phosphate
<i>PKA</i>	Protein kinase A
<i>PKC</i>	Protein kinase C
<i>PLA2</i>	Phospholipase A2
<i>PLAA</i>	Phospholipase A2-activating protein
<i>PTEN</i>	Phosphatase and tensin homolog deleted on chromosome 10
<i>PTH</i>	Parathyroid hormone
<i>PTH LH</i>	Parathyroid hormone like hormone
<i>RAF/MAPK</i>	Rapidly Accelerated Fibrosarcoma / Mitogen-Activated Protein Kinase
<i>RIA</i>	Radioimmunoassay
<i>SRC 1-3</i>	Steroid receptor coactivator family 1-3
<i>STM</i>	Soft-tissue mineralization
<i>TME</i>	Tumor microenvironment
<i>Trpv6</i>	Transient Receptor Potential Vanilloid 6
<i>TSA</i>	Trichostatin A
<i>UVB</i>	Ultraviolet B
<i>VDD</i>	Vitamin D deficiency
<i>VDR</i>	Vitamin D receptor
<i>VDRE</i>	Vitamin D response element

VDT Vitamin D Toxicity

References

1. Sahota, O. Understanding vitamin D deficiency. *Age Ageing* **2014**, *43*, 589–591.
2. McKenna, M.J.; Murray, B.F.; O’Keane, M.; Kilbane, M.T. Rising trend in vitamin D status from 1993 to 2013: Dual concerns for the future. *Endocr. Connect.* **2015**, *4*, 163–171.
3. Kuchuk, N.O.; Pluijm, S.M.F.; Van Schoor, N.M.; Looman, C.W.N.; Smit, J.H.; Lips, P. Relationships of Serum 25-Hydroxyvitamin D to Bone Mineral Density and Serum Parathyroid Hormone and Markers of Bone Turnover in Older Persons. *J. Clin. Endocrinol. Metab.* **2009**, *94*, 1244–1250.
4. Holick, M.F. Vitamin D Deficiency. *N. Engl. J. Med.* **2007**, *357*, 266–281.
5. Chan, J.; Jaceldo-Siegl, K.; Fraser, G.E. Serum 25-hydroxyvitamin D status of vegetarians, partial vegetarians, and nonvegetarians: The Adventist Health Study-21234. *Am. J. Clin. Nutr.* **2009**, *89*, 1686S–1692S.
6. Looker, A.C.; Bowman, B.A.; Hollis, B.W.; Gillespie, C.; Allen, C.; Dougherty, C.; Gunter, E.W.; Cogswell, M.E.; Nesby-O’Dell, S.; Scanlon, K.S. Hypovitaminosis D prevalence and determinants among African American and white women of reproductive age: Third National Health and Nutrition Examination Survey, 1988–1994. *Am. J. Clin. Nutr.* **2002**, *76*, 187–192.
7. Vitamin D Deficiency: The Ultimate Guide. Available online: <https://www.your.md/blog/vitamin-d-deficiency/> (accessed on 21 March 2019).
8. Van Schoor, N.; Lips, P. Global Overview of Vitamin D Status. *Endocrinol. Metab. Clin. N. Am.* **2017**, *46*, 845–870.
9. 90% of UAE Population Vitamin D Deficient, Says DHA Official. Available online: <https://gulfnews.com/going-out/society/90-of-uae-population-vitamin-d-deficient-says-dha-official-1.2113556> (accessed on 22 March 2019).
10. Buyukuslu, N.; Esin, K.; Hizli, H.; Sunal, N.; Yiğit, P.; Garipağaoğlu, M. Clothing preference affects vitamin D status of young women. *Nutr. Res.* **2014**, *34*, 688–693.
11. Vitamin D Deficiency Could Cause Deadly Diseases, Warn UAE Doctors—Khaleej Times. Available online: <https://www.khaleejtimes.com/nation/abu-dhabi/vitamin-d-deficiency-could-cause-deadly-diseases-warn-uae-doctors> (accessed on 22 March 2019).
12. Dunn, P.M. Francis Glisson (1597–1677) and the “Discovery” of Rickets. *Arch. Dis. Child. Fetal Neonatal Ed.* **1998**, *78*, F154–F155.
13. O’Riordan, J.L.H.; Bijvoet, O.L.M. Rickets before the discovery of vitamin D. *BoneKEy Rep.* **2014**, *3*, 3.
14. The History of Vitamin D | Drops® Blog. Available online: <http://blog.vitaminndrops.com/the-history-of-vitamin-d/> (accessed on 22 March 2019).
15. McCollum, E.V.; Simmonds, N.; Becker, J.E.; Shipley, P.G. Studies on experimental rickets. *Nutr. Rev.* **1975**, *33*, 48–50.
16. Cantorna, M.T.; Hayes, C.E.; DeLuca, H.F. 1,25-Dihydroxycholecalciferol Inhibits the Progression of Arthritis in Murine Models of Human Arthritis. *J. Nutr.* **1998**, *128*, 68–72.
17. Chun, R.F.; Peercy, B.E.; Orwoll, E.S.; Nielson, C.M.; Adams, J.S.; Hewison, M. Vitamin D and DBP: The free hormone hypothesis revisited. *J. Steroid Biochem. Mol. Biol.* **2014**, *144*, 132–137.
18. Christakos, S.; Hewison, M.; Gardner, D.G.; Wagner, C.L.; Sergeev, I.N.; Rutten, E.; Pittas, A.G.; Boland, R.; Ferrucci, L.; Bikle, D.D. Vitamin D: Beyond bone: Vitamin D: Beyond bone. *Ann. N. Y. Acad. Sci.* **2013**, *1287*, 45–58.
19. Hewison, M. Vitamin D and the immune system: New perspectives on an old theme. *Endocrinol. Metab. Clin. N. Am.* **2010**, *39*, 365–379.
20. Welsh, J. Cellular and molecular effects of vitamin D on carcinogenesis. *Arch. Biochem. Biophys.* **2012**, *523*, 107–114.
21. Mondul, A.M.; Weinstein, S.J.; Layne, T.M.; Albanes, D. Vitamin D and Cancer Risk and Mortality: State of the Science, Gaps, and Challenges. *Epidemiol. Rev.* **2017**, *39*, 28–48.
22. Guillot, X.; Semerano, L.; Saidenberg-Kermanac’h, N.; Falgarone, G.; Boissier, M.-C. Vitamin D and inflammation. *Joint Bone Spine* **2010**, *77*, 552–557.

23. Wolf, M.; Shah, A.; Gutierrez, O.; Ankers, E.; Monroy, M.; Tamez, H.; Steele, D.; Chang, Y.; Camargo, C.; Tonelli, M.; et al. Vitamin D levels and early mortality among incident hemodialysis patients. *Kidney Int.* **2007**, *72*, 1004–1013.
24. Vimalaswaran, K.S.; Cavadino, A.; Berry, D.J.; Jorde, R.; Dieffenbach, A.K.; Lu, C.; Alves, A.C.; Heerspink, H.J.L.; Tikkanen, E.; Eriksson, J.; et al. Association of vitamin D status with arterial blood pressure and hypertension risk: A mendelian randomisation study. *Lancet Diabetes Endocrinol.* **2014**, *2*, 719–729.
25. Vanlint, S. Vitamin D and Obesity. *Nutrients* **2013**, *5*, 949–956.
26. Pilz, S.; Grübler, M.; Gaksch, M.; Schwetz, V.; Trummer, C.; Hartaigh, B.Ó.; Verheyen, N.; Tomaschitz, A.; März, W. Vitamin D and Mortality. *Anticancer Res.* **2016**, *36*, 1379–1387.
27. Penckofer, S.; Kouba, J.; Wallis, D.E.; Emanuele, M.A. Vitamin D and Diabetes Let the Sunshine in. *Diabetes Educ.* **2008**, *34*, 939–954.
28. Jeong, H.Y.; Park, K.M.; Lee, M.J.; Yang, D.H.; Kim, S.H.; Lee, S.-Y. Vitamin D and Hypertension. *Electrolyte Blood Press* **2017**, *15*, 1–11.
29. Low Vitamin D Levels in Adults | Empoweryourhealth.org. Available online: <https://www.empoweryourhealth.org/issue-1/Low-Vitamin-D-LEVELS-IN-ADULTS> (accessed on 31 March 2019).
30. Schmidt, J.A. Measurement of 25-Hydroxyvitamin D Revisited. *Clin. Chem.* **2006**, *52*, 2304–2305.
31. Terry, A.H.; Sandrock, T.; Meikle, A.W. Measurement of 25-Hydroxyvitamin D by the Nichols ADVANTAGE, DiaSorin LIAISON, DiaSorin RIA, and Liquid Chromatography-Tandem Mass Spectrometry. *Clin. Chem.* **2005**, *51*, 1565–1566.
32. Shah, I.; James, R.; Barker, J.; Petroczi, A.; Naughton, D.P. Misleading measures in Vitamin D analysis: A novel LC-MS/MS assay to account for epimers and isobars. *Nutr. J.* **2011**, *10*, 46.
33. De Koning, L.; Al-Turkmani, M.R.; Berg, A.H.; Shkreta, A.; Law, T.; Kellogg, M.D. Variation in clinical vitamin D status by DiaSorin Liaison and LC-MS/MS in the presence of elevated 25-OH vitamin D₂. *Clin. Chim. Acta* **2013**, *415*, 54–58.
34. Vitamin D Deficiency: How Common Is It and What Can I Do about It? | Ddrops® Blog. Available online: <http://blog.vitaminddrops.com/vitamin-d-deficiency-common-can/> (accessed on 22 March 2019).
35. Nair, R.; Maseeh, A. Vitamin D: The “sunshine” vitamin. *J. Pharmacol. Pharmacother.* **2012**, *3*, 118.
36. *Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium Dietary Reference Intakes for Calcium and Vitamin D*; Ross, A.C., Taylor, C.L., Yaktine, A.L., Del Valle, H.B., Eds.; The National Academies Collection: Reports Funded by National Institutes of Health; National Academies Press (US): Washington, DC, USA, 2011.
37. Jäpelt, R.B.; Jakobsen, J. Vitamin D in plants: A review of occurrence, analysis, and biosynthesis. *Front. Plant Sci.* **2013**, *4*, 136.
38. Wierzbicka, J.; Piotrowska, A.; Żmijewski, M.A. The renaissance of vitamin D. *Acta Biochim. Pol.* **2014**, *61*, 679–686.
39. Prohormones—A Comprehensive Guide to Prohormones—Updated | Predator Nutrition. Available online: <https://www.predatornutrition.com/articlesdetail?cid=prohormones-a-comprehensive-guide-to-prohormones-updated> (accessed on 26 March 2019).
40. Vitamin D | Hormone Health Network. Available online: <https://www.hormone.org/hormones-and-health/hormones/vitamin-d> (accessed on 26 March 2019).
41. Maestro, M.A.; Molnár, F.; Mouriño, A.; Carlberg, C. Vitamin D receptor 2016: Novel ligands and structural insights. *Expert Opin. Ther. Patents* **2016**, *26*, 1291–1306.
42. Seuter, S.; Ryyänen, J.; Carlberg, C. The ASAP2 gene is a primary target of 1,25-dihydroxyvitamin D₃ in human monocytes and macrophages. *J. Steroid Biochem. Mol. Biol.* **2014**, *144*, 12–18.
43. Norman, A.W.; Röth, J.; Orci, L. The Vitamin D Endocrine System: Steroid Metabolism, Hormone Receptors, and Biological Response (Calcium Binding Proteins). *Endocr. Rev.* **1982**, *3*, 331–366.
44. Menegaz, D.; Mizwicki, M.T.; Barrientos-Duran, A.; Chen, N.; Henry, H.L.; Norman, A.W. Vitamin D Receptor (VDR) Regulation of Voltage-Gated Chloride Channels by Ligands Preferring a VDR-Alternative Pocket (VDR-AP). *Mol. Endocrinol.* **2011**, *25*, 1289–1300.
45. Jovičić, S.; Ignjatovic, S.; Majkić-Singh, N. Biochemistry and metabolism of vitamin D/Biohemija i metabolizam vitamina D. *J. Med. Biochem.* **2012**, *31*, 309–315.
46. Wikvall, K. Cytochrome P450 enzymes in the bioactivation of vitamin D to its hormonal form (Review). *Int. J. Mol. Med.* **2001**, *7*, 201–209.

47. Jones, G.; Prosser, D.E.; Kaufmann, M. Cytochrome P450-mediated metabolism of vitamin D. *J. Lipid Res.* **2014**, *55*, 13–31.
48. Hollis, B.W. Comparison of equilibrium and disequilibrium assay conditions for ergocalciferol, cholecalciferol and their major metabolites. *J. Steroid Biochem.* **1984**, *21*, 81–86.
49. Hewison, M. Antibacterial effects of vitamin D. *Nat. Rev. Endocrinol.* **2011**, *7*, 337–345.
50. Bikle, D. Vitamin D: Production, Metabolism, and Mechanisms of Action. In *Endotext*; Feingold, K.R., Anawalt, B., Boyce, A., Chrousos, G., Dungan, K., Grossman, A., Hershman, J.M., Kaltsas, G., Koch, C., Kopp, P., et al., Eds.; MDText.com, Inc.: South Dartmouth, MA, USA, 2000.
51. Kamao, M.; Tatematsu, S.; Hatakeyama, S.; Sakaki, T.; Sawada, N.; Inouye, K.; Ozono, K.; Kubodera, N.; Reddy, G.S.; Okano, T. C-3 Epimerization of Vitamin D₃ Metabolites and Further Metabolism of C-3 Epimers 25-hydroxyvitamin D₃ is metabolized to 3-epi-25-hydroxyvitamin D₃ and subsequently metabolized through C-1 α or C-24 hydroxylation. *J. Biol. Chem.* **2004**, *279*, 15897–15907.
52. Bailey, D.; Veljkovic, K.; Yazdanpanah, M.; Adeli, K. Analytical measurement and clinical relevance of vitamin D₃ C₃-epimer. *Clin. Biochem.* **2013**, *46*, 190–196.
53. Cashman, K.D.; Hayes, A.; Galvin, K.; Merkel, J.; Jones, G.; Kaufmann, M.; Hoofnagle, A.N.; Carter, G.D.; Durazo-Arvizu, R.A.; Sempos, C.T. Significance of Serum 24,25-Dihydroxyvitamin D in the Assessment of Vitamin D Status: A Double-edged Sword? *Clin. Chem.* **2015**, *61*, 636–645.
54. Michio, I.; Kuniyama, K.; Toru, Y.; Toshiaki, O. A simple and sensitive assay for 25-hydroxyvitamin D, 24,25-dihydroxyvitamin D and 1,25-dihydroxyvitamin D in human serum. *Clin. Chim. Acta* **1982**, *124*, 63–73.
55. Somjen, D.; Somjen, G.J.; Weisman, Y.; Binderman, I.; Dokoh, S.; Haussler, M.R.; Pike, J.W.; Danan, J.L.; Mathieu, H.; Lawson, D.E.M.; et al. Evidence for 24,25-dihydroxycholecalciferol receptors in long bones of newborn rats. *Biochem. J.* **1982**, *204*, 31–36.
56. Knutson, J.C.; Hollis, B.W.; LeVan, L.W.; Valliere, C.; Gould, K.G.; Bishop, C.W. Metabolism of 1 alpha-hydroxyvitamin D₂ to activated dihydroxyvitamin D₂ metabolites decreases endogenous 1 alpha, 25-dihydroxyvitamin D₃ in rats and monkeys. *Endocrinology* **1995**, *136*, 4749–4753.
57. Reddy, G.S.; Muralidharan, K.R.; Okamura, W.H.; Tserng, K.Y.; McLane, J.A. Metabolism of 1alpha,25-dihydroxyvitamin D(3) and its C-3 epimer 1alpha,25-dihydroxy-3-epi-vitamin D(3) in neonatal human keratinocytes. *Steroids* **2001**, *66*, 441–450.
58. Brown, A.J.; Ritter, C.; Slatopolsky, E.; Muralidharan, K.R.; Okamura, W.H.; Reddy, G.S. 1Alpha,25-dihydroxy-3-epi-vitamin D₃, a natural metabolite of 1alpha,25-dihydroxyvitamin D₃, is a potent suppressor of parathyroid hormone secretion. *J. Cell. Biochem.* **1999**, *73*, 106–113.
59. Kaseda, R.; Hosojima, M.; Sato, H.; Saito, A. Role of Megalin and Cubilin in the Metabolism of Vitamin D₃. *Ther. Apher. Dial.* **2011**, *15*, 14–17.
60. Romagnoli, E.; Pepe, J.; Piemonte, S.; Cipriani, C.; Minisola, S. Value and limitations of assessing vitamin D nutritional status and advised levels of vitamin D supplementation. *Eur. J. Endocrinol.* **2013**, *169*, R59–R69.
61. Chun, R.F.; Percy, B.E.; Adams, J.S.; Hewison, M. Vitamin D Binding Protein and Monocyte Response to 25-Hydroxy Vitamin D and 1,25-Dihydroxy Vitamin D: Analysis by Mathematical Modeling. *PlosOne.* **2012**, *7*, E30773, doi:10.1371/journal.pone.0030773.
62. Heaney, R.P.; Armas, L.A.G.; Shary, J.R.; Bell, N.H.; Binkley, N.; Hollis, B.W. 25-Hydroxylation of vitamin D₃: Relation to circulating vitamin D₃ under various input conditions. *Am. J. Clin. Nutr.* **2008**, *87*, 1738–1742.
63. Nussey, S.; Whitehead, S. *Endocrinology: An Integrated Approach*; BIOS Scientific Publishers: Oxford, UK, 2001; ISBN 978-1-85996-252-7.
64. Rowling, M.J.; Taffany, D.A.; Welsh, J.; Kemmis, C.M. Megalin-Mediated Endocytosis of Vitamin D Binding Protein Correlates with 25-Hydroxycholecalciferol Actions in Human Mammary Cells. *J. Nutr.* **2006**, *136*, 2754–2759.
65. Luo, W.; Karpf, A.R.; Deeb, K.K.; Muindi, J.R.; Morrison, C.D.; Johnson, C.S.; Trump, D.L. Epigenetic Regulation of Vitamin D 24-Hydroxylase/CYP24A1 in Human Prostate Cancer. *Cancer Res.* **2010**, *70*, 5953–5962.
66. Mangelsdorf, D.J.; Thummel, C.; Beato, M.; Herrlich, P.; Schütz, G.; Umesono, K.; Blumberg, B.; Kastner, P.; Mark, M.; Chambon, P.; et al. The nuclear receptor superfamily: The second decade. *Cell* **1995**, *83*, 835–839.

67. Rastinejad, F.; Huang, P.; Chandra, V.; Khorasanizadeh, S. Understanding nuclear receptor form and function using structural biology. *J. Mol. Endocrinol.* **2013**, *51*, T1–T21.
68. Luong, K.V.Q.; Nguyen, L.T.H.; Luong, K.V.Q.; Nguyễn, L.T.H. The beneficial role of vitamin D and its analogs in cancer treatment and prevention. *Crit. Rev. Oncol.* **2010**, *73*, 192–201.
69. Schwartz, Z.; Sylvia, V.L.; Larsson, D.; Nemere, I.; Casasola, D.; Dean, D.D.; Boyan, B.D. $1\alpha,25(\text{OH})_2\text{D}_3$ Regulates Chondrocyte Matrix Vesicle Protein Kinase C (PKC) Directly via G-protein-dependent Mechanisms and Indirectly via Incorporation of PKC during Matrix Vesicle Biogenesis. *J. Biol. Chem.* **2002**, *277*, 11828–11837.
70. Zhang, C.; Tang, W.; Li, Y.; Yang, F.; Dowd, D.R.; Macdonald, P.N. Osteoblast-Specific Transcription Factor Osterix Increases Vitamin D Receptor Gene Expression in Osteoblasts. *PLoS ONE* **2011**, *6*, e26504.
71. Valdivielso, J.M.; Fernández, E. Vitamin D receptor polymorphisms and diseases. *Clin. Chim. Acta* **2006**, *371*, 1–12.
72. Vdr MGI Mouse Gene Detail—MGI:103076—Vitamin D (1,25-Dihydroxyvitamin D₃) Receptor. Available online: <http://www.informatics.jax.org/marker/MGI:103076> (accessed on 16 April 2019).
73. Garcia-Vallvé, S.; Palau, J. Nuclear receptors, nuclear-receptor factors, and nuclear-receptor-like orphans form a large paralog cluster in Homo sapiens. *Mol. Biol. Evol.* **1998**, *15*, 665–682.
74. Kouzmenko, A.; Ohtake, F.; Fujiki, R.; Kato, S. Chapter 12—Epigenetic Modifications in Vitamin D Receptor-mediated Transrepression. In *Vitamin D*, 3rd ed.; Feldman, D., Pike, J.W., Adams, J.S., Eds.; Academic Press: San Diego, CA, USA, 2011; pp. 227–234, ISBN 978-0-12-381978-9.
75. Wei, P.; Inamdar, N.; Vedeckis, W.V. Transrepression of c-jun Gene Expression by the Glucocorticoid Receptor Requires Both AP-1 Sites in the c-jun Promoter. *Mol. Endocrinol.* **1998**, *12*, 1322–1333.
76. Chen, S.; Wu, J.; Hsieh, J.C.; Whitfield, G.K.; Jurutka, P.W.; Haussler, M.R.; Gardner, D.G. Suppression of ANP Gene Transcription by Liganded Vitamin D Receptor. *Hypertension* **1998**, *31*, 1338–1342.
77. Heikkinen, S.; Väisänen, S.; Pehkonen, P.; Seuter, S.; Benes, V.; Carlberg, C. Nuclear hormone $1\alpha,25$ -dihydroxyvitamin D₃ elicits a genome-wide shift in the locations of VDR chromatin occupancy. *Nucleic Acids Res.* **2011**, *39*, 9181–9193.
78. Meyer, M.B.; Goetsch, P.D.; Pike, J.W. VDR/RXR and TCF4/ β -catenin cistromes in colonic cells of colorectal tumor origin: Impact on c-FOS and c-MYC gene expression. *Mol. Endocrinol.* **2012**, *26*, 37–51.
79. Pike, J.W.; Meyer, M.B. Regulation of Mouse Cyp24a1 Expression via Promoter-Proximal and Downstream-Distal Enhancers Highlights New Concepts of 1,25-Dihydroxyvitamin D₃ Action. *Arch. Biochem. Biophys.* **2012**, *523*, 2–8.
80. Carlberg, C.; Seuter, S.; De Mello, V.D.F.; Schwab, U.; Voutilainen, S.; Pulkki, K.; Nurmi, T.; Virtanen, J.; Tuomainen, T.-P.; Uusitupa, M. Primary Vitamin D Target Genes Allow a Categorization of Possible Benefits of Vitamin D₃ Supplementation. *PLoS ONE* **2013**, *8*, e71042.
81. Norman, A.W.; Mizwicki, M.T.; Norman, D.P.G. Steroid-hormone rapid actions, membrane receptors and a conformational ensemble model. *Nat. Rev. Drug Discov.* **2004**, *3*, 27–41.
82. Rochel, N.; Wurtz, J.-M.; Mitschler, A.; Klaholz, B.; Moras, D. The Crystal Structure of the Nuclear Receptor for Vitamin D Bound to Its Natural Ligand. *Mol. Cell* **2000**, *5*, 173–179.
83. Bhattacharjee, S.; Renganaath, K.; Mehrotra, R.; Mehrotra, S. Combinatorial Control of Gene Expression. *BioMed Res. Int.* **2013**, *2013*, 1–11.
84. Uitterlinden, A.G.; Fang, Y.; Van Meurs, J.B.; Pols, H.A.; Van Leeuwen, J.P. Genetics and biology of vitamin D receptor polymorphisms. *Gene* **2004**, *338*, 143–156.
85. Huang, J.; Xie, Z.-F. Polymorphisms in the vitamin D receptor gene and multiple sclerosis risk: A meta-analysis of case-control studies. *J. Neurol. Sci.* **2012**, *313*, 79–85.
86. Hii, C.S.; Ferrante, A. The Non-Genomic Actions of Vitamin D. *Nutrients* **2016**, *8*, 135.
87. Chen, J.; Doroudi, M.; Cheung, J.; Grozier, A.L.; Schwartz, Z.; Boyan, B.D. Plasma membrane Pdia3 and VDR interact to elicit rapid responses to $1\alpha,25(\text{OH})_2\text{D}_3$. *Cell Signal.* **2013**, *25*, 2362–2373.
88. Nemere, I.; Farach-Carson, M.C.; Rohe, B.; Sterling, T.M.; Norman, A.W.; Boyan, B.D.; Safford, S.E. Ribozyme knockdown functionally links a 1,25(OH)₂D₃ membrane binding protein (1,25D₃-MARRS) and phosphate uptake in intestinal cells. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 7392–7397.
89. Nemere, I. Immunochemical studies on the putative plasmalemmal receptor for 1,25-dihydroxyvitamin D₃. III. Vitamin D status. *Steroids* **2000**, *65*, 451–457.
90. Pike, J.W.; Meyer, M.B. The Vitamin D Receptor: New Paradigms for the Regulation of Gene Expression by 1,25-Dihydroxyvitamin D₃. *Endocrinol. Metab. Clin. N. Am.* **2010**, *39*, 255–269.

91. Larriba, M.J.; González-Sancho, J.M.; Bonilla, F.; Muñoz, A. Interaction of vitamin D with membrane-based signaling pathways. *Front. Physiol.* **2014**, *5*, 60.
92. Zhao, G.; Simpson, R.U. Interaction between vitamin D receptor with caveolin-3 and regulation by 1,25-dihydroxyvitamin D₃ in adult rat cardiomyocytes. *J. Steroid Biochem. Mol. Biol.* **2010**, *121*, 159–163.
93. Garbi, N.; Tanaka, S.; Momburg, F.; Hämmerling, G.J. Impaired assembly of the major histocompatibility complex class I peptide-loading complex in mice deficient in the oxidoreductase ERp57. *Nat. Immunol.* **2006**, *7*, 93–102.
94. Wang, Y.; Chen, J.; Lee, C.S.; Nizkorodov, A.; Riemenschneider, K.; Martin, D.; Hyzy, S.; Schwartz, Z.; Boyan, B.D. Disruption of Pdia3 gene results in bone abnormality and affects 1 α ,25-dihydroxy-vitamin D₃-induced rapid activation of PKC. *J. Steroid Biochem. Mol. Biol.* **2010**, *121*, 257–260.
95. Nemere, I.; Garbi, N.; Hämmerling, G.J.; Khanal, R.C. Intestinal Cell Calcium Uptake and the Targeted Knockout of the 1,25D₃-MARRS (Membrane-associated, Rapid Response Steroid-binding) Receptor/PDIA3/Erp57. *J. Boil. Chem.* **2010**, *285*, 31859–31866.
96. Boyan, B.D.; Sylvania, V.L.; McKinney, N.; Schwartz, Z. Membrane actions of vitamin D metabolites 1 α ,25(OH)₂D₃ and 24R,25(OH)₂D₃ are retained in growth plate cartilage cells from vitamin D receptor knockout mice. *J. Cell. Biochem.* **2003**, *90*, 1207–1223.
97. Patel, H.H.; Murray, F.; Insel, P.A. Caveolae as Organizers of Pharmacologically Relevant Signal Transduction Molecules. *Annu. Rev. Pharmacol. Toxicol.* **2008**, *48*, 359–391.
98. Bosch, M.; Marí, M.; Herms, A.; Fernández, A.; Fajardo, A.; Kassan, A.; Giralt, A.; Colell, A.; Balgoma, D.; Barbero, E.; et al. Caveolin-1 deficiency causes cholesterol dependent mitochondrial dysfunction and apoptotic susceptibility. *Curr. Boil.* **2011**, *21*, 681–686.
99. Mesbah, M.; Nemere, I.; Papagerakis, P.; Nefussi, J.-R.; Nessmann, C.; Berdal, A.; Orestes-Cardoso, S.; Orestes-Cardoso, S. Expression of a 1,25-Dihydroxyvitamin D₃ Membrane-Associated Rapid-Response Steroid Binding Protein During Human Tooth and Bone Development and Biomineralization. *J. Bone Miner. Res.* **2002**, *17*, 1588–1596.
100. Norman, A.W. Receptors for 1 α ,25(OH)₂D₃: Past, Present, and Future. *J. Bone Miner. Res.* **1998**, *13*, 1360–1369.
101. Richard, C.L.; Farach-Carson, M.C.; Rohe, B.; Nemere, I.; Meckling, K.A. Involvement of 1,25D₃-MARRS (membrane associated, rapid response steroid-binding), a novel vitamin D receptor, in growth inhibition of breast cancer cells. *Exp. Cell Res.* **2010**, *316*, 695–703.
102. Farach-Carson, M.; Nemere, I. Membrane Receptors for Vitamin D Steroid Hormones: Potential New Drug Targets. *Curr. Drug Targets* **2003**, *4*, 67–76.
103. Sitrin, M. Rapid effects of 1,25(OH)₂ vitamin D₃ on signal transduction systems in colonic cells. *Steroids* **1999**, *64*, 137–142.
104. Chen, A.; Davis, B.H.; Bissonnette, M.; Scaglione-Sewell, B.; Brasitus, T.A. 1,25-Dihydroxyvitamin D₃ Stimulates Activator Protein-1-dependent Caco-2 Cell Differentiation. *J. Boil. Chem.* **1999**, *274*, 35505–35513.
105. Doroudi, M.; Boyan, B.D.; Schwartz, Z. Rapid 1 α ,25(OH)₂D₃ membrane-mediated activation of Ca²⁺/calmodulin-dependent protein kinase II in growth plate chondrocytes requires Pdia3, PLAA and caveolae. *Connect. Tissue Res.* **2014**, *55*, 125–128.
106. Rohe, B.; Safford, S.E.; Nemere, I.; Farach-Carson, M.C. Identification and characterization of 1,25D₃-membrane-associated rapid response, steroid (1,25D₃-MARRS)-binding protein in rat IEC-6 cells. *Steroids* **2005**, *70*, 458–463.
107. Nemere, I.; Safford, S.E.; Rohe, B.; DeSouza, M.M.; Farach-Carson, M.C. Identification and characterization of 1,25D₃-membrane-associated rapid response, steroid (1,25D₃-MARRS) binding protein. *J. Steroid Biochem. Mol. Biol.* **2004**, *89*, 281–285.
108. Pillai, S.; Bikle, D.D.; Su, M.J.; Ratnam, A.; Abe, J. 1,25-Dihydroxyvitamin D₃ upregulates the phosphatidylinositol signaling pathway in human keratinocytes by increasing phospholipase C levels. *J. Clin. Investig.* **1995**, *96*, 602–609.
109. Alkon, D.; Rasmussen, H. A spatial-temporal model of cell activation. *Science* **1988**, *239*, 998–1005.
110. Sato, K.; Imaki, T.; Toraya, S.; Demura, H.; Tanaka, M.; Kasajima, T.; Takeuchi, A.; Kobayashi, T. Increased 1,25-(OH)₂D₂ Concentration in a Patient with Malignancy-Associated Hypercalcemia Receiving Intravenous Hydration Inadvertently Supplemented with Vitamin D₂. *Intern. Med.* **1993**, *32*, 886–890.

111. Boyan, B.D.; Wang, L.; Wong, K.L.; Jo, H.; Schwartz, Z. Plasma membrane requirements for $1\alpha,25(\text{OH})_2\text{D}_3$ dependent PKC signaling in chondrocytes and osteoblasts. *Steroids* **2006**, *71*, 286–290.
112. Schwartz, Z.; Shaked, D.; Hardin, R.; Gruwell, S.; Dean, D.; Sylvia, V.; Boyan, B.; Dean, D. $1\alpha,25(\text{OH})_2\text{D}_3$ causes a rapid increase in phosphatidylinositol-specific PLC- β activity via phospholipase A2-dependent production of lysophospholipid. *Steroids* **2003**, *68*, 423–437.
113. Peehl, D.M.; Skowronski, R.J.; Leung, G.K.; Wong, S.T.; Stamey, T.A.; Feldman, D. Antiproliferative effects of 1,25-dihydroxyvitamin D3 on primary cultures of human prostatic cells. *Cancer Res.* **1994**, *54*, 805–810.
114. Lin, R.; Nagai, Y.; Sladek, R.; Bastien, Y.; Ho, J.; Petrecca, K.; Sotiropoulou, G.; Diamandis, E.P.; Hudson, T.J.; White, J.H. Expression Profiling in Squamous Carcinoma Cells Reveals Pleiotropic Effects of Vitamin D3 Analog EB1089 Signaling on Cell Proliferation, Differentiation, and Immune System Regulation. *Mol. Endocrinol.* **2002**, *16*, 1243–1256.
115. Bao, B.-Y.; Yao, J.; Lee, Y.-F. $1\alpha, 25$ -dihydroxyvitamin D3 suppresses interleukin-8-mediated prostate cancer cell angiogenesis. *Carcinogenesis* **2006**, *27*, 1883–1893.
116. Hilliard, G.M.; Cook, R.G.; Weigel, N.L.; Pike, J.W. $1,25$ -Dihydroxyvitamin D3 modulates phosphorylation of serine 205 in the human vitamin D receptor: Site-directed mutagenesis of this residue promotes alternative phosphorylation. *Biochemistry* **1994**, *33*, 4300–4311.
117. Wu, W.; Beilhartz, G.; Roy, Y.; Richard, C.L.; Curtin, M.; Brown, L.; Cadieux, D.; Coppolino, M.; Farach-Carson, M.C.; Nemere, I.; et al. Nuclear translocation of the $1,25\text{D}_3$ -MARRS (membrane associated rapid response to steroids) receptor protein and NFkappaB in differentiating NB4 leukemia cells. *Exp. Cell Res.* **2010**, *316*, 1101–1108.
118. Barletta, F.; Freedman, L.P.; Christakos, S. Enhancement of VDR-Mediated Transcription by Phosphorylation: Correlation with Increased Interaction between the VDR and DRIP205, a Subunit of the VDR-Interacting Protein Coactivator Complex. *Mol. Endocrinol.* **2002**, *16*, 301–314.
119. Norman, A.W. Vitamin D Receptor: New Assignments for an Already Busy Receptor. *Endocrinology* **2006**, *147*, 5542–5548.
120. Deeb, K.K.; Trump, D.L.; Johnson, C.S. Vitamin D signalling pathways in cancer: Potential for anticancer therapeutics. *Nat. Rev. Cancer* **2007**, *7*, 684–700.
121. Nussey, S.; Whitehead, S. *The Parathyroid Glands and Vitamin D*; BIOS Scientific Publishers: Didcott, UK, 2001.
122. Tanaka, Y.; DeLuca, H. Bone mineral mobilization activity of $1,25$ -dihydroxycholecalciferol, a metabolite of vitamin D. *Arch. Biochem. Biophys.* **1971**, *146*, 574–578.
123. Parathyroid Hormone | You and Your Hormones from the Society for Endocrinology. Available online: <http://www.yourhormones.info/hormones/parathyroid-hormone/> (accessed on 8 April 2019).
124. Garabedian, M.; Holick, M.; DeLuca, H.F.; Boyle, I.T. Control of 25 -Hydroxycholecalciferol Metabolism by Parathyroid Glands. *Proc. Natl. Acad. Sci. USA* **1972**, *69*, 1673–1676.
125. Brenza, H.L.; DeLuca, H.F. Regulation of 25 -Hydroxyvitamin D3 1α -Hydroxylase Gene Expression by Parathyroid Hormone and $1,25$ -Dihydroxyvitamin D3. *Arch. Biochem. Biophys.* **2000**, *381*, 143–152.
126. Chambers, T.J.; Magnus, C.J. Calcitonin alters behaviour of isolated osteoclasts. *J. Pathol.* **1982**, *136*, 27–39.
127. Shinki, T.; Ueno, Y.; DeLuca, H.F.; Suda, T. Calcitonin is a major regulator for the expression of renal 25 -hydroxyvitamin D3- 1α -hydroxylase gene in normocalcemic rats. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 8253–8258.
128. Genetics Home Reference. CYP24A1 Gene. Available online: <https://ghr.nlm.nih.gov/gene/CYP24A1> (accessed on 8 April 2019).
129. Vuolo, L.; Di Somma, C.; Faggiano, A.; Colao, A. Vitamin D and cancer. *Front. Endocrinol.* **2012**, *3*, 58.
130. Gross, M.; Kost, S.B.; Ennis, B.; Stumpf, W.; Kumar, R. Effect of $1,25$ -dihydroxyvitamin D3 on mouse mammary tumor (GR) cells: Evidence for receptors, cellular uptake, inhibition of growth and alteration in morphology at physiologic concentrations of hormone. *J. Bone Miner. Res.* **1986**, *1*, 457–467.
131. Narvaez, C.J.; Matthews, D.; LaPorta, E.; Simmons, K.M.; Beaudin, S.; Welsh, J. The impact of vitamin D in breast cancer: Genomics, pathways, metabolism. *Front. Physiol.* **2014**, *5*, 5.
132. Fleet, J.C. Molecular Actions of Vitamin D Contributing to Cancer Prevention. *Mol. Asp. Med.* **2008**, *29*, 388–396.
133. Skowronski, R.J.; Peehl, D.M.; Feldman, D. Vitamin D and prostate cancer: $1,25$ dihydroxyvitamin D3 receptors and actions in human prostate cancer cell lines. *Endocrinology* **1993**, *132*, 1952–1960.
134. Basit, S. Vitamin D in health and disease: A literature review. *Br. J. Biomed. Sci.* **2013**, *70*, 161–72.

135. Mahendra, A.; Karishma; Choudhury, B.K.; Sharma, T.; Bansal, N.; Bansal, R.; Gupta, S. Vitamin D and gastrointestinal cancer. *J. Lab. Physicians* **2018**, *10*, 1–5.
136. Leigh-Clare, J.L. A note on the vitamin D content of the stomach oil of the australasian petrel (*aestralata lessoni*). *Biochem. J.* **1927**, *21*, 725–727.
137. Selye, H.; Bois, P. On the role of corticoids in conditioning the gastric mucosa to certain toxic actions of ergocalciferol. *Br. J. Nutr.* **1957**, *11*, 18–22.
138. Stumpf, W.; Sar, M.; Reid, F.; Tanaka, Y.; DeLuca, H. Target cells for 1,25-dihydroxyvitamin D₃ in intestinal tract, stomach, kidney, skin, pituitary, and parathyroid. *Science* **1979**, *206*, 1188–1190.
139. Kirui, N.A.; Weisbrode, S.E.; Kindig, O.R. The role of dietary calcium on the development of soft tissue mineralization due to 1,25-dihydroxyvitamin D₃ intoxication in rats. *Virchows Arch. B Cell Pathol. Incl. Mol. Pathol.* **1981**, *37*, 251–263.
140. Selking, Ö.; Borch, K.; Johansson, H.; Ljunghall, S.; Wide, L. Evaluation of Parathyroid Function in Patients with Hypergastrinaemia and Pernicious Anaemia. *Uppsala J. Med. Sci.* **1982**, *87*, 215–222.
141. Kurose, T.; Seino, Y.; Ishida, H.; Tsuji, K.; Fukumoto, H.; Koh, G.; Takeda, J.; Kitano, N.; Inagaki, N.; Tsuda, K.; et al. Effect of vitamin D on gastrin and gastric somatostatin secretion from the isolated perfused rat stomach. *Life Sci.* **1988**, *42*, 1995–2001.
142. Axelson, J.; Persson, P.; Gagnemo-Persson, R.; Hakanson, R. *Gut* **1991**, *32*, 1298–1302.
143. Holick, M.; Holick, M. Noncalcemic actions of 1,25-dihydroxyvitamin D₃ and clinical applications. *Bone* **1995**, *17*, S107–S111.
144. Stumpf, W.E. Vitamin D sites and mechanisms of action: A histochemical perspective. Reflections on the utility of autoradiography and cytopharmacology for drug targeting. *Histochem. Cell Boil.* **1995**, *104*, 417–427.
145. Ikezaki, S.; Nishikawa, A.; Furukawa, F.; Tanakamura, Z.; Kim, H.C.; Mori, H.; Takahashi, M. Chemopreventive effects of 24R,25-dihydroxyvitamin D₃, a vitamin D₃ derivative, on glandular stomach carcinogenesis induced in rats by N-methyl-N'-nitro-N-nitrosoguanidine and sodium chloride. *Cancer Res.* **1996**, *56*, 2767–2770.
146. Gagnemo-Persson, R.; Persson, P.; Bryngelsson, T.; Green, B.; Håkanson, R. Rat stomach ECL-cell histidine decarboxylase activity is suppressed by ergocalciferol but unaffected by parathyroid hormone and calcitonin. *Regul. Pept.* **1999**, *79*, 131–139.
147. Stumpf, W.E. Vitamin D and the digestive system. *Eur. J. Drug Metab. Pharmacokinet.* **2008**, *33*, 85–100.
148. Häkkinen, I.; Lindgren, I. The localization of tetracycline in the metastatic calcifications in the stomach of rat induced by overdose of dihydrotachysterol and vitamin D₃. *Acta Pathol. Microbiol. Scand.* **2009**, *59*, 428–434.
149. Sahin, H.H.; Cumbul, A.; Uslu, Ünal; Yilmaz, Z.; Ercan, F.; Alican, I. The effect of 1,25 dihydroxyvitamin D₃ on HCl/Ethanol-induced gastric injury in rats. *Tissue Cell* **2018**, *51*, 68–76.
150. Paterson, C.R.; Woods, C.G. Search for osteomalacia in 1228 patients after gastrectomy and other operations on the stomach. *Lancet* **1965**, *2*, 1085–1088.
151. Kimura, K.; Nozawa, Y.; Kitamura, S.; Takahashi, H.; Ota, M.; Norimatsu, H.; Wada, H. An autopsy case of hypervitaminosis D. *Pathol. Int.* **1967**, *17*, 377–386.
152. Lawrence, W. Nutritional consequences of surgical resection of the gastrointestinal tract for cancer. *Cancer Res.* **1977**, *37*, 2379–2386.
153. Giovannucci, E.; Liu, Y.; Willett, W.C. Cancer Incidence and Mortality and Vitamin D in Black and White Male Health Professionals. *Cancer Epidemiol. Biomark. Prev.* **2006**, *15*, 2467–2472.
154. Chen, W.; Dawsey, S.M.; Qiao, Y.-L.; Mark, S.D.; Dong, Z.-W.; Taylor, P.R.; Zhao, P.; Abnet, C.C. Prospective study of serum 25(OH)-vitamin D concentration and risk of oesophageal and gastric cancers. *Br. J. Cancer* **2007**, *97*, 123–128.
155. Pan, L.; Matloob, A.F.; Du, J.; Pan, H.; Dong, Z.; Zhao, J.; Feng, Y.; Zhong, Y.; Huang, B.; Lu, J. Vitamin D stimulates apoptosis in gastric cancer cells in synergy with trichostatin A /sodium butyrate-induced and 5-aza-2'-deoxycytidine-induced PTEN upregulation. *FEBS J.* **2010**, *277*, 989–999.
156. Antico, A.; Tozzoli, R.; Giavarina, D.; Tonutti, E.; Bizzaro, N. Hypovitaminosis D as predisposing factor for atrophic type A gastritis: A case-control study and review of the literature on the interaction of Vitamin D with the immune system. *Clin. Rev. Allergy Immunol.* **2012**, *42*, 355–364.

157. Park, M.R.; Lee, J.H.; Park, M.S.; Hwang, J.E.; Shim, H.J.; Cho, S.H.; Chung, I.-J.; Bae, W.K. Suppressive Effect of 19-nor-1 α -25-Dihydroxyvitamin D₂ on Gastric Cancer Cells and Peritoneal Metastasis Model. *J. Korean Med. Sci.* **2012**, *27*, 1037–1043.
158. Kopic, S.; Geibel, J.P. Gastric Acid, Calcium Absorption, and Their Impact on Bone Health. *Physiol. Rev.* **2013**, *93*, 189–268.
159. Guo, L.; Chen, W.; Zhu, H.; Chen, Y.; Wan, X.; Yang, N.; Xu, S.; Yu, C.; Chen, L. Helicobacter pylori induces increased expression of the vitamin D receptor in immune responses. *Helicobacter* **2014**, *19*, 37–47.
160. Wen, Y.; Da, M.; Zhang, Y.; Peng, L.; Yao, J.; Duan, Y. Alterations in vitamin D signaling pathway in gastric cancer progression: A study of vitamin D receptor expression in human normal, premalignant, and malignant gastric tissue. *Int. J. Clin. Exp. Pathol.* **2015**, *8*, 13176–13184.
161. Bashir, M.; Prietl, B.; Tauschmann, M.; Mautner, S.I.; Kump, P.K.; Treiber, G.; Wurm, P.; Gorkiewicz, G.; Högenauer, C.; Pieber, T.R. Effects of high doses of vitamin D₃ on mucosa-associated gut microbiome vary between regions of the human gastrointestinal tract. *Eur. J. Nutr.* **2016**, *55*, 1479–1489.
162. Vyas, N.; Companioni, R.C.; Tiba, M.; AlKhawam, H.; Catalano, C.; Sogomonian, R.; Baum, J.; Walfish, A. Association between serum vitamin D levels and gastric cancer: A retrospective chart analysis. *World J. Gastrointest. Oncol.* **2016**, *8*, 688–694.
163. Du, C.; Yang, S.; Zhao, X.; Dong, H. Pathogenic roles of alterations in vitamin D and vitamin D receptor in gastric tumorigenesis. *Oncotarget* **2017**, *8*, 29474–29486.
164. Yıldırım, O.; Yıldırım, T.; Seçkin, Y.; Osanmaz, P.; Bilgic, Y.; Mete, R. The influence of vitamin D deficiency on eradication rates of Helicobacter pylori. *Adv. Clin. Exp. Med.* **2017**, *26*, 1377–1381.
165. El Shahawy, M.S.; Hemida, M.H.; El Metwaly, I.; Shady, Z.M. The effect of vitamin D deficiency on eradication rates of Helicobacter pylori infection. *JGH Open* **2018**, *2*, 270–275.
166. Fletcher, J.; Cooper, S.C.; Ghosh, S.; Hewison, M. The Role of Vitamin D in Inflammatory Bowel Disease: Mechanism to Management. *Nutrients* **2019**, *11*, 1019.
167. Clark, I.; Smith, M.R. Effects of hypervitaminosis A and D on skeletal metabolism. *J. Biol. Chem.* **1964**, *239*, 1266–1271.
168. DeWind, L.T. Hypervitaminosis D with Osteosclerosis. *Arch. Dis. Child.* **1961**, *36*, 373–380.
169. Compston, J. Vitamin D. Molecular Biology, Physiology and Clinical Applications. *Gut* **2000**, *46*, 584.
170. Albrechtsson, E.; Jonsson, T.; Möller, S.; Höglund, M.; Ohlsson, B.; Axelsson, J. Vitamin D receptor is expressed in pancreatic cancer cells and a vitamin D₃ analogue decreases cell number. *Pancreatology* **2003**, *3*, 41–46.
171. Hewison, M. Vitamin D and the intracrinology of innate immunity. *Mol. Cell. Endocrinol.* **2010**, *321*, 103–111.
172. Sandgren, M.E.; Brönnegård, M.; DeLuca, H.F. Tissue distribution of the 1,25-dihydroxyvitamin D₃ receptor in the male rat. *Biochem. Biophys. Res. Commun.* **1991**, *181*, 611–616.
173. Nilas, L.; Christiansen, C.; Christiansen, J. Regulation of vitamin D and calcium metabolism after gastrectomy. *Gut* **1985**, *26*, 252–257.
174. Chakhtoura, M.T.; Nakhoul, N.F.; Akl, E.A.; Safadi, B.Y.; Mantzoros, C.S.; Fuleihan, G.E.-H. Vitamin D supplementation for obese adults undergoing bariatric surgery. *Cochrane Database Syst. Rev.* **2015**, doi:10.1002/14651858.CD011800.
175. Carrasco, F.; Basfi-Fer, K.; Rojas, P.; Valencia, A.; Csendes, A.; Codoceo, J.; Inostroza, J.; Ruz, M. Changes in Bone Mineral Density After Sleeve Gastrectomy or Gastric Bypass: Relationships with Variations in Vitamin D, Ghrelin, and Adiponectin Levels. *Obes. Surg.* **2014**, *24*, 877–884.
176. Eddy, R.L. Metabolic bone disease after gastrectomy. *Am. J. Med.* **1971**, *50*, 442–449.
177. Zittel, T.T.; Zeeb, B.; Maier, G.W.; Kaiser, G.W.; Zwirner, M.; Liebich, H.; Starlinger, M.; Becker, H.D. *Am. J. Surg.* **1997**, *174*, 431–438.
178. Rino, Y.; Oshima, T.; Yoshikawa, T. Changes in fat-soluble vitamin levels after gastrectomy for gastric cancer. *Surg. Today* **2017**, *47*, 145–150.
179. Schubert, M.L.; Peura, D.A. Control of Gastric Acid Secretion in Health and Disease. *Gastroenterology* **2008**, *134*, 1842–1860.
180. Feng, J.; Petersen, C.D.; Coy, D.H.; Jiang, J.-K.; Thomas, C.J.; Pollak, M.R.; Wank, S.A. Calcium-sensing receptor is a physiologic multimodal chemosensor regulating gastric G-cell growth and gastrin secretion. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 17791–17796.

181. Kantham, L.; Quinn, S.J.; Egbuna, O.I.; Baxi, K.; Butters, R.; Pang, J.L.; Pollak, M.R.; Goltzman, D.; Brown, E.M. The calcium-sensing receptor (CaSR) defends against hypercalcemia independently of its regulation of parathyroid hormone secretion. *Am. J. Physiol. Metab.* **2009**, *297*, E915–E923.
182. Christiansen, J.; Rehfeld, J.F.; Stadil, F. The effect of calcium on gastric acid and gastrin secretion in antrectomized subjects. *Gut* **1974**, *15*, 622–625.
183. Harris, S.S. Vitamin D and African Americans. *J. Nutr.* **2006**, *136*, 1126–1129.
184. Dixon, M.F. Pathology of Gastritis and Peptic Ulceration. In *Helicobacter Pylori: Physiology and Genetics*; Mobley, H.L., Mendz, G.L., Hazell, S.L., Eds.; ASM Press: Washington, DC, USA, 2001; ISBN 978-1-55581-213-3.
185. Waterhouse, M.; Hope, B.; Krause, L.; Morrison, M.; Protani, M.M.; Zakrzewski, M.; Neale, R.E. Vitamin D and the gut microbiome: A systematic review of in vivo studies. *Eur. J. Nutr.* **2018**, 1–16, doi:10.1007/s00394-018-1842-7.
186. Gigeck, C.O.; Chen, E.S.; Calcagno, D.Q.; Wisnieski, F.; Burbano, R.R.; Smith, M.A.C. Epigenetic mechanisms in gastric cancer. *Epigenomics* **2012**, *4*, 279–294.
187. De Ruijter, A.J.; Van Gennip, A.H.; Caron, H.N.; Kemp, S.; Van Kuilenburg, A.B. Histone deacetylases (HDACs): Characterization of the classical HDAC family. *Biochem. J.* **2003**, *370*, 737–749.
188. Meshorer, E.; Misteli, T. Chromatin in pluripotent embryonic stem cells and differentiation. *Nat. Rev. Mol. Cell Biol.* **2006**, *7*, 540–546.
189. Dokmanovic, M.; Clarke, C.; Marks, P.A. Histone Deacetylase Inhibitors: Overview and Perspectives. *Mol. Cancer Res.* **2007**, *5*, 981–989.
190. Jones, P.A. The Role of DNA Methylation in Mammalian Epigenetics. *Science* **2001**, *293*, 1068–1070.
191. Gowher, H.; Jeltsch, A. Mechanism of inhibition of DNA methyltransferases by cytidine analogs in cancer therapy. *Cancer Boil. Ther.* **2004**, *3*, 1062–1068.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).