Supplementary table 1. Number and percentage of animals with change in body condition, coat and grooming, and ears, eyes and whiskers in different groups.

Group		Body Condition					Coat and grooming					Eyes, Ears and Whiskers				
			-1 Week	0 Week	1 Week	2 Week		-1 Week	0 Week	1 Week	2 Week		-1 Week	0 Week	1 Week	2 Week
I (n = 10)	Wild-type parecoxib	0	10 (100.0%)	10 (100.0%)	10 (100.0%)	10 (100.0%)	0	10 (100.0%)	10 (100.0%)	10 (100.0%)	10 (100.0%)	0	10 (100.0%)	10 (100.0%)	10 (100.0%)	10 (100.0%)
		1					1					1				
		2					2					2				
							3									
II (n = 11)	Wild-type control	0	11 (100.0%)	11 (100.0%)	11 (100.0%)	11 (100.0%)	0	11 (100.0%)	11 (100.0%)	11 (100.0%)	11 (100.0%)	0	11 (100.0%)	11 (100.0%)	11 (100.0%)	11 (100.0%)
		1					1					1				
		2					2					2				
							3									
III (n = 11)	HPV parecoxib	0	11 (100.0%)	8 (72.7%)	8 (72.7%)	7 (63.6%)	0	11 (100.0%)	11 (100.0%)	9 (81.8%)	10 (90.9%)	0	10 (90.9%)	10 (90.9%)	8 (72.7%)	7 (63.6%)
		1		3 (27.3%)	3 (27.3%)	4 (36.4%)	1			2 (18.2%)	(9.10%)	1	1 (9.10%)	(9.10%)	3 (27.3%)	4 (36.4%)
		2					2					2				
							3									
IV (n = 11)	HPV control	0	11 (100.0%)	10 (90.9%)	10 (90.9%)	10 (90.9%)	0	11 (100.0%)	11 (100.0%)	10 (90.9%)	8 (72.7%)	0	11 (100.0%)	11 (100.0%)	10 (90.9%)	11 (100.0%)
		1		1 (9.10%)	1 (9.10%)	(9.10%)	1			1 (9.10%)	3 (27.3%)	1			1 (9.10%)	
		2					2					2				
							3									