



1 Supporting information

2 High-Throughput Screening for the Optimization of 3 D-Glycerate Dehydratase Activity


4 Benjamin Begander ¹, Anna Huber ¹, Manuel Döring ¹, Josef Sperl ¹ and Volker Sieber ^{1,2,3,*}

5 ¹ Chair of Chemistry of Biogenic Resources, Campus Straubing for Biotechnology and Sustainability,
6 Technical University of Munich, Schulgasse 16, D-94315 Straubing, Germany; benjamin.begander@tum.de
7 (B.B.); anna97.huber@tum.de (A.H.); manuel.doering@tum.de (M.D.); josef.sperl@tum.de (J.S.)

8 ² Catalysis Research Center, Technical University of Munich, 85748 Garching, Germany

9 ³ School of Chemistry and Molecular Biosciences, The University of Queensland, St. Lucia, QLD 4072,
10 Australia

11 * Correspondence: sieber@tum.de; Tel.: +49-9421-187-300



12

13 **Figure S1.** Effect of spiking with NADH on the absorption difference of pyruvate standard samples.
14 standard = pyruvate in water; w/ NADH = pyruvate in water, spiked with the indicated concentration
15 of NADH.

16



© 2019 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

17