
Supplementary Figure 1.


Dotplots showing the individual levels of CSF biomarkers by ASIA grade.

- 600 • 9 ŝ IL-16 200 400 IL-15 1L-13 3 4 N C C B ASIA B ASIA Ď B ASIA А D А А D IL-2Ra 10 15 20 25 IL-6 0 40000 60000 80000 8 • • 9 IL-18 4 20000 2 . • S 8. 0 • . B D ASIA B ASIA B ASIA D A Ď А А IL-7 20 30 40 50 150 30000 . . . IL-9 50 100 IL-8 10000 20000 9 • 0 • 0 0 B ASIA B ASIA B ASIA D A D A D А IP-10 10000 20000 30000 150 • CTACK 10 20 30 40 50 60 RANTES 50 100 2 • • . 8 2 0 0 B ASIA A B ASIA D D B ASIA D А А MCP-1 1000 2000 3000 MIP-1b 0 500 100015002000 80 . MIP-1a . BASIA . 1 0 0 00 BASIA D Å B ASIA Ď Å A D 8. - 25 . . . 10 TRAIL 0 15 20 IFN-a2 TNF-a 10 4 . 0 5 B ASIA B ASIA Ď Å Ď B D ASIA Å Å
- a) Cytokines and chemokines

b) Growth and other factors



c) Soluble cell adhesion molecules and Neurological biomarkers



d) Other Biomarkers.

