

Supplementary data

Article

The antimicrobial peptide Temporin G: anti-biofilm, anti-persister activities and potentiator effect of tobramycin efficacy against *Staphylococcus aureus*

Bruno Casciaro^{1,*}, Maria Rosa Loffredo², Floriana Cappiello², Guendalina Fabiano², Luisa Torrini², Maria Luisa Mangoni^{2,*}

¹ Center For Life Nano Science@Sapienza, Istituto Italiano di Tecnologia, Viale Regina Elena 291,

00161 Rome, Italy (B.C.);

² Laboratory affiliated to Pasteur Italia-Fondazione Cenci Bolognetti, Department of Biochemical Sciences, Sapienza University of Rome, P.le Aldo Moro 5, 00185 Rome, Italy (M.R.L, F.C., G.F., L.T., M.L.M.);

* Correspondence: bruno.casciaro@iit.it, (B.C.); marialuisa.mangoni@uniroma1.it; Tel.: +39 0649910838 (M.L.M.);

Supplementary data.

Table S1. Minimum inhibitory concentrations of the tested compounds after 20 h treatment.

Strain	Rifampicin (ng/mL)	Tobramycin (µg/mL)	TG (µM)
<i>S. aureus</i> ATCC 25923	1.9	0.5	25
<i>S. aureus</i> 1a	1.9	64	12.5
<i>S. aureus</i> 1b	1.9	64	25
<i>S. aureus</i> 1c	R	32	12.5

R=resistant, >250 µg/mL

Table S2. Resistance phenotypes of the bacterial strains

Strain	Resistance phenotype ^a
<i>S. aureus</i> 1a	AMC-AMP-CIP-ERY-GEN-OXA
<i>S. aureus</i> 1b	AMC-AMP-CIP-ERY-GEN-OXA
<i>S. aureus</i> 1c	AMC-AMP-CIP-ERY-GEN-OXA-RIF

AMC, amoxicillin; AMP, ampicillin; CIP, ciprofloxacin; erythromycin; GEN, gentamicin; oxacillin; RIF, rifampicin.