

**Supplementary Table S1: Demographic and clinical parameters of the 27 Hispanic study participants with HCC.** Data are presented as frequency (%) or as mean (range) - median. BMI: body mass index; AFP: alpha-fetoprotein.

<b>Parameters</b>	
<b>Male (n=27)</b>	18 (66.7%)
<b>Age (n=27)</b>	66.9 (50.0-88.0) - 68.0
<b>BMI (n=25)</b>	31.2 (23.2-50.7) - 29.0
<b>Obese (BMI <math>\geq 30</math>)</b>	10 (40%)
<b>Diabetes (n=27)</b>	20 (74.1%)
<b>Family History of Cancer (n=27)</b>	13 (48.1%)
<b>Stage (n=27)</b>	
<b>I</b>	5 (18.5%)
<b>II</b>	4 (14.8%)
<b>III</b>	8 (29.6%)
<b>IV</b>	10 (37.0%)
<b>Child-Pugh Score (n=27)</b>	
<b>A</b>	20 (74.1%)
<b>B</b>	7 (25.9%)
<b>Multiple tumors (n=25)</b>	15 (55.6%)
<b>AFP (n=26)</b>	3857.2 (2.8-36000.0) - 89.8
<b>Differentiation (n=17)</b>	
<b>Well</b>	8 (47.1%)
<b>Moderate</b>	5 (29.4%)
<b>Poor</b>	4 (23.5%)