

Supplementary Materials: Somatic Copy Number Alterations and Associated Genes in Clear-Cell Renal-Cell Carcinoma in Brazilian Patients



Flávia Gonçalves Fernandes¹, Henrique Cesar Santejo Silveira¹, João Neif Antonio Júnior², Rosana Antunes da Silveira¹, Luis Eduardo Zucca², Flavio Mavignier Cárcano^{2,3}, André Octavio Nicolau Sanches², Luciano Neder⁴, Cristovam Scapulatempo-Neto⁴, Sergio Vicente Serrano^{2,3}, Eric Jonasch⁵, Rui Manuel Reis^{1,6,7*}  and Adriane Feijó Evangelista^{1*} 

Table S1. Summary of the main risk factors associated with ccRCC patients.

Clinical Data	Category	Patients (Frequency)
Obesity	Yes	31 (33.6%)
	No	61 (66.3%)
Smoke	Yes	14 (15.2%)
	No	56 (60.8%)
	Former smoker	17 (18.4%)
	Unknown	5 (5.4%)
Alcohol	Yes	19 (20.6%)
	No	67 (72.8%)
	Unknown	6 (6.5%)
Conditions	Diabetes	21 (22.8%)
	Hypertension	50 (54.3%)
	Diabetes and Hypertension	20 (21.7%)
	Other	8 (8.6%)




Figure S1. Overall survival of the 92 ccRCC patients.




Figure S2. Kaplan Meier plots according to stage (A), histological grade (B), metastasis (C) and tumor size (D).