

SUPPLEMENTARY FIGURES

Single-walled carbon nanotubes modify leaf micromorphology, chloroplast ultrastructure and photosynthetic activity of pea plants

Violeta Velikova ^{1, *}, Nia Petrova ², László Kovács ³, Asya Petrova ¹, Dimitrina Koleva ⁴, Tsonko Tsonev ², Stefka Taneva ², Petar Petrov ⁵, Sashka Krumova ^{2, *}

¹ Institute of Plant Physiology and Genetics, Bulgarian Academy of Sciences, Acad G. Bonchev Str. Bl. 21, 1113 Sofia, Bulgaria; e-mail@ violet@bio21.bas.bg (V.V.), petrova_asya@abv.bg (A.P.)

² Institute of Biophysics and Biomedical Engineering, Bulgarian Academy of Sciences, Acad G. Bonchev Str. Bl. 21, 1113 Sofia, Bulgaria; e-mail@ zlatkova.nia@gmail.com (N.P.), tsonev@gmail.com (T.T.), sgtaneva@gmail.com (S.T.), sashka.b.krumova@gmail.com (S.K.)

³ Institute of Plant Biology, Biological Research Center, Temesvári krt. 62, 6726 Szeged, Hungary; e-mail@ kovacs.laszlo@brc.hu

⁴ Faculty of Biology, Sofia University “St. K. Ohridski”, 8 Dragan Tsankov, 1164 Sofia, Bulgaria; e-mail@ koleva@biofac.uni-sofia.bg

⁵ Institute of Polymers, Bulgarian Academy of Sciences, Acad G. Bonchev Str. Bl. 103, 1113 Sofia, Bulgaria; e-mail@ ppetrov@polymer.bas.bg

* Correspondence: e-mail@ violet@bio21.bas.bg (V.V.), sashka.b.krumova@gmail.com (S.K.)




Figure S1. Cuticle and epicuticular waxes at the adaxial (A) and abaxial (B) leaf surface of pea plants seven days after spraying with P₃₀₀.




Figure S2. Representative electron micrographs of chloroplast cross section taken from intact pea leaves seven days after spraying with different concentrations of “Pluronic” P-85 (A, P₁₀; B, P₁₀₀; P₃₀₀, C). G – grana, S – stroma lamellae.