

Supplementary Table S1. Sites of lymphadenopathy and biopsy sites in iMCD-NOS cases.

| Subtype | Age/Sex | Biopsy site | Site of lymphadenopathy | Extranodal lesion |
|---------|---------|-----------------------------|--|---|
| IPL | 47/M | inguinal | mediastinum, hilar, inguinal | multiple lung nodules |
| | 35/F | cervical | cervical, supraclavicular fossa, axillary, para-aortic, mesenterium, inguinal | |
| | 55/M | inguinal | mediastinum, inguinal | multiple lung nodules |
| | 62/M | -* | systemic, unspecified * | multiple lung nodules, kidney |
| | 55/F | inguinal | systemic, unspecified * | |
| | 76/M | cervical | cervical, axillary, inguinal | skin |
| | 61/F | axillary | systemic, unspecified * | |
| | 37/M | -* | cervical, axillary, abdominal | |
| | 39/M | inguinal | systemic, unspecified * | multiple lung nodules |
| | 49/F | mediastinum | infraclavicular, mediastinum, axillary, inguinal | |
| | 48/F | axillary | axillary | multiple lung nodules |
| | 59/M | cervical | cervical, axillary, abdominal, inguinal | multiple lung nodules, skin |
| | 64/M | inguinal | cervical, axillary, mesenterium, pelvis, inguinal | |
| | 70/M | cervical | cervical, supraclavicular fossa, mediastinum, axillary, para-aortic | |
| | 54/F | cervical | cervical, axillary | multiple lung nodules |
| | 65/M | cervical | systemic, unspecified * | |
| | 62/F | cervical | cervical (bilateral, multiple) | tonsil swelling |
| | 37/M | inguinal | axillary, inguinal | splenomegaly |
| | 55/F | -* | cervical, supraclavicular fossa, mediastinum, axillary, para-aortic, abdominal, inguinal | |
| | 52/F | axillary | mediastinum, hilar, axillary | |
| | 74/F | mediastinum | mediastinum, abdominal | multiple lung nodules |
| | 34/M | mediastinum | cervical, mediastinum, hilar | hepatosplenomegaly |
| | 57/M | inguinal | cervical, axillary, para-aortic, iliac, inguinal | |
| | 59/M | cervical | cervical, axillary | |
| | 52/F | inguinal | cervical, para-aortic, inguinal | |
| | 70/M | cervical | cervical, mediastinum, para-aortic, | |
| | 43/M | inguinal | supraclavicular fossa, inguinal | |
| | 41/M | cervical | cervical, supraclavicular, axillary, inguinal, | skin |
| | 49/M | axillary | cervical, supraclavicular, mediastinum, axillary, inguinal | hepatosplenomegaly, kidney |
| | 48/F | axillary | axillary, para-aortic, abdominal | hepatosplenomegaly |
| | 38/F | inguinal | cervical, mediastinum, inguinal | multiple lung nodules, splenomegaly, skin |
| | 76/M | cervical | cervical, subclavicular, axillary, para-aortic, pelvis, inguinal | |
| | 67/M | inguinal | axillary, inguinal | |
| | 72/M | cervical | cervical (bilateral, multiple) | |
| non-IPL | 70/M | supraclavicular fossa | supraclavicular, mediastinum | |
| | 52/F | axillary | cervical, axillary, inguinal | kidney |
| | 89/M | para-aortic (needle biopsy) | mediastinum, retroperitoneum (multiple) | |
| | 73/F | axillary | mediastinum, axillary | |
| | 49/F | axillary | systemic, unspecified * | multiple lung nodules, skin |
| | 32/M | cervical | systemic, unspecified * | hepatosplenomegaly |
| | 49/F | cervical | systemic, unspecified * | kidney, skin |
| | 49/F | cervical | systemic, unspecified * | kidney |

Multiple lymphadenopathies were noted in all cases. Sampling site was unknown in 3 cases of IPL group. All cases had excisional biopsy except for one non-IPL case who had a core needle biopsy. * Details unknown.