

Table S1. Primer sequence for qRT-PCR

primers	Forward (5'-3')	Reverse(5'-3')
<i>GAPDH</i>	GGCATTGCTCTCAATGACAA	TGTGAGGGAGATGCTCAGTG
<i>Nestin</i>	GGAGGGCAGAGAAGACAGTG	TGACATCCTGGACCTTGACA
<i><math>\beta</math>-actin</i>	GGCTGTATTCCCCTCCATCG	TAAAGACCTCTATGCCAACACAGT
<i>GFAP</i>	CACGAACGAGTCCCTAGAGC	TCACATCACCACGTCCTTGT
<i><math>\beta</math>III-tubulin</i>	GAATGACCTGGTGTCCGAGT	CAGAGCCAAGTGGACTCACA
<i>Cyclin D2</i>	CTGTGCGCTACCGACTTCAA	GCAGAGCTTCGATTTGCTCC
<i>Rasal2</i>	ATGGAGCTGTCTCCGTCGT	GCCTTTTACATCGAACACCCG
<i>Cyclin D1</i>	CCCTGACACCAATCTCCTCA	TCTTCTTCAAGGGCTCCAGG
<i>Cdkn1B</i>	TCAAACGTGAGAGTGTCTAACG	CCGGGCCGAAGAGATTTCTG
<i>Trp53bp2</i>	AGTAAAGGCTCTAAAGCTCACCC	GTAAGAGGTCGGCATTGGAAG
<i>Ccnyl1</i>	GAGAGGCTCCTAACTTATGCTGA	CCTGGTCATCCCAAACCTTG
<i>Cyclin B1</i>	AAGGTGCCTGTGTGTGAACC	GTCAGCCCCATCATCTGCG
<i>Cyclin A2</i>	GCCTTCACCATTCATGTGGAT	TTGCTGCGGGTAAAGAGACAG

Figure S1




Figure S2




Figure S3

