

Effect of the Synthetic Approach on the Formation and Magnetic Properties of Iron-Based Nanophase in Branched Polyester Polyol Matrix

Artur Khannanov ¹, Anastasia Burmatova ¹, Klara Ignatyeva ¹, Farit Vagizov ², Airat Kiyamov ^{2*}, Dmitrii Tayurskii ², Mikhail Cherosov ², Alexander Gerasimov ¹, Evtugyn Vladimir ³ and Marianna Kutyreva ¹

¹ Butlerov Chemistry Institute, Kazan Federal University, Kazan 420008, Russia

² Institute of Physics, Kazan Federal University, Kazan 420008, Russia

³ Interdisciplinary Center "Analytical Microscopy", Kazan Federal University, Kazan 420008, Russia

* Correspondence: ajrat.kiyamov@kpfu.ru




Figure S1. Theoretical structure of Boltorn H20






Figure S2. A) ChemRed-NP Change in the color of solutions during the synthesis of $\text{Fe}_m\text{O}_n/\text{H}_2\text{O}$ nanoparticles with the addition of $\text{N}_2\text{H}_4 \times \text{H}_2\text{O}$ and color of the dispersed system containing composite $\text{Fe}_m\text{O}_n/\text{H}_2\text{O}$ nanoparticles at the end of the synthesis reaction; B) TermRed-NP reaction mixture during washing before centrifugation; C) US/TermRed-NP reaction mixture during washing before centrifugation;; D) US-NP after centrifugation.




FigureS3. SEM image of US/TermRed-NP nanoparticles to confirm the statistics.




MDC AM 1 μm Mag = 15.00 K X Signal A = STEM EHT = 25.00 kV I Probe = 300 pA
STEM Q1 = Off STEM Q2 = Inverted STEM Q3 = Inverted


FigureS4. SEM image of US-NP nanoparticles to confirm the statistics.




FigureS5. Ft-IR spectra: A) BH20; B) ChemRed-NP; C) TermRed-NP; D) US/TermRed-NP; E) US-NP.



FigureS6 Full XRD spectra of **ChemRed-NP**.



FigureS7 Full XRD spectra of **US-NP**.



FigureS8. Full XRD spectra of **US/TermRed**.




Figure S9 Mossbauer spectra of US/TermRed-NP with deconvolution: A) at 300 K; B) at 77 K.




Figure S10 Mossbauer spectra of US-NP with deconvolution: A) at 300 K; B) at 80 K.





Figure S11 Mossbauer spectra of ChemRed-NP with deconvolution: A) at 76 K; B) at 180 K; C) at 300 K ; D) Isomer shifts at different temperatures; E) quadrupole splitting; F) Thermal deposition of ChemRed-NP



Figuere S12. NTA analysis ChemRed-NP




Figure S13. NTA analysis TermRed-NP




Figure S14. NTA analysis US/TermRed-NP




Figure S15. NTA analysis US-NP.