

Combined Model-Based Prediction for Non-Invasive Prenatal Screening

So-Yun Yang ^{1,†}, Kyung Min Kang ^{2,†}, Sook-Young Kim ³, Seo Young Lim ⁴, Hee Yeon Jang ^{1,2}, Kirim Hong ⁵, Dong Hyun Cha ^{2,5}, Sung Han Shim ^{1,2,*} and Je-Gun Joung ^{1,3,6*}

- ¹ Department of Biomedical Science, College of Life Science, CHA University, Seongnam, Gyunggi-do 13488, Korea
- ² Center for Genome Diagnostics, CHA Biotech Inc, Seoul 06135, Korea
- ³ CHA Future Medicine Research Institute, CHA Bundang Medical Center, Seongnam, Gyunggi-do 13520, Korea
- ⁴ Department of Biomedical Engineering, Hankuk University of Foreign Studies, Seoul, Korea
- ⁵ Department of Obstetrics and Gynecology, CHA Gangnam Medical Center, CHA University, Seoul 06135, Korea
- ⁶ Institute for Biomedical Informatics, CHA University School of Medicine, CHA University, Seongnam, Republic of Korea.
- * Correspondence: e-mail@e-mail.com; Tel.:
- † These authors contributed equally to this work




Figure S1. Z-scores of chromosomes 13, 18, and 21 calculated using three different methods.

Table S1. Number of trisomies predicted using three different methods.

| | STD | NCV | WSRB | Confirmation |
|------------|-----|-----|------|--------------|
| Trisomy 13 | 5 | 5 | 3 | 0 |
| Trisomy 18 | 9 | 10 | 9 | 5 |
| Trisomy 21 | 33 | 34 | 33 | 33 |
| Total | 47 | 49 | 45 | 38 |