

Complex Formation between Cytochrome *c* and a Tetra-alanino-calix[4]arene

Stefano Volpi ¹, Aishling Doolan ², Laura Baldini ^{1,*}, Alessandro Casnati ¹, Peter B. Crowley ² and Francesco Sansone ¹

¹ Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale, Università degli Studi di Parma, Viale delle Scienze, 17/A, 43124 Parma, Italy

² School of Biological and Chemical Sciences, University of Galway, University Road, H91 TK33 Galway, Ireland

* Correspondence: laura.baldini@unipr.it

Supporting Information

Table of contents

Figure S1: ¹H NMR spectrum (D₂O, 25°C, 500 MHz) of compound **1** (page S2).

Figure S2: Overlaid ¹H-¹⁵N HSQC full-spectra of 0.1 mM cytochrome *c* in the absence or in the presence of 0.1–3.1 mM **1** (page S2).

Figure S3: Gel formation observed during crystallization tests for the cytochrome *c* - **1** complex (page S3).




Figure S1: ^1H NMR spectrum (D₂O, 25°C, 500 MHz) of compound **1**.




Figure S2: Overlaid ^1H - ^{15}N HSQC full-spectra of 0.1 mM cytochrome *c* in the absence (black contours) or presence of 0.1–3.1 mM **1** (blue scale).




Figure S3: Gel formation observed during crystallization tests for the cytochrome *c* - 1 complex performed by **a)** a robot sitting drop and **b)** manual hanging drop vapor diffusion method (1.7 mM protein, 10 mM ligand). The reservoir solutions were **a)** 20% PEG 6000, 1M LiCl, 100mM sodium citrate buffer (pH 4) and **b)** 25% PEG 8000, 50mM sodium citrate buffer (pH 5).