



# **Integrin Signaling in the Central Nervous System in Animals and Human Brain Diseases**

Hiroko Ikeshima-Kataoka <sup>1,2,\*</sup>, Chikatoshi Sugimoto <sup>1</sup> and Tatsuya Tsubokawa <sup>1</sup>

- <sup>1</sup> Department of Biology, Keio University, 4-1-1, Hiyoshi, Kohoku-ku, Yokohama-shi 223-8521, Japan; chikatoshi.sugimoto@keio.jp (C.S.); tsubo@a8.keio.jp (T.T.)
- <sup>2</sup> Faculty of Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan
- Correspondence: ikeshima@1988.jukuin.keio.ac.jp

Abstract: The integrin family is involved in various biological functions, including cell proliferation, differentiation and migration, and also in the pathogenesis of disease. Integrins are multifunctional receptors that exist as heterodimers composed of  $\alpha$  and  $\beta$  subunits and bind to various ligands, including extracellular matrix (ECM) proteins; they are found in many animals, not only vertebrates (e.g., mouse, rat, and teleost fish), but also invertebrates (e.g., planarian flatworm, fruit fly, nematodes, and cephalopods), which are used for research on genetics and social behaviors or as models for human diseases. In the present paper, we describe the results of a phylogenetic tree analysis of the integrin family among these species. We summarize integrin signaling in teleost fish, which serves as an excellent model for the study of regenerative systems and possesses the ability for replacing missing tissues, especially in the central nervous system, which has not been demonstrated in mammals. In addition, functions of astrocytes and reactive astrocytes, which contain neuroprotective subpopulations that act in concert with the ECM proteins tenascin C and osteopontin via integrin are also reviewed. Drug development research using integrin as a therapeutic target could result in breakthroughs for the treatment of neurodegenerative diseases and brain injury in mammals.

**Keywords:** astrocyte; axon; CNS; invertebrate; mammal; mouse; neuron; phylogenetic tree; regeneration; teleost fish

# 1. Introduction

Integrins are multifunctional receptors mainly responsible for binding to extracellular matrix (ECM) proteins that mediate cell adhesion. Cell-cell and cell-ligand adhesions were observed and determined to be important processes based in tissue culture studies conducted by Harrison in the early 20th century [1]. Tissue culture techniques came to be widely used for cancer research in the 1960s and early 1970s. In one study reported, in 1976, that when fibronectin, among the earliest of the identified ECM proteins, was added to a culture of viral-transformed cells, the cells reverted to their normal phenotype. In 1978, receptors for fibronectin were biochemically purified as a 140 kD protein complex and immunologically identified by monoclonal antibodies, and were named integrins, to reflect their integral membrane-protein nature and their role in linking the cytoskeleton to the ECM (fibronectin) [2,3]. Then, in 1986, the cDNA of the  $\beta$  integrin subunit was cloned. Bioassays have revealed that the Arg-Gly-Asp-Ser (RGDS) motif is sufficient to inhibit the binding of fibronectin to integrins [4]. Monoclonal antibodies directed against lymphocytes have been shown to share a common subunit and adhesion-related processes with leucocyte-function-associated antigen (LFA1) and macrophage 1 antigen (MAC1). The Drosophila melanogaster position-specific antigen (PS) was characterized as a heterodimeric complex that shared a common subunit. All of these findings were reported between 1986 and 1987 [5-7].

In regard to their molecular structure, integrins are composed of a large extracellular domain that binds to the ECM and other proteins, a transmembrane domain, and a short



Citation: Ikeshima-Kataoka, H.; Sugimoto, C.; Tsubokawa, T. Integrin Signaling in the Central Nervous System in Animals and Human Brain Diseases. *Int. J. Mol. Sci.* 2022, 23, 1435. https://doi.org/10.3390/ ijms23031435

Academic Editors: Anthony Lemarié and Sylvie Monferran

Received: 28 December 2021 Accepted: 25 January 2022 Published: 27 January 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). intercellular domain that interacts with signal transduction proteins composed of about 20–60 amino acids. However, the cytoplasmic domain of integrin  $\beta$ 4 is unusually large (about 1000 a.a.) that interact with cytoskeletal components of hemidesmosomes [8]. The fully functional integrins exist as heterodimers composed of one  $\alpha$  integrin subunit and one  $\beta$  integrin subunit. Currently, mammals have been reported to have 18 different  $\alpha$  subunits and 8  $\beta$  subunits of integrin that form 24  $\alpha/\beta$  pairs in a specific way [9]. Some pairs ( $\alpha$ 1 $\beta$ 1, etc.) are known as collagen receptors, others ( $\alpha$ D $\beta$ 2, etc.) as leucocyte integrins, and some others ( $\alpha$ 9 $\beta$ 1, etc.) as osteopontin receptors. In the regeneration of the peripheral nervous system (PNS), ID1/3 transcription factors regulate the expression of  $\alpha$ 6 integrin, which has been shown to mediate interactions between axons and skin fibroblasts to promote axon extension [10]. In central nervous system (CNS) regeneration, fibrotic scars formed by fibroblasts may exert inhibitory effects both via the production of inhibitory molecules (CSPG, collagen, etc.) and by physical abrogation of aligned pathways for regenerating axons crossing the lesion [11].

In mammals, some of the integrin receptors are involved in peripheral nerve regeneration [12,13]. The localization of the integrin family in the adult nervous system and the roles of integrins in axonal regeneration have been elegantly summarized in some review articles [14,15]. Integrin function in concert with ECM in synaptic plasticity and memory are important in the brain, and the roles of integrins in brain diseases have been summarized [16].

Integrins have various roles in neuronal connectivity and development of the brain [17,18]. In the adult mammalian CNS, injured axons fail to regenerate; while they have the capacity to regenerate during development, this ability is lost with maturity. Many researchers have focused on the molecules that contribute to CNS development to induce regeneration of the CNS after injury, however, good results have not yet been obtained. For example, the axonal growth-promoting integrins continue to be transported until adulthood in the axons of the PNS, while such transport declines with maturity in axons of the CNS. In fact, integrin and Rab11, a regulator of recycling endosomes, manipulated into mature CNS axons in vitro, enabled axonal regeneration after laser injury, but long-distance regeneration in vivo remains to be observed [13].

In the mammalian CNS, astrocytes are among the most abundant of glial cells that maintain the physiological integrity of the blood brain barrier (BBB), and play roles in glucose sensing (so-called astrocyte-neuron lactate shuttle hypothesis), blood flow regulation, synaptic functioning at tripartite synapses, gliotransmission, etc. [19–21]. Recently, we summarized the metabolic plasticity of astrocytes and aging of the brain in a review article [22]. The distribution and expressions of three astrocytic markers, namely, glial fibrillary acidic protein (GFAP), Ca<sup>2+</sup> binding peptide (S100 $\beta$ ), and N-Myc downstream-regulated gene 2 (NDRG2), in different mouse cerebral regions has been reported [23]. Since astrocytes also express integrins and are potentially involved in CNS regeneration and development of neuroimmunological diseases, we focus on the role of astrocytes participating in the integrin signaling in the following sections.

### 2. Integrins and Regeneration Studies in Invertebrates and Fish

All multicellular animals express integrins that exist as heterodimers composed of one  $\alpha$  and one  $\beta$  subunit. The fully sequenced genome of the fruit fly *Drosophila melanogaster* contains five  $\alpha$  and two  $\beta$  integrin subunits. The nematode roundworm *Caenorhabditis elegans* has been shown to have two orthologues of  $\alpha$  integrin (INA-2, PAT-2) and one ortholog of  $\beta$  integrin (PAT-3) [24]. Unicellular organisms also have integrins, a recent genetic analysis had shown that many of the key genes of integrin adhesion complex that had formerly been cited as crucial for metazoan origins have a much earlier origin [25].

On the basis of a phylogenetic reconstruction, Johnson et al. [24] divided vertebrate and invertebrate animal  $\alpha$  integrins into six clusters, termed PS1 (vertebrate  $\alpha$ 3, 6, and 7), PS2 (vertebrate  $\alpha$ 5, IIb, V, and  $\beta$ ), PS3, PS4 (vertebrate  $\alpha$ 4/9), collagen receptors (vertebrate  $\alpha$ 1, 2, 10, and 11), and leucocyte integrins of the immune system (vertebrate  $\alpha$ E, L, M,

D, and X). The PS3 group seems to be specific for insects. The PS4 group includes two mammalian  $\alpha$  subunits, namely  $\alpha$ 4 and  $\alpha$ 9, can be considered to be a separate subgroup. These integrins recognize, in addition to ECM proteins, certain plasma proteins such as osteopontin, receptors belonging to the immunoglobulin superfamily, and vascular endothelial cell growth factors.

We have researched TN-C and OPN [26–31], and have revealed that either of the molecules are indispensable for mammalian brain functions in concerted with  $\alpha$ 9 and  $\beta$ 1 integrin as the receptors (see Sections 4.4 and 4.5). Furthermore, recently, it has been reported that  $\alpha$ 9 integrin could be a therapeutic target for nerve injury [32]. Thus, we focused on  $\alpha$ 9 integrin and a subgroup member  $\alpha$ 4 integrin, and performed a molecular phylogenetic tree analysis for integrins of the PS4 cluster. We clearly showed that the vertebrate  $\alpha$ 4 subunit and  $\alpha$ 9 subunit, which are distinct from the invertebrate  $\alpha$ 4 subunit and  $\alpha$ 9 subunit, which are distinct from the  $\alpha$ 4 subunit before the appearance of the first vertebrates (Figure 1). While integrin  $\alpha$ 9 is thought to be vertebrate-specific, integrins of the PS4 group (including Drosophila PS4) are independent from other PS groups and might have common functions in the CNS (see below).



**Figure 1.** Phylogenetic relationship of PS4 integrin group. Protein sequences of the vertebrate integrins  $\alpha 4$  and  $\alpha 9$  and invertebrate integrins ( $\alpha$  ina-1 and pat-2 of *Caenorhabditis elegans*,  $\alpha 4$  of *Octopus sinensis*,  $\alpha PS4$  of *Drosophila melanogaster*, and  $\alpha$  int-4 of *Schmidtea mediterranea*) obtained from the NCBI were aligned using the ClustalW algorithm, and a phylogenetic tree was drawn with Neighbor-Joining on MEGA11 [33].

Johnson et al. [24] divided the vertebrate and invertebrate animal  $\beta$  integrins into three clusters, namely, vertebrate A ( $\beta$ 1, 2, and 7), vertebrate B ( $\beta$ 3, 4, 5, 6, and 8), and invertebrate clusters.

### 2.1. Invertebrate (Planaria, Nematode, Fruit Fly, and Cephalopods)

Regeneration is a process that is ubiquitous among multicellular organisms. Some species of planarian flatworms (including the freshwater triclad, *Schmidtea mediterranea*) show autotomy, namely, when one adult flatworm is cut into two, it grows into two adults, and animals have adult regeneration ability for need to regulate, using specific signaling, cell proliferation, specification and patterning induced by injury. Bonar and Peterson identified that inhibition of one of four planarian integrin-alpha subunits, that is,  $\alpha$  integrin 2, resulted in phenocopying of the regeneration of the normal brain architecture, but without normal cell-type regionalization, suggesting that a specific receptor controls brain organization through regeneration [34].

The nematode (roundworm) *Caenorhabditis elegans* is a well-known model for almost complete identification of the whole adult cell lineage and a fast life cycle. It has been reported that the TNS-1 (*C. elegans* homolog of vertebrate tensin) PTB domain is required for its association with the cytoplasmic region of the integrin  $\beta$  subunit PAT-3, and that this integrin pathway also promotes axonal regeneration through activation of the JNK cascade [35]. It has also been demonstrated that the integrin  $\alpha$  subunit INA-1 activates the JNK pathway through the signaling complex CED-2–CED-5–CED-12, to promote axonal regeneration [36], and that the *C. elegans* integrin signaling network regulates axonal regeneration via the Src–RhoGEF–RhoA axis [37]. Single cell lineage analysis has also shown glia-derived neurons in adults whose progenitor cells are fully differentiated glia [38].

The fruit fly *Drosophila melanogaster* has also been shown to be an excellent model for genetic research. Drosophila neurons express integrins, heterodimeric cell surface receptors consisting of an  $\alpha$  subunit and a  $\beta$  subunit. Neuron-specific overexpression of myospheroid (mys, a  $\beta$  integrin subunit) and multiple edematous (mew, an  $\alpha$  integrin subunit) increase dendrite adhesion to the ECM [39].

Cephalopods (including the east Asian common octopus (*Octopus sinensis*)) are not only an attractive target for ethologic studies of animals with higher intelligence, but also a promising model for research in genomics [40]. Since cephalopods show regeneration of the arms, shell, and nerves [41,42], integrins of cephalopods may also contribute to these regenerations, as in vertebrates.

### 2.2. Vertebrates (Fish)

Many animals are used as models, but teleost fish are an excellent model for the study of regenerative systems of vertebrates and possess the ability to replace missing organs. These fish work as an ideal pathological model because of higher similarities with humans, neuroanatomical, neurochemical, and emotional/social behaviorally.

The CNS neurons of lower vertebrates, such as fish and amphibians, have been known to have the ability to regenerate after axotomy since Sperry's work in the 1950s [43]. In contrast to the case in mammals and birds, neurons in the CNS of fish, such as the retinal ganglion cells, are capable of axonal regeneration and vision restoration. The axonal regeneration seems to be accomplished successfully in the fish visual pathway with some particular properties of the glial cells and neurons [44].

Mammalian neurons have been shown to express enhanced neurite outgrowth on salmon fibrin gel via the mediation of integrins [45]. Integrin of not only mammalian neurons, but also fish microglia and macrophages have been revealed to be involved in self optic nerve regeneration in fish [46,47]. Inhibition of tenascin C (TN-C) ameliorated axon regeneration and recovered the swimming behavior after spinal cord injury in adults, indicating that inhibitors of axonal regeneration are not completely lacking in the CNS environment of fish. Thus, similar to chondroitin sulfate proteoglycans (CSPGs), tenascin R may play a role in guidance during axonal regeneration. The lack of inhibitory molecules or cells and the presence of positive factors may induce CNS regeneration in fish. As potential therapeutics of CNS injuries, it may be valuable to consider some of these factors [48]. After traumatic brain injury, transformation of quiescent type I radial glial cells into proliferative

type II radial glial cells similar to mammalian reactive astrocytes has been shown to be significantly suppressed in zebrafish harboring mutant glial maturation factor  $\beta$  [49].

A wide variety of fish species have been used for the study of regeneration, such as the lamprey, salmon, medaka, goldfish, and zebrafish (larva and adult), and have their pros and cons. Finally, we recommend one teleost fish, the Japanese eel (*Anguilla japonica*), known for its long spawning migration (about 3000 km) from Japan to the West Mariana Ridge (seamount chain) in the western North Pacific region [50]. They have also been used in the study of endocrinology because of their toughness to injury, and in behavioral experiments [51]. We developed an easy method for ablation of the telencephalon of the Japanese eel [52] and have observed their CNS regeneration and behavioral aspects of spatial learning and social behavior [53].

### 3. Integrins in Vertebrates (Mammals)

### 3.1. Integrin Signaling between Neurons and Astrocytes

Thy-1 is an abundantly expressed glycoprotein in neurons, which functions as an inhibitor of neurite outgrowth and acts as both a ligand, such as for integrins and other receptors, and as a receptor in Thy-1-expressing cells [54]. In the interaction between neuron and astrocytes, the  $\alpha\nu\beta3$  integrin, as a receptor, directly binds to the Thy-1 present on neurons, stimulating the astrocytes [55]. Furthermore,  $\alpha\nu\beta3$  integrin in astrocytes limits the neurites growth and also induces retraction of already existing processes by Thy-1 clustering induction [56]; it also triggers neurite retraction via the RhoA/ROCK pathway [57]. Moreover, upregulation of integrins in the astrocytes by inflammation induces neurite retraction by binding Thy-1, and formation of a ternary complex, i.e., Thy-1– $\alpha\nu\beta3$  integrin and Syndecan-4, alters astrocyte contractility and inhibits neurite outgrowth [58]. Astrocytic migration requires a large increase in the intracellular Ca<sup>2+</sup> concentration and connexin 43 hemichannel opening, which are induced by Thy-1 via  $\alpha\nu\beta3$  integrin in astrocytes [59,60]. Modulation of the astrocyte–neuron interaction in relation to this mechanism might contribute to axonal repair after CNS injury.

### 3.2. Role of Astroglial Integrins in the Functioning of the CNS

The largest integrin subfamily, the  $\beta$ 1 integrin, is highly expressed in the mammalian neural stem cells (NSCs), and plays a role in the "niche" by regulating the growth factor responsiveness of NSC into either remaining as stem cells or differentiating and migrating away to shape the developing cortex [61]. Moreover, activation of integrin-linked kinase (ILK) in cultured adult NSCs from  $\beta$ 1 integrin ablated mice reduced astrocytic differentiation, as  $\beta$ 1 integrin restricted astrocytic differentiation of adult hippocampal NSCs mediated by ILK [62].

Selective ablation of  $\alpha v$  or  $\beta 8$  integrin in astrocytes has revealed that  $\alpha v \beta 8$  integrin signaling links astrocytes to blood vessels and is essential for retinal angiogenesis [63]. In addition,  $\beta 1$  integrin is required for efficient blood vessel-guided migration of neuroblasts towards an infarct area and also their stable adhesion to laminin-rich astrocytes [64]. Thus, astroglial integrin signaling is indispensable for precise functioning of the CNS.

## 3.3. Integrins in Neurodegenerative Diseases

For neurons, integrin activation in neurite outgrowth and axonal regeneration have been extensively summarized [65]. Furthermore, integrin functions have, recently, been reviewed for effects of neuronal connectivity disruption in neurodevelopmental diseases, axonal outgrowth, synaptogenesis and synaptic maturation [17], and also for neuropsychiatric disorders [66].

Age-related macular degeneration (AMD) is characterized by a deficit in autophagy associated with deficiency of calcium and integrin binding protein 2 (CIB2), which are essential for visual functions [67]. Defects in the CIB2 protein have been reported to be related to deafness and/or vision deficits in humans, zebrafish, and drosophila [68].

Multiple sclerosis (MS) is a chronic auto-neuroinflammatory disease of the CNS driven by peripheral immune cells invading the target tissue to cause demyelination and neuronal death [69]. To overcome the BBB, the T cell  $\alpha 4\beta 1$  integrin interacts with endothelial vascular cell adhesion molecule 1 (V-CAM1) [70], and an antibody against  $\alpha 4\beta 1$  integrin (Natalizumab) has been commonly used in the treatment of MS [71]. The mechanism of T cell infiltration into the brain has not yet been elucidated, but recently, Zondler et al. reported that melanoma cell adhesion molecule (MCAM) signaling via phosphorylated PLC $\gamma 1$  induced intracellular signaling leading to  $\beta 1$  integrin activation on human memory T cells, resulting in increased brain T cell infiltration [72].

### 3.4. Integrins in the Regeneration of the CNS

In adult mammals, CNS axons fail to regenerate, while PNS neurons support robust axonal regeneration over long distances. Petrova and Eva reported selective axonal transport of integrins as regeneration-associated receptors [13]. A key difference in the regeneration capability between the CNS and PNS is whether integrin is transport into the axons or not. Integrins can mediate PNS regeneration, but are excluded from CNS axons after development [73,74]. Thus, when integrins and carrier Rab11 endosomes in the CNS axons were manipulated in vitro, axonal regeneration was observed in the dorsal root ganglion (DRG), although the findings remain to be seen in vivo [75,76].

Expression of the integrin activator kindlin-1 in the DRG neurons only promoted limited regeneration in the spinal cord, whereas co-expression of  $\alpha$ 9 integrin with kindlin-1 induced extensive regeneration and functional recovery [77,78]. This latter strategy, however, failed in the CNS axons, because of the exclusion of integrins from the CNS axons due to changes in trafficking with the maturation of neurons [73].

The roles of integrins related to glycoproteins, such as laminin-, fibronectin-, and collagen-associated integrins, in axonal regeneration in PNS injury have been intensively examined, but no effective results for CNS injury have been obtained yet. A possible reason is that most axon guidance repulsive molecules embark inactivation of integrins [15]. The authors claim that modulation of integrins could be a general approach to promote axonal regeneration in the CNS. Collectively, in this review, we are going to focus on the integrin functions in relation to astrocytes and reactive astrocytes rather than neurons.

# 4. Integrin Involvement in the Roles of Astrocytes in Physiological and Pathological States

### 4.1. Integrins Involved in Astrocyte Functions

Neurons are generated from NSCs in the subventricular zone (SVZ) of the lateral ventricle in the adult mammalian brain [79,80]. In the SVZ, slow-dividing cells generate fast-dividing cells that migrate tangentially within astrocytic tubes along the rostral migratory stream (RMS) to the olfactory bulb.  $\beta$ 1 integrins are required for the maintenance of these astrocytic tubes, and defects lead to ectopic migration of neuroblasts into the tissues around the RMS [81]. Thus,  $\beta$ 1 integrin promotes cell–cell interactions that link neuroblasts to astrocytic chains.

*Clostridium botulinum* C3 transferase (C3bot) peptides enhance neuronal regenerative growth and connectivity, and seemed to show potential as treatment for spinal cord injury (SCI) [82,83]. C3bot harboring a mutated integrin-binding RGD (arginine-glycine-aspartate) motif (C3bot-G89I) showed reduced binding to even astrocytes releasing the intermediate filament protein vimentin, suggesting that vimentin and  $\beta$ 1 integrin are binding partners and that their binding is involved in the internalization of C3bot into cultured astrocytes and astrocytes in SCI rats [84]. Thus, this mechanism of  $\beta$ 1 integrin signaling initiated by C3bot with release of vimentin from astrocytes could be used as a therapeutic target for SCI treatment.

Prominent labeling for  $\beta$ 1 integrin was observed in astrocyte processes near both unmyelinated nerve and myelinated nerve fibers in the normal optic nerve, while in a mouse model of glaucoma, in the presence of elevated intraocular pressure (IOP), apop-

totic astrocytes were observed [85]. Thus, astrocytes normally exhibit junctions with the basement membrane, which are disrupted by IOP elevation. Membrane-bound  $\beta$ 1 integrin is involved in mechano-sensitive signaling and is thought to be potentially beneficial or detrimental to axonal health.

Galectins, which are a family of  $\beta$ -galactoside-binding lectins consisting of 15 members in mammals, and their ligand glycoconjugates, including integrins, play diverse roles in both the physiology and pathology of the CNS [86]. In the adult mouse brain, galectin-1 binding activity has been detected on the  $\beta$ 1 integrin-expressing cells in the subependymal zone (SEZ). Furthermore,  $\beta$ 1 integrin has been shown to be required for galectin-1 functioning in neural progenitor cell (NPC) adhesion in vitro [87]. A genome-wide expression analysis, conducted to compare adult NSCs isolated from the SEZ and reactive astrocytes isolated 5 days after stab wound injury from the adult mouse cerebral cortex, revealed higher expression levels of galectin-1 and -3 levels in both types of cells, and addition of galectin-3 fully rescued the NSC phenotype in mutant animals [88,89]. Taken together, galectin-1 and -3 may contribute to providing a stem cell niche and play differential roles in neural generation via  $\beta$ 1 integrin.

### 4.2. Integrin Signaling in Reactive Astrocytes

Astrocytes become reactive after brain/spinal cord injury or neuroinflammation involving the CNS, and reactive astrocytes show morphological changes and upregulation of some intermediate filament proteins such as GFAP, vimentin, and nestin, as well as of ECM molecules such as TN-C and chondroitin sulfate proteoglycans (CSPGs) [90-93]. It remains under debate as to whether this is a favorable reaction as an essential state for wound healing (good thing and beneficial), or an unfavorable reaction inhibiting axonal regrowth and neurite outgrowth (bad thing and detrimental) [94,95]. On the one hand, corroborating the notion that it is an unfavorable reaction, activated astrocytes form a glial scar which acts as a physical barrier to regenerating axons in the mammalian brain, and neurotoxic reactive astrocytes induce cell death in the CNS [96,97]. On the other hand, as evidence of its being a favorable reaction, a glial scar prevents the spread of neuronal death and inflammation around the wound site [98,99]. Moreover, Zamanian et al. termed A1 reactive astrocytes, which showed strong upregulation of various genes, as destructive to neurons (neurotoxic) and A2 reactive astrocytes, which release many neuroprotective neurotrophic factors [100]. A2 astrocytes show upregulated expressions of HMGB1 and  $\beta$ 2 integrin, and promote adhesive interactions between the brain endothelium and endothelial progenitor cells [101,102].

Next, we focused on integrin signaling in reactive astrocytes to determine whether reactive astrocytes represent a favorable or unfavorable reaction to injury. Brain damage (i.e., traumatic brain injury, inflammation, or ischemia) is associated with upregulation of one of the astrocyte surface proteins, i.e., integrins, and proinflammatory cytokines and ECM proteins are released to generate a stiffer matrix, which leads to the formation of a glial scar. It is known that interaction between upregulated type I collagen and its receptor β1 integrin and N-cadherin-mediated cell adhesion play major functions around the lesion site of a SCI [103]. Furthermore, administration of anti- $\beta$ 1 integrin antibody during the scar formation period has been found to inhibit the interaction between type 1 collagen and Ras, and prevented transformation of reactive astrocytes to scar-forming astrocytes, thereby, preventing scar formation, promoting axonal regrowth, and improving functional recovery in the injured spinal cord [104]. They showed that normal migration of reactive astrocytes to limit the compact lesion area induced functional recovery, whereas their impaired migration resulted in a wide lesion area and poor functional recovery after the SCI. During the scar-forming phase, type I collagen is highly expressed in the injured spinal cord and induces astrocytic scar formation via the integrin-N-cadherin pathway [105].

Astrocyte adhesion and contractility are induced by  $\alpha\nu\beta3$  integrin/Thy-1 signaling, and are enhanced and accelerated by mechanical stress, as demonstrated using a magnetic bead assay [106]. The mechanoreceptor  $\alpha\nu\beta3$  integrin could be a new target for the negative

effects of reactive astrocytes in an injured brain. In accordance with these data, astrocytes are responsive as mechanosensitive cells via the mediation of integrins.

In an injured brain, expression of the actin-binding protein drebrin (DBN) is upregulated in reactive astrocytes, and is essential for glial scar formation and maintenance in vivo. DBN-mediated membrane trafficking via  $\beta$ 1 integrin, in inducing reactivity of astrocytes, has been shown to be an important neuroprotective mechanism following traumatic brain injury in mice [107]. Type I collagen is highly expressed in the spinal cord during astrocytic scar formation phase via the integrin-N-cadherin pathway, and its blockade prevents glial scar formation, resulting in improved axonal regrowth [105]. Therefore,  $\beta$ 1 integrin could be effectively targeted to promote SCI recovery.

It has been reported that dentin sialophosphoprotein (DSPP), a member of the small integrin-binding ligand N-linked glycoprotein family, interacts with  $\beta6$  integrin to promote cell proliferation, differentiation, and migration [108]. In fact, overexpression of DSPP increased activation of the Akt/mTOR pathway in astrocytes, which might promote proliferation and migration of reactive astrocytes [109]. Thus, targeting DSPP appears to be an effective therapeutic strategy for treating CNS injury related to glial scar formation. As Adams et al. advocate [94], we propose to target the coordination of reactive astrocytes and integrin signaling for developing therapies for neurodegenerative diseases (Figure 2).

#### 4.3. Integrins in Neuroinflammation Mediated by Astrocytes

Astrocytes interact not only with neurons, but also with microglial cells, oligodendrocytes, endothelial cells, and peripheral immune cells in the event of neuroinflammation [110,111]. Yoshizaki et al. revealed that after chronic spinal cord injury in mice, microglial inflammation was enhanced by reactive astrocytes via the fibronectin/ $\beta$ 1 integrin pathway. They showed that intralesional administration of anti- $\beta$ 1 integrin antibody in the subacute phase of SCI suppressed the production and secretion of tumor necrosis factor  $\alpha$ , an inflammatory cytokine, and glial scar formation by switching of the microglial phenotype to an anti-inflammatory phenotype [112]. Targeting  $\beta$ 1 integrin signaling as a potential therapy for SCI could have a potential risk due to the fact that it could form a lot of potential heterodimers, and that might induce integrin crosstalk.

It has been reported that anti- $\beta$ 1 integrin antibody exerts therapeutic effect by preventing glial scar formation [105]. Furthermore, the antibody also effectively blocks the interactions between reactive astrocytes and microglial cells via fibronectin. From these data, modulation of integrin signaling could be attempted for developing treatments against neuroimmunological diseases.

Another inhibitor, ATN-161, of  $\alpha 5\beta 1$  integrin, has been reported to reduce the infarct volume, edema, and infiltration of immune cells into the brain and functional deficit, and to be neuroprotective in MCAO mice, a mouse model of ischemic stroke [113].

In diabetic retinopathy, angiopoietin (Ang2) induces astrocyte apoptosis in the presence of high glucose via the  $\alpha v\beta 5$  integrin/GSK3 $\beta$ / $\beta$ -catenin pathway [114]. Thus, Ang2/integrin signaling could be targeted for inhibiting vascular leakage by astrocyte loss in early diabetic retinopathy.

Astrocytes are also known to be involved in the pathogenesis of Alzheimer's disease (AD). Functional changes, such as disrupt gliotransmission, neurotransmitter uptake, and altered calcium signaling in astrocytes have been observed in the brains of individuals with AD in the presence of amyloid beta. Actually, downregulation of astrocytic cytoskeletal proteins, such as actin  $\beta$ , dynein, and  $\alpha$  integrin, is observed in human AD patients [115]. Astrocyte  $\alpha v / \beta 1$  integrin receptor binding to recombinant human Tau, which mediates the entry of Tau fibrils into astrocytes, and induction of integrin signaling leads NF $\kappa$ B activation, causing upregulation of proinflammatory cytokines and chemokines, induction of a subgroup of neurotoxic astrocyte markers, and release of neurotoxic factors in primary cultures of astrocytes [116]. Since ablation of reactive astrocytes has been shown to exacerbate memory deficits in a mouse model of AD [117], manipulation of astrocytes could be useful for clarifying the mechanisms of memory loss in such conditions as tauopathies.



**Figure 2.** Involvement of integrins in CNS regeneration in collaboration with potential factors. Modulation of integrins with potential factor(s) stated in the main text, and possibly also the environment, could convert astrocytes or reactive astrocytes to healthy neurons for induction of neuro-regeneration in the mammalian CNS. The number(s) in square brackets beside the factor names are the reference numbers cited in the main text [28–31,106–109,112].

Glial scar, formed by activated astrocytes, have been thought to interfere with CNS regeneration. Anderson et al. showed that preventing astrocyte scar formation significantly reduced axonal regrowth, and their RNA sequencing revealed that astrocytes at the site of SCI expressed multiple axon-growth supporting molecules [118]. Furthermore, the transcriptome profile has been reported of neuroinflammatory astrocyte subtypes induced by bacterial cell wall endotoxin lipopolysaccharide (LPS) injection in mice [119]. Since astrocytes and reactive astrocytes have been implicated in traumatic SCI [120], targeting of glial scar formation after a SCI might be helpful to induce CNS regeneration.

### 4.4. TN-C Function in Astrocytes via Integrins

The authors have focused on TN-C and osteopontin (OPN), which are ligands for the integrin family of receptors in the brain, to investigate the neuroinflammatory functions of astrocytes. For example, TN-C has functional roles such as the BBB recovery from breakdown caused by a brain injury, proliferation of astrocytes, and balancing between neuroprotective-and neurotoxic-cytokine production, which were examined after a stab wound injury to mice cerebral cortex and in primary culture of astrocytes [26–28]. First, we summarize the roles that TN-C plays in integrin signaling involved in the functioning of astrocytes.

TN-C is one of the ECM glycoproteins of the CNS that is involved in CNS development, and its expression is upregulated after brain or spinal cord injury [121,122]. TN-C is also known to have immunomodulatory roles in neurodegenerative disorders, and to be highly associated with astrocyte activation and glial scar formation [123].

Neurons transfected with  $\alpha$ 9 integrin and kindlin-1 in vitro grew axons for long distances on TN-C, however, a much lower degree of regeneration was observed in vivo [124]. They hypothesized that major inhibitory molecules, not present in the in vitro culture system, in the adult CNS might inactivate integrins, preventing them from binding with TN-C. Actually, Tan et al. revealed using adeno-associated virus vector expressing  $\alpha 9$  integrin in the adult rat DRG neurons, a number of axons growing into the dorsal root entry zone in vivo. However, the inhibitory substrates CSPG and Nogo blocked the growth by inactivating integrins [77]. Since expression of the  $\alpha 9$  integrin subunit was downregulated in the mature corticospinal tract (CST), grafting of lentivirus-mediated  $\alpha 9$  integrin-overexpressing induced pluripotent stem cell (iPSC)-derived human neural progenitor cells (hNPCs) resulted in increased neurite outgrowth in the presence of TN-C in the developing CST [125]. Using a scratch wound in the primary culture of astrocytes, it was shown that extracellular vesicle functions and wound recovery are mediated by  $\beta 1$  integrin related to IL-1 $\beta$  treated TN-C-bound-primary culture [126]. Although TN-C exerts an inhibitory effect on mature axons, when specifically bound to  $\alpha 9\beta 1$  integrin, TN-C can promote neurite outgrowth and axonal regeneration [127].

Recently, we reported TN-C as an inhibitor of neuronal death after stab wound injury in mouse brain and in a primary culture with LPS in the presence of 2-carba-cyclic phosphatidic acid (2ccPA) [31]. In this study, TN-C secreted from astrocytes had a neuroprotective role, but only in the presence of 2ccPA. Thus, although TN-C and integrins are becoming well-known as being neuroprotective, additional factor(s) and conditions seem to be necessary for their inducing neuronal regeneration in vivo.

### 4.5. OPN Functions in Astrocytes Mediated via Integrins

Next, we summarize the functions of OPN mediated via integrins playing a role in astrocyte functions.

OPN is a proinflammatory cytokine inducer and a pleiotropic protein involved in a wide range of cell functions, such as bone development and remodeling, vascularization, wound healing, immune response, and inflammation, and also tumor metastasis [128,129]. OPN is weakly expressed in the brain under normal conditions, while it is strongly expressed in the microglia and astrocytes, exerting a neuroprotective effect after brain injury [130].

Recently, we demonstrated that in the injured cerebral cortex after a stab wound and in a primary culture of astrocytes with LPS, an inflammation inducer, reactive astrocytes also expressed OPN, and that OPN exerted its neuroprotective role both in vivo and in vitro via the mediation of  $\alpha 9\beta 1$  integrin [29,30]. Although OPN utilizes various combinations of integrin heterodimers, binding through the RGD domain, the precise receptors it binds to depends on the pathological conditions. Actually, several cell surface integrins, such as  $\alpha \nu \beta 1$ ,  $\alpha \nu \beta 3$ ,  $\alpha \nu \beta 5$ ,  $\alpha 4\beta 1$ ,  $\alpha 8\beta 3$ , and  $\alpha 9\beta 1$ , and also CD44 variants are known to serve as receptors for OPN [131,132].

OPN has been reported as a potential cerebrospinal fluid biomarker in amyotrophic lateral sclerosis [133,134] and also in neuromyelitis optica (NMO), an inflammatory disease of the CNS affecting the optic nerve and spinal cord [135]. It has been shown that OPN in the cerebrospinal fluid in NMO patients bound with integrin  $\alpha\nu\beta$ 3 to promote macrophage chemotaxis by activating the phosphoinositide 3-kinase and MEK1/2 signaling pathways.

The RGD domain of OPN interacts with the integrins  $\alpha\nu\beta3$  and  $\beta1$ , and seems to play a role in the neuroprotective actions of OPN in models of Parkinson's disease. Broom et al. reported that both OPN and the synthetic 15-mer RGD domain of OPN were equally protective of increased glial-derived neurotrophic factor and brain-derived neurotrophic factor levels against 1-methyl-4-phenylpyridinium exposure of ventral mesencephalic cultures, accompanied by a decrease in the numbers of activated microglia, but no change in the astrocyte number [136]. This finding suggests that other domains of OPN may also be needed for their neuroprotective role.

Human immunodeficiency virus does not enter or replicate in neurons, but can infect microglia and astrocytes to induce neurotoxicity and neuroinflammation. Zhu et al. showed that extracellular OPN blocked the negative effects of the envelope on neurite growth via the integrin mTORC1/2 signaling pathway, which is involved in long-term potentiation

and long-term depression [137], learning and memory, neuronal survival, differentiation, and morphogenesis [138].

OPN signaling through  $\beta 1$  integrin is not only required for stimulation and maintenance of neurite growth in cultured primary rat cortical neurons, but also to protect neurons from excitotoxicity [139]. Thus,  $\beta 1$  integrin is the key factor involved in higher brain functions mediated by OPN.

### 4.6. Integrins Functions in the BBB

The BBB, a picket line, constitutes a neurovascular unit formed by microvascular endothelial cells (EC), pericytes, and astrocytes that prevent certain molecules from passing from the blood into the brain. Astrocyte endfeet and pericytes that contact the cerebrovascular endothelium, and specialized connections with the tight junction (TJ), i.e., claudin-5, between the EC, and zonula occludens 1 (ZO-1) and occludin are involved in the integrity of the ECM of the basal lamina [140,141].

Integrins contribute significantly to angiogenesis and vascular remodeling mediated by thrombospondin-1/integrin/YAP signaling [142,143]. The role of integrins and ECM functions in the endothelial cells of the BBB in stroke or EAE have already been reported in previous papers [144–146]. In the present paper, we mainly focus on pericyte and astrocyte functions in the BBB mediated via integrin signaling.

Loss of the tight connections of astrocytes to endothelial cells and reduced expressions of pericyte markers and of various integrins such as  $\alpha$ 6,  $\alpha$ 3 $\beta$ 1, and  $\beta$ 1 with loss of TJ integrity have been associated with enhanced permeability of the BBB in mice [147]. Nakamura et al. showed, in an ischemic stroke mouse model, that perlecan, a major heparan sulfate proteoglycan of the basement membrane, regulated the pericyte dynamics through  $\alpha$ 5 $\beta$ 1 integrin in the maintenance and repair of the BBB [148]. These data suggest that  $\beta$ 1 integrin is involved in the integrity of the BBB.

ECM epidermal growth factor-like protein 7 (EGFL7) secreted by the ECM is increased in patients with MS and in EAE mouse model of MS. T lymphocytes expressing  $\alpha\nu\beta3$ integrin adhered to EGFL7. Recombinant EGFL7 has been shown to improve EAE, tighten the BBB, and reduce neuroinflammation in mouse [149]. Thus, EGFL7/ $\alpha\nu\beta3$  integrin signaling could serve as a new therapeutic target for MS.

Chronic mild hypoxia (CMH) promoted a strong vascular remodeling response in the spinal cord blood vessels via an  $\alpha 5\beta 1$  integrin-mediated mechanism in mice [150]. CMH triggered astrocyte activation specifically in the gray matter and the newly generated vessels grew towards neuron-rich areas and were also closely associated with astrocytic endfeet.

### 5. Conclusions

Integrin signaling mediates a variety of functions, not only in neurons but also in astrocytes in the brains of mammals, fish, as well as invertebrates. In this review, we have summarized the roles of integrin signaling in the functioning of astrocytes and reactive astrocytes, which could provide further insight for unlocking the potential for valuable neuronal regeneration in CNS injury and disease in humans.

**Author Contributions:** H.I.-K., writing original draft, writing review and editing, supervision and conceptualization; C.S., writing original draft, writing review and editing, performed molecular phylogenetic tree analysis, drawing schema (Figure 2); T.T., writing original draft, writing review and editing, performed molecular phylogenetic tree analysis. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by Keio Gijuku Academic Development Funds to T.T. and H.I.-K.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

### Abbreviations

BBB	blood brain barrier
CNS	central nervous system
CSPG	chondroitin sulfate proteoglycan
DRG	dorsal root ganglion
ECM	extracellular cellular matrix
GFAP	glial fibrillary acidic protein
LPS	lipopolysaccharide
NSCs	neural stem cells
OPN	osteopontin
PNS	peripheral nervous system
RGD	Arg-Gly-Asp
SCI	spinal cord injury
SVZ	subventricular zone
TJ	tight junction
TN-C	tenascin C

### References

- 1. Harrison, R.G. Observations on the livining developing nerve fiber. Proc. Soc. Exp. Biol. Med. 1907, 4, 140–143. [CrossRef]
- 2. Hynes, R.O.; Pearistein, E.S. Investigations of the possible role of proteases in altering surface proteins of virally transformed hamster fibroblasts. *J. Supramol. Struct.* **1976**, *4*, 1–14. [CrossRef] [PubMed]
- 3. Schwarzbauer, J.E.; Paul, J.I.; Hynes, R.O. On the origin of species of fibronectin (RNA splicing/protein variants/fusion pr oteins). *Proc. Natl. Acad. Sci. USA* **1985**, *82*, 1424–1428. [CrossRef] [PubMed]
- 4. Hynes, R.O.; Destree, A.T. Relationships between fibronectin (LETS protein) and actin. Cell 1978, 15, 875–886. [CrossRef]
- 5. Horwitz, A.; Duggan, K.; Buck, C.; Beckerle, M.C.; Burridge, K. Interaction of plasma membrane fibronectin receptor with talin—A transmembrane linkage. *Nature* **1986**, *320*, 531–533. [CrossRef]
- 6. Horwitz, A.R. The Origins of the Molecular Era of Adhesion Research. Nat. Rev. Mol. Cell Biol. 2012, 13, 805811. [CrossRef]
- Tamkun, J.W.; DeSimone, D.W.; Fonda, D.; Patel, R.S.; Buck, C.; Horwitz, A.F.; Hynes, R.O. Structure of integrin, a glycoprotein involved in the transmembrane linkage between fibronectin and actin. *Cell* 1986, 46, 271–282. [CrossRef]
- Spinardi, L.; Ren, Y.L.; Sanders, R.; Giancotti, F.G. The β4 Subunit Cytoplasmic Domain Mediates the Interaction of α6β4 Integrin with the Cytoskeleton of Hemidesmosomes. *Mol. Biol. Cell* **1993**, *4*, 871–884. [CrossRef]
- Mezu-Ndubuisi, O.J.; Maheshwari, A. The Role of Integrins in Inflammation and Angiogenesis. *Pediatr. Res.* 2021, 89, 1619–1626. [CrossRef]
- Chen, Z.; Shen, G.; Tan, X.; Qu, L.; Zhang, C.; Ma, L.; Luo, P.; Cao, X.; Yang, F.; Liu, Y.; et al. ID1/ID3 Mediate the Contribution of Skin Fibroblasts to Local Nerve Regeneration through Itgα6 in Wound Repair. *Stem Cells Transl. Med.* 2021, 10, 1637–1649. [CrossRef]
- 11. Kawano, H.; Kimura-Kuroda, J.; Komuta, Y.; Yoshioka, N.; Li, H.P.; Kawamura, K.; Li, Y.; Raisman, G. Role of the Lesion Scar in the Response to Damage and Repair of the Central Nervous System. *Cell Tissue Res.* **2012**, *349*, 169–180. [CrossRef]
- 12. Eva, R.; Fawcett, J. Integrin Signalling and Traffic during Axon Growth and Regeneration. *Curr. Opin. Neurobiol.* **2014**, 27, 179–185. [CrossRef]
- 13. Petrova, V.; Eva, R. The Virtuous Cycle of Axon Growth: Axonal Transport of Growth-Promoting Machinery as an Intrinsic Determinant of Axon Regeneration. *Dev. Neurobiol.* **2018**, *78*, 898–925.
- 14. Fawcett, J.W.; Verhaagen, J. Intrinsic Determinants of Axon Regeneration. Dev. Neurobiol. 2018, 78, 890–897. [CrossRef]
- 15. Nieuwenhuis, B.; Haenzi, B.; Andrews, M.R.; Verhaagen, J.; Fawcett, J.W. Integrins Promote Axonal Regeneration after Injury of the Nervous System. *Biol. Rev.* 2018, *93*, 1339–1362. [CrossRef]
- 16. Park, Y.K.; Goda, Y. Integrins in synapse regulation. Nat. Rev. Neurosci. 2016, 17, 745–765. [CrossRef]
- 17. Lilja, J.; Ivaska, J. Integrin Activity in Neuronal Connectivity. J. Cell Sci. 2018, 131, jcs212803. [CrossRef]
- De Arcangelis, A.; Georges-Labouesse, E. Integrin and ECM Functions in Vertebrate Development. *Trends Genet.* 2000, 16, 389–395. [CrossRef]
- 19. Allen, N.J.; Eroglu, C. Cell Biology of Astrocyte-Synapse Interactions. Neuron 2017, 96, 697–708. [CrossRef]
- 20. Verkhratsky, A.; Nedergaard, M. Physiology of Astroglia. Physiol. Rev. 2018, 98, 239–389. [CrossRef]
- 21. Khakh, B.S.; Deneen, B. The Emerging Nature of Astrocyte Diversity. Annu. Rev. Neurosci. 2019, 42, 187–207. [CrossRef] [PubMed]
- 22. Morita, M.; Ikeshima-Kataoka, H.; Kreft, M.; Vardjan, N.; Zorec, R.; Noda, M. Metabolic Plasticity of Astrocytes and Aging of the Brain. *Int. J. Mol. Sci.* 2019, 20, 941. [CrossRef] [PubMed]
- Zhang, Z.; Ma, Z.; Zou, W.; Guo, H.; Liu, M.; Ma, Y.; Zhang, L. The Appropriate Marker for Astrocytes: Comparing the Distribution and Expression of Three Astrocytic Markers in Different Mouse Cerebral Regions. *BioMed Res. Int.* 2019, 2019, 9605265. [CrossRef] [PubMed]
- Johnson, M.S.; Lu, N.; Denessiouk, K.; Heino, J.; Gullberg, D. Integrins during Evolution: Evolutionary Trees and Model Organisms. *Biochim. Biophys. Acta Biomembr.* 2009, 1788, 779–789. [CrossRef]

- 25. Sebé-Pedrósa, A.; Rogerb, A.J.; Langc, F.B.; Kingd, N.; Ruiz-Trillo, I. Ancient origin of the integrin-mediated adhesion and signaling machinery. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 10142–10147. [CrossRef]
- Ikeshima-Kataoka, H.; Saito, S.; Yuasa, S. Tenascin-C Is Required for Proliferation of Astrocytes in Primary Culture. *In Vivo* 2007, 21, 629–634.
- Ikeshima-Kataoka, H.; Yuasa, S. Selective Ablation of an Astroglial Subset by Toxic Gene Expression Driven by Tenascin Promoter. *Neurol. Res.* 2008, 30, 701–709. [CrossRef]
- 28. Ikeshima-Kataoka, H.; Shen, J.-S.; Eto, Y.; Saito, S.; Yuasa, S. Alteration of Inflammatory Cytokine Production in the Injured Central Nervous System of Tenascin-Deficient Mice. *In Vivo* **2008**, *22*, 409–414.
- 29. Ikeshima-Kataoka, H.; Abe, Y.; Abe, T.; Yasui, M. Immunological Function of Aquaporin-4 in Stab-Wounded Mouse Brain in Concert with a pro-Inflammatory Cytokine Inducer, Osteopontin. *Mol. Cell. Neurosci.* **2013**, *56*, 65–75. [CrossRef]
- Ikeshima-Kataoka, H.; Matsui, Y.; Uede, T. Osteopontin Is Indispensable for Activation of Astrocytes in Injured Mouse Brain and Primary Culture. *Neurol. Res.* 2018, 40, 1071–1079. [CrossRef]
- Nakashima, M.; Gotoh, M.; Hashimoto, K.; Endo, M.; Murakami-Murofushi, K.; Ikeshima-Kataoka, H.; Miyamoto, Y. The Neuroprotective Function of 2-Carba-Cyclic Phosphatidic Acid: Implications for Tenascin-C via Astrocytes in Traumatic Brain Injury. J. Neuroimmunol. 2021, 361, 577749. [CrossRef]
- Xu, S.; Zhang, T.; Cao, Z.; Zhong, W.; Zhang, C.; Li, H.; Song, J. Integrin-α9β1 as a Novel Therapeutic Target for Refractory Diseases: Recent Progress and Insights. *Front. Immunol.* 2021, *12*, 638400. [CrossRef]
- Tamura, K.; Stecher, G.; Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Mol. Biol. Evol. 2021, 38, 3022–3027. [CrossRef]
- Bonar, N.A.; Petersen, C.P. Integrin Suppresses Neurogenesis and Regulates Brain Tissue Assembly in Planarian Regeneration. Development 2017, 144, 784–794. [CrossRef]
- 35. Hisamoto, N.; Shimizu, T.; Asai, K.; Sakai, Y.; Pastuhov, S.I.; Hanafusa, H.; Matsumoto, K.C. Elegans Tensin Promotes Axon Regeneration by Linking the Met-like SVH-2 and Integrin Signaling Pathways. *J. Neurosci.* **2019**, *39*, 5662–5672. [CrossRef]
- Pastuhov, S.I.; Fujiki, K.; Tsuge, A.; Asai, K.; Ishikawa, S.; Hirose, K.; Matsumoto, K.; Hisamoto, N. The Core Molecular Machinery Used for Engulfment of Apoptotic Cells Regulates the JNK Pathway Mediating Axon Regeneration in Caenorhabditis Elegans. J. Neurosci. 2016, 36, 9710–9721. [CrossRef]
- Sakai, Y.; Tsunekawa, M.; Ohta, K.; Shimizu, T.; Pastuhov, S.I.; Hanafusa, H.; Hisamoto, N.; Matsumoto, K. The Integrin Signaling Network Promotes Axon Regeneration via the Src–Ephexin–RhoA GTPase Signaling Axis. J. Neurosci. 2021, 41, 4754–4767. [CrossRef]
- Sammut, M.; Cook, S.J.; Nguyen, K.C.Q.; Felton, T.; Hall, D.H.; Emmons, S.W.; Poole, R.J.; Barrios, A. Glia-Derived Neurons Are Required for Sex-Specific Learning in *C. Elegans. Nature* 2015, 526, 385–390. [CrossRef]
- De Vault, L.; Li, T.; Izabel, S.; Thompson-Peer, K.L.; Jan, L.Y.; Jan, Y.N. Dendrite Regeneration of Adult Drosophila Sensory Neurons Diminishes with Aging and Is Inhibited by Epidermal-Derived Matrix Metalloproteinase 2. *Genes Dev.* 2018, 32, 402–414. [CrossRef]
- Davison, A.; Neiman, M. Mobilizing Molluscan Models and Genomes in Biology. *Philos. Trans. R. Soc. B Biol. Sci.* 2021, 376, 20200163. [CrossRef]
- 41. Imperadore, P.; Fiorito, G. Cephalopod Tissue Regeneration: Consolidating over a Century of Knowledge. *Front. Physiol.* **2018**, *9*, 593. [CrossRef]
- 42. Imperadore, P.; Parazzoli, D.; Oldani, A.; Duebbert, M.; Büschges, A.; Fiorito, G. From Injury to Full Repair: Nerve Regeneration and Functional Recovery in the Common Octopus, Octopus Vulgaris. *J. Exp. Biol.* **2019**, 222, jeb209965. [CrossRef]
- Matsukawa, T.; Arai, K.; Koriyama, Y.; Liu, Z.; Kato, S. Axonal Regeneration of Fish Optic Nerve after Injury. *Biol. Pharm. Bull.* 2004, 27, 445–451. [CrossRef]
- 44. Stuermer, C.A.; Bastmeyer, M.; Bähr, M.; Strobel, G.; Paschke, K. Trying to Understand Axonal Regeneration in the CNS of Fish. *J. Neurobiol.* **1992**, *23*, 537–550. [CrossRef]
- 45. Ju, Y.; Janmey, P.A.; Mccormick, M.E.; Sawyer, E.S.; Flanagan, L.A. Enhanced Neurite Growth from Mammalian Neurons in Three-Dimensional Salmon Fibrin Gels. *Biomaterials* **2007**, *28*, 2097–2108. [CrossRef]
- Dowding, A.J.; Maggs, A.; Scholes, J. Diversity amongst the Microglia in Growing and Regenerating Fish CNS: Immunohistochemical Characterization Using FL. 1, an Anti-macrophage Monoclonal Antibody. *Glia* 1991, 4, 345–364. [CrossRef]
- 47. Var, S.R.; Byrd-Jacobs, C.A. Role of Macrophages and Microglia in Zebrafish Regeneration. *Int. J. Mol. Sci.* 2020, 21, 4768. [CrossRef]
- Rasmussen, J.P.; Sagasti, A. Learning to Swim, Again: Axon Regeneration in Fish. *Exp. Neurol.* 2017, 287, 318–330. [CrossRef] [PubMed]
- 49. Yin, G.; Du, M.; Li, R.; Li, K.; Huang, X.; Duan, D.; Ai, X.; Yao, F.; Zhang, L.; Hu, Z.; et al. Glia Maturation Factor Beta Is Required for Reactive Gliosis after Traumatic Brain Injury in Zebrafish. *Exp. Neurol.* **2018**, *305*, 129–138. [CrossRef] [PubMed]
- 50. Inoue, J.G.; Miya, M.; Miller, J.; Sado, T.; Hanel, R. Deep-Ocean Origin of the Freshwater Eels. *Biol. Lett.* 2010, *6*, 363–366. [CrossRef] [PubMed]
- 51. Watanabe, S.; Shinozuka, K. Spatial Learning in Japanese Eels (Anguilla japonica). Anim. Cogn. 2020, 23, 233–236. [CrossRef]
- 52. Tsubokawa, T.; Tsukada, T. Experimental Study: Ablation in the Telencephalon of Japanese Eel (*Anguilla japonica*). *Mannou News* **2020**, *11*, 1–2. (In Japanese)

- Tsubokawa, T.; Saito, K.; Kawano, H.; Kawamura, K.; Shinozuka, K.; Watanabe, S. Pharmacological Effects on Mirror Approaching Behavior and Neurochemical Aspects of the Telencephalon in the Fish, Medaka (*Oryzias latipes*). Soc. Neurosci. 2009, 4, 276–286. [CrossRef]
- 54. Morris, R.J. Thy-1, a Pathfinder Protein for the Post-Genomic Era. Front. Cell Dev. Biol. 2018, 6, 173. [CrossRef]
- 55. Hermosilla, T.; Muñoz, D.; Herrera-Molina, R.; Valdivia, A.; Muñoz, N.; Nham, S.U.; Schneider, P.; Burridge, K.; Quest, A.F.G.; Leyton, L. Direct Thy-1/αVβ3 Integrin Interaction Mediates Neuron to Astrocyte Communication. *Biochim. Biophys. Acta Mol. Cell Res.* 2008, 1783, 1111–1120. [CrossRef]
- 56. Herrera-Molina, R.; Frischknecht, R.; Maldonado, H.; Seidenbecher, C.I.; Gundelfinger, E.D.; Hetz, C.; de la Luz Aylwin, M.; Schneider, P.; Quest, A.F.G.; Leyton, L. Astrocytic αvβ3 Integrin Inhibits Neurite Outgrowth and Promotes Retraction of Neuronal Processes by Clustering Thy-1. *PLoS ONE* 2012, 7, e34295. [CrossRef]
- 57. Maldonado, H.; Calderon, C.; Burgos-Bravo, F.; Kobler, O.; Zuschratter, W.; Ramirez, O.; Härtel, S.; Schneider, P.; Quest, A.F.G.; Herrera-Molina, R.; et al. Astrocyte-to-Neuron Communication through Integrin-Engaged Thy-1/CBP/Csk/Src Complex Triggers Neurite Retraction via the RhoA/ROCK Pathway. *Biochim. Biophys. Acta Mol. Cell Res.* 2017, 1864, 243–254. [CrossRef]
- Burgos-Bravo, F.; Martínez-Meza, S.; Quest, A.F.G.; Wilson, C.A.M.; Leyton, L. Application of Force to a Syndecan-4 Containing Complex with Thy-1–AVβ3 Integrin Accelerates Neurite Retraction. *Front. Mol. Biosci.* 2020, 7, 582257. [CrossRef]
- Lagos-Cabré, R.; Brenet, M.; Díaz, J.; Pérez, R.D.; Pérez, L.A.; Herrera-Molina, R.; Quest, A.F.G.; Leyton, L. Intracellular Ca<sup>2+</sup> Increases and Connexin 43 Hemichannel Opening Are Necessary but Not Sufficient for Thy-1-Induced Astrocyte Migration. *Int.* J. Mol. Sci. 2018, 19, 2179. [CrossRef]
- 60. Lagos-Cabré, R.; Burgos-Bravo, F.; Avalos, A.M.; Leyton, L. Connexins in Astrocyte Migration. *Front. Pharmacol.* **2020**, *10*, 1546. [CrossRef]
- 61. Campos, L.S. B1 Integrins and Neural Stem Cells: Making Sense of the Extracellular Environment. *BioEssays* 2005, 27, 698–707. [CrossRef]
- 62. Brooker, S.M.; Bond, A.M.; Peng, C.Y.; Kessler, J.A. B1-Integrin Restricts Astrocytic Differentiation of Adult Hippocampal Neural Stem Cells. *Glia* 2016, 64, 1235–1251. [CrossRef]
- 63. Hirota, S.; Liu, Q.; Lee, H.S.; Hossain, M.G.; Lacy-Hulbert, A.; Mccarty, J.H. The Astrocyte-Expressed Integrin αvβ8 Governs Blood Vessel Sprouting in the Developing Retina. *Development* **2011**, *138*, 5157–5166. [CrossRef]
- Fujioka, T.; Kaneko, N.; Ajioka, I.; Nakaguchi, K.; Omata, T.; Ohba, H.; Fässler, R.; García-Verdugo, J.M.; Sekiguchi, K.; Matsukawa, N.; et al. B1 Integrin Signaling Promotes Neuronal Migration along Vascular Scaffolds in the Post-Stroke Brain. *EBioMedicine* 2017, 16, 195–203. [CrossRef]
- 65. Cheah, M.; Andrews, M.R. Integrin Activation: Implications for Axon Regeneration. Cells 2018, 7, 20. [CrossRef]
- Jaudon, F.; Thalhammer, A.; Cingolani, L.A. Integrin adhesion in brain assembly: From molecular structure to neuropsychiatric disorders. *Eur. J. Neurosci.* 2021, 53, 3831–3850. [CrossRef]
- Sethna, S.; Scott, P.A.; Giese, A.P.J.; Duncan, T.; Jian, X.; Riazuddin, S.; Randazzo, P.A.; Redmond, T.M.; Bernstein, S.L.; Riazuddin, S.; et al. CIB2 Regulates MTORC1 Signaling and Is Essential for Autophagy and Visual Function. *Nat. Commun.* 2021, 12, 3906. [CrossRef] [PubMed]
- Riazuddin, S.; Belyantseva, I.A.; Giese, A.P.J.; Lee, K.; Indzhykulian, A.A.; Nandamuri, S.P.; Yousaf, R.; Sinha, G.P.; Lee, S.; Terrell, D.; et al. Alterations of the CIB2 Calcium-and Integrin-Binding Protein Cause Usher Syndrome Type 1J and Nonsyndromic Deafness DFNB48. *Nat. Genet.* 2012, 44, 1265–1271. [CrossRef] [PubMed]
- 69. Reich, D.S.; Lucchinetti, C.F.; Calabresi, P.A. Multiple Sclerosis. N. Engl. J. Med. 2018, 378, 169–180. [CrossRef] [PubMed]
- 70. Glatigny, S.; Duhen, R.; Arbelaez, C.; Kumari, S.; Bettelli, E. Integrin Alpha L Controls the Homing of Regulatory T Cells during CNS Autoimmunity in the Absence of Integrin Alpha 4. *Sci. Rep.* **2015**, *5*, 7834. [CrossRef]
- Polman, C.; O'connor, P.; Havrdova, E.; Hutchinson, M.; Kappos, L.; Miller, D.; Phillips, J.; Lublin, F.; Giovannoni, G.; Wajgt, A.; et al. A Randomized, Placebo-Controlled Trial of Natalizumab for Relapsing Multiple Sclerosis. *N. Engl. J. Med.* 2006, 354, 899–910. [CrossRef]
- 72. Zondler, L.; Herich, S.; Kotte, P.; Körner, K.; Schneider-Hohendorf, T.; Wiendl, H.; Schwab, N.; Zarbock, A. MCAM/CD146 Signaling via PLCγ1 Leads to Activation of β1-Integrins in Memory T-Cells Resulting in Increased Brain Infiltration. *Front. Immunol.* 2020, *11*, 599936. [CrossRef]
- 73. Franssen, E.H.P.; Zhao, R.R.; Koseki, H.; Kanamarlapudi, V.; Hoogenraad, C.C.; Eva, R.; Fawcett, J.W. Exclusion of Integrins from CNS Axons Is Regulated by Arf6 Activation and the AIS. *J. Neurosci.* **2015**, *35*, 8359–8375. [CrossRef]
- Andrews, M.R.; Soleman, S.; Cheah, M.; Tumbarello, D.A.; Mason, M.R.J.; Moloney, E.; Verhaagen, J.; Bensadoun, J.C.; Schneider, B.; Aebischer, P.; et al. Axonal Localization of Integrins in the CNS Is Neuronal Type and Age Dependent. *eNeuro* 2016, 3, 5546–5557. [CrossRef]
- 75. Eva, R.; Koseki, H.; Kanamarlapudi, V.; Fawcett, J.W. EFA6 Regulates Selective Polarised Transport and Axon Regeneration from the Axon Initial Segment. *J. Cell Sci.* 2017, 130, 3663–3675. [CrossRef]
- 76. Koseki, H.; Donegá, M.; Lam, B.Y.; Petrova, V.; van Erp, S.; Yeo, G.S.; Kwok, J.C.; Ffrench-Constant, C.; Eva, R.; Fawcett, J.W. Selective Rab11 Transport and the Intrinsic Regenerative Ability of CNS Axons. *eLife* 2017, *6*, e26956. [CrossRef]
- 77. Tan, C.L.; Kwok, J.C.F.; Patani, R.; Ffrench-Constant, C.; Chandran, S.; Fawcett, J.W. Integrin Activation Promotes Axon Growth on Inhibitory Chondroitin Sulfate Proteoglycans by Enhancing Integrin Signaling. J. Neurosci. 2011, 31, 6289–6295. [CrossRef]

- Cheah, M.; Andrews, M.R.; Chew, D.J.; Moloney, E.B.; Verhaagen, J.; Fässler, R.; Fawcett, J.W. Expression of an Activated Integrin Promotes Long-Distance Sensory Axon Regeneration in the Spinal Cord. J. Neurosci. 2016, 36, 7283–7297. [CrossRef]
- 79. Obernier, K.; Alvarez-Buylla, A. Neural Stem Cells: Origin, Heterogeneity and Regulation in the Adult Mammalian Brain. *Development* 2019, 146, dev156059. [CrossRef]
- Bond, A.M.; Ming, G.; Song, H. Ontogeny of Adult Neural Stem Cells in the Mammalian Brain. In *Current Topics in Developmental Biology*; Academic Press Inc.: Cambridge, MA, USA, 2021; Volume 142, pp. 67–98. ISBN 9780128152812.
- Belvindrah, R.; Hankel, S.; Walker, J.; Patton, B.L.; Müller, U. B1 Integrins Control the Formation of Cell Chains in the Adult Rostral Migratory Stream. J. Neurosci. 2007, 27, 2704–2717. [CrossRef]
- Boato, F.; Hendrix, S.; Huelsenbeck, S.C.; Hofmann, F.; Große, G.; Djalali, S.; Klimaschewski, L.; Auer, M.; Just, I.; Ahnert-Hilger, G.; et al. C3 Peptide Enhances Recovery from Spinal Cord Injury by Improved Regenerative Growth of Descending Fiber Tracts. J. Cell Sci. 2010, 123, 1652–1662. [CrossRef] [PubMed]
- Loske, P.; Boato, F.; Hendrix, S.; Piepgras, J.; Just, I.; Ahnert-Hilger, G.; Höltje, M. Minimal Essential Length of Clostridium Botulinum C3 Peptides to Enhance Neuronal Regenerative Growth and Connectivity in a Non-Enzymatic Mode. *J. Neurochem.* 2012, 120, 1084–1096. [CrossRef] [PubMed]
- Adolf, A.; Rohrbeck, A.; Münster-Wandowski, A.; Johansson, M.; Kuhn, H.G.; Kopp, M.A.; Brommer, B.; Schwab, J.M.; Just, I.; Ahnert-Hilger, G.; et al. Release of Astroglial Vimentin by Extracellular Vesicles: Modulation of Binding and Internalization of C3 Transferase in Astrocytes and Neurons. *Glia* 2019, 67, 703–717. [CrossRef]
- Quillen, S.; Schaub, J.; Quigley, H.; Pease, M.; Korneva, A.; Kimball, E. Astrocyte Responses to Experimental Glaucoma in Mouse Optic Nerve Head. *PLoS ONE* 2020, 15, e0238104. [CrossRef]
- Nio-Kobayashi, J.; Itabashi, T. Galectins and Their Ligand Glycoconjugates in the Central Nervous System Under Physiological and Pathological Conditions. *Front. Neuroanat.* 2021, 15, 767330. [CrossRef] [PubMed]
- Sakaguchi, M.; Imaizumi, Y.; Shingo, T.; Tada, H.; Hayama, K.; Yamada, O.; Morishita, T.; Kadoya, T.; Uchiyama, N.; Shimazaki, T.; et al. Regulation of Adult Neural Progenitor Cells by Galectin-1/β1 Integrin Interaction. *J. Neurochem.* 2010, 113, 1516–1524. [CrossRef]
- Götz, M.; Sirko, S.; Beckers, J.; Irmler, M. Reactive Astrocytes as Neural Stem or Progenitor Cells: *In Vivo* Lineage, *In Vitro* Potential, and Genome-Wide Expression Analysis. *Glia* 2015, 63, 1452–1468. [CrossRef]
- 89. Sirko, S.; Irmler, M.; Gascón, S.; Bek, S.; Schneider, S.; Dimou, L.; Obermann, J.; de Souza Paiva, D.; Poirier, F.; Beckers, J.; et al. Astrocyte Reactivity after Brain Injury-: The Role of Galectins 1 and 3. *Glia* 2015, *63*, 2340–2361. [CrossRef]
- 90. Escartin, C.; Guillemaud, O.; Carrillo-de Sauvage, M.A. Questions and (Some) Answers on Reactive Astrocytes. *Glia* 2019, 67, 2221–2247. [CrossRef]
- Pekny, M.; Wilhelmsson, U.; Tatlisumak, T.; Pekna, M. Astrocyte Activation and Reactive Gliosis—A New Target in Stroke? Neurosci. Lett. 2019, 689, 45–55. [CrossRef]
- 92. Sofroniew, M.V. Astrocyte Reactivity: Subtypes, States, and Functions in CNS Innate Immunity. *Trends Immunol.* **2020**, *41*, 758–770. [CrossRef]
- 93. Williamson, M.R.; Fuertes, C.J.A.; Dunn, A.K.; Drew, M.R.; Jones, T.A. Reactive Astrocytes Facilitate Vascular Repair and Remodeling after Stroke. *Cell Rep.* 2021, *35*, 109048. [CrossRef]
- 94. Adams, K.L.; Gallo, V. The Diversity and Disparity of the Glial Scar. Nat. Neurosci. 2018, 21, 9–15. [CrossRef]
- Escartin, C.; Galea, E.; Lakatos, A.; O'Callaghan, J.P.; Petzold, G.C.; Serrano-Pozo, A.; Steinhäuser, C.; Volterra, A.; Carmignoto, G.; Agarwal, A.; et al. Reactive Astrocyte Nomenclature, Definitions, and Future Directions. *Nat. Neurosci.* 2021, 24, 312–325. [CrossRef]
- 96. Silver, J.; Miller, J.H. Regeneration beyond the Glial Scar. Nat. Rev. Neurosci. 2004, 5, 146–156. [CrossRef]
- Guttenplan, K.A.; Weigel, M.K.; Prakash, P.; Wijewardhane, P.R.; Hasel, P.; Rufen-Blanchette, U.; Münch, A.E.; Blum, J.A.; Fine, J.; Neal, M.C.; et al. Neurotoxic Reactive Astrocytes Induce Cell Death via Saturated Lipids. *Nature* 2021, 599, 102–107. [CrossRef]
- Rolls, A.; Shechter, R.; Schwartz, M. The Bright Side of the Glial scar in CNS Repair. *Nat. Rev. Neurosci.* 2009, 10, 235–241. [CrossRef]
- 99. Sofroniew, M.V. Molecular Dissection of Reactive Astrogliosis and Glial Scar Formation. *Trends Neurosci.* 2009, 32, 638–647. [CrossRef]
- Zamanian, J.L.; Xu, L.; Foo, L.C.; Nouri, N.; Zhou, L.; Giffard, R.G.; Barres, B.A. Genomic Analysis of Reactive Astrogliosis. J. Neurosci. 2012, 32, 6391–6410. [CrossRef]
- 101. Hayakawa, K.; Pham, L.D.D.; Arai, K.; Lo, E.H. Reactive Astrocytes Promote Adhesive Interactions between Brain Endothelium and Endothelial Progenitor Cells via HMGB1 and Beta-2 Integrin Signaling. *Stem Cell Res.* **2014**, *12*, 531–538. [CrossRef]
- 102. Li, K.; Li, J.; Zheng, J.; Qin, S. Reactive Astrocytes in Neurodegenerative Diseases. Aging Dis. 2019, 10, 664–675. [CrossRef] [PubMed]
- 103. Nathan, F.M.; Li, S. Environmental Cues Determine the Fate of Astrocytes after Spinal Cord Injury. *Neural Regen. Res.* 2017, *12*, 1964–1970. [PubMed]
- Okada, S.; Hara, M.; Kobayakawa, K.; Matsumoto, Y.; Nakashima, Y. Astrocyte Reactivity and Astrogliosis after Spinal Cord Injury. *Neurosci. Res.* 2018, 126, 39–43. [CrossRef]

- 105. Hara, M.; Kobayakawa, K.; Ohkawa, Y.; Kumamaru, H.; Yokota, K.; Saito, T.; Kijima, K.; Yoshizaki, S.; Harimaya, K.; Nakashima, Y.; et al. Interaction of Reactive Astrocytes with Type I Collagen Induces Astrocytic Scar Formation through the Integrin-N-Cadherin Pathway after Spinal Cord Injury. *Nat. Med.* 2017, 23, 818–828. [CrossRef]
- 106. Pérez, L.A.; Rashid, A.; Combs, J.D.; Schneider, P.; Rodríguez, A.; Salaita, K.; Leyton, L. An Outside-In Switch in Integrin Signaling Caused by Chemical and Mechanical Signals in Reactive Astrocytes. *Front. Cell Dev. Biol.* **2021**, *9*, 712627. [CrossRef]
- 107. Schiweck, J.; Murk, K.; Ledderose, J.; Münster-Wandowski, A.; Ornaghi, M.; Vida, I.; Eickholt, B.J. Drebrin Controls Scar Formation and Astrocyte Reactivity upon Traumatic Brain Injury by Regulating Membrane Trafficking. *Nat. Commun.* 2021, 12, 1490. [CrossRef]
- Bellahcène, A.; Castronovo, V.; Ogbureke, K.U.E.; Fisher, L.W.; Fedarko, N.S. Small Integrin-Binding LIgand N-Linked Glycoproteins (SIBLINGs): Multifunctional Proteins in Cancer. *Nat. Rev. Cancer* 2008, *8*, 212–226. [CrossRef]
- 109. Gao, H.; Fan, S.; Xie, L.; Hu, S.; Zhang, Y. Overexpression of DSPP Promotes the Proliferation and Migration of Astrocytes. *Neurosci. Lett.* **2021**, *754*, 135775. [CrossRef] [PubMed]
- Linnerbauer, M.; Wheeler, M.A.; Quintana, F.J. Astrocyte Crosstalk in CNS Inflammation. *Neuron* 2020, 108, 608–622. [CrossRef]
  [PubMed]
- Liu, L.R.; Liu, J.C.; Bao, J.S.; Bai, Q.Q.; Wang, G.Q. Interaction of Microglia and Astrocytes in the Neurovascular Unit. *Front. Immunol.* 2020, 11, 1024. [CrossRef] [PubMed]
- 112. Yoshizaki, S.; Tamaru, T.; Hara, M.; Kijima, K.; Tanaka, M.; Konno, D.; Matsumoto, Y.; Nakashima, Y.; Okada, S. Microglial Inflammation after Chronic Spinal Cord Injury Is Enhanced by Reactive Astrocytes via the Fibronectin/β1 Integrin Pathway. J. Neuroinflamm. 2021, 18, 12. [CrossRef]
- Edwards, D.N.; Salmeron, K.; Lukins, D.E.; Trout, A.L.; Fraser, J.F.; Bix, G.J. Integrin α5β1 Inhibition by ATN-161 Reduces Neuroinflammation and Is Neuroprotective in Ischemic Stroke. J. Cereb. Blood Flow Metab. 2020, 40, 1695–1708. [CrossRef]
- 114. Yun, J.H.; Park, S.W.; Kim, J.H.; Park, Y.J.; Cho, C.H.; Kim, J.H. Angiopoietin 2 Induces Astrocyte Apoptosis via αvβ5-Integrin Signaling in Diabetic Retinopathy. *Cell Death Dis.* **2016**, *7*, e2101. [CrossRef]
- 115. Dossi, E.; Vasile, F.; Rouach, N. Human Astrocytes in the Diseased Brain. Brain Res. Bull. 2018, 136, 139–156. [CrossRef]
- 116. Wang, P.; Ye, Y. Filamentous Recombinant Human Tau Activates Primary Astrocytes via an Integrin Receptor Complex. *Nat. Commun.* **2021**, *12*, 95. [CrossRef]
- 117. Katsouri, L.; Birch, A.M.; Renziehausen, A.W.J.; Zach, C.; Aman, Y.; Steeds, H.; Bonsu, A.; Palmer, E.O.C.; Mirzaei, N.; Ries, M.; et al. Ablation of Reactive Astrocytes Exacerbates Disease Pathology in a Model of Alzheimer's Disease. *Glia* 2020, 68, 1017–1030. [CrossRef]
- 118. Anderson, M.A.; Burda, J.E.; Ren, Y.; Ao, Y.; O'Shea, T.M.; Kawaguchi, R.; Coppola, G.; Khakh, B.S.; Deming, T.J.; Sofroniew, M.v. Astrocyte Scar Formation Aids Central Nervous System Axon Regeneration. *Nature* 2016, 532, 195–200. [CrossRef]
- 119. Hasel, P.; Rose, I.V.L.; Sadick, J.S.; Kim, R.D.; Liddelow, S.A. Neuroinflammatory Astrocyte Subtypes in the Mouse Brain. *Nat. Neurosci.* 2021, 24, 1475–1487. [CrossRef]
- Ahuja, C.S.; Wilson, J.R.; Nori, S.; Kotter, M.R.N.; Druschel, C.; Curt, A.; Fehlings, M.G. Traumatic Spinal Cord Injury. *Nat. Rev. Dis. Primers* 2017, 3, 17018. [CrossRef]
- 121. Roll, L.; Faissner, A. Tenascins in CNS Lesions. Semin. Cell Dev. Biol. 2019, 89, 118–124. [CrossRef]
- 122. Tucić, M.; Stamenković, V.; Andjus, P. The Extracellular Matrix Glycoprotein Tenascin C and Adult Neurogenesis. *Front. Cell Dev. Biol.* **2021**, *9*, 674199. [CrossRef]
- Wiemann, S.; Reinhard, J.; Faissner, A. Immunomodulatory Role of the Extracellular Matrix Protein Tenascin-C in Neuroinflammation. *Biochem. Soc. Trans.* 2019, 47, 1651–1660. [CrossRef]
- 124. Fawcett, J.W. An Integrin Approach to Axon Regeneration. Eye 2017, 31, 206–208. [CrossRef]
- 125. Forbes, L.H.; Andrews, M.R. Grafted Human IPSC-Derived Neural Progenitor Cells Express Integrins and Extend Long-Distance Axons within the Developing Corticospinal Tract. *Front. Cell. Neurosci.* **2019**, *13*, 26. [CrossRef]
- Sutter, P.A.; Rouillard, M.E.; Alshawi, S.A.; Crocker, S.J. Extracellular Matrix Influences Astrocytic Extracellular Vesicle Function in Wound Repair. *Brain Res.* 2021, 1763, 147462. [CrossRef]
- 127. Quraishe, S.; Forbes, L.H.; Andrews, M.R. The Extracellular Environment of the CNS: Influence on Plasticity, Sprouting, and Axonal Regeneration after Spinal Cord Injury. *Neural Plast.* **2018**, 2018, 2952386. [CrossRef]
- 128. Uede, T. Osteopontin, Intrinsic Tissue Regulator of Intractable Inflammatory Diseases. Pathol. Int. 2011, 61, 265–280. [CrossRef]
- 129. Icer, M.A.; Gezmen-Karadag, M. The Multiple Functions and Mechanisms of Osteopontin. *Clin. Biochem.* 2018, 59, 17–24. [CrossRef]
- 130. Zhou, Y.; Yao, Y.; Sheng, L.; Zhang, J.; Zhang, J.H.; Shao, A. Osteopontin as a Candidate of Therapeutic Application for the Acute Brain Injury. *J. Cell. Mol. Med.* 2020, 24, 8918–8929. [CrossRef]
- Denhardt, D.T.; Noda, M. Osteopontin Expression and Function: Role in Bone Remodeling. J. Cell. Biochem. 1998, 72, 92–102.
  [CrossRef]
- 132. Weber, G.F. The Metastasis Gene Osteopontin: A Candidate Target for Cancer Therapy. *Biochim. Biophys. Acta* 2001, 1552, 61–85. [CrossRef]
- 133. Von Neuhoff, N.; Oumeraci, T.; Wolf, T.; Kollewe, K.; Bewerunge, P.; Neumann, B.; Brors, B.; Bufler, J.; Wurster, U.; Schlegelberger, B.; et al. Monitoring CSF Proteome Alterations in Amyotrophic Lateral Sclerosis: Obstacles and Perspectives in Translating a Novel Marker Panel to the Clinic. *PLoS ONE* 2012, 7, e44401. [CrossRef] [PubMed]

- 134. Mary Varghese, A.; Sharma, A.; Mishra, P.; Vijayalakshmi, K.; Chandregowda Harsha, H.; Sathyaprabha, T.N.; Bharath, S.M.; Nalini, A.; Anand Alladi, P.; Raju, T.R. Chitotriosidase-a Putative Biomarker for Sporadic Amyotrophic Lateral Sclerosis. *Clin. Proteom.* 2013, 10, 19. [CrossRef] [PubMed]
- 135. Kariya, Y.; Kariya, Y.; Saito, T.; Nishiyama, S.; Honda, T.; Tanaka, K.; Yoshida, M.; Fujihara, K.; Hashimoto, Y. Increased Cerebrospinal Fluid Osteopontin Levels and Its Involvement in Macrophage Infiltration in Neuromyelitis Optica. *BBA Clin.* 2015, 3, 126–134. [CrossRef] [PubMed]
- 136. Broom, L.; Jenner, P.; Rose, S. Increased Neurotrophic Factor Levels in Ventral Mesencephalic Cultures Do Not Explain the Protective Effect of Osteopontin and the Synthetic 15-Mer RGD Domain against MPP + Toxicity. *Exp. Neurol.* 2015, 263, 1–7. [CrossRef]
- 137. Zhu, P.J.; Chen, C.J.; Mays, J.; Stoica, L.; Costa-Mattioli, M. MTORC2, but Not MTORC1, Is Required for Hippocampal MGluR-LTD and Associated Behaviors. *Nat. Neurosci.* 2018, *21*, 799–802. [CrossRef]
- 138. Switon, K.; Kotulska, K.; Janusz-Kaminska, A.; Zmorzynska, J.; Jaworski, J. Molecular Neurobiology of MTOR. *Neuroscience* 2017, 341, 112–153. [CrossRef]
- Calvez, M.; Hseeh, G.; Benzer, S.; Brown, A.M. Osteopontin Counters Human Immunodeficiency Virus Type 1–Induced Impairment of Neurite Growth through Mammalian Target of Rapamycin and Beta-Integrin Signaling Pathways. J. Neurovirol. 2019, 25, 384–396. [CrossRef]
- 140. Morita, K.; Sasaki, H.; Furuse, M.; Tsukita, S. Endothelial Claudin: Claudin-5/TMVCF Constitutes Tight Junction Strands in Endothelial Cells. *J. Cell Biol.* **1999**, *147*, 185–194. [CrossRef]
- 141. Nitta, T.; Hata, M.; Gotoh, S.; Seo, Y.; Sasaki, H.; Hashimoto, N.; Furuse, M.; Tsukita, S. Size-Selective Loosening of the Blood-Brain Barrier in Claudin-5-Deficient Mice. J. Cell Biol. 2003, 161, 653–660. [CrossRef]
- 142. Yamashiro, Y.; Bui, T.Q.; Ramirez, K.; Shin, J.; Kohata, T.; Ohata, S.; Nguyen, A.V.; Ohtsuki, S.; Nagayama, K.; Yanagisawa, H. Matrix Mechanotransduction Mediated by Thrombospondin-1/Integrin/YAP in the Vascular Remodeling. *Proc. Natl. Acad. Sci. USA* 2020, 117, 9896–9905. [CrossRef]
- 143. Mrugacz, M.; Bryl, A.; Falkowski, M.; Zorena, K. Integrins: An Important Link between Angiogenesis, Inflammation and Eye Diseases. *Cells* **2021**, *10*, 1703. [CrossRef]
- 144. Welser, J.; Halder, S.K.; Kant, R.; Boroujerdi, A.; Milner, R. Endothelial α6β4 Integrin Protects during Experimental Autoimmune Encephalomyelitis-Induced Neuroinflammation by Maintaining Vascular Integrity and Tight Junction Protein Expression. J. Neuroinflamm. 2017, 14, 217. [CrossRef]
- 145. Labus, J.; Wöltje, K.; Stolte, K.N.; Häckel, S.; Kim, K.S.; Hildmann, A.; Danker, K. IL-1β Promotes Transendothelial Migration of PBMCs by Upregulation of the FN/α5β1 Signalling Pathway in Immortalised Human Brain Microvascular Endothelial Cells. *Exp. Cell Res.* 2018, 373, 99–111. [CrossRef]
- Edwards, D.N.; Bix, G.J. Roles of Blood-Brain Barrier Integrins and Extracellular Matrix in Stroke. *Am. J. Physiol. Cell Physiol.* 2019, 316, C252–C263. [CrossRef]
- 147. Bhowmick, S.; D'Mello, V.; Caruso, D.; Wallerstein, A.; Abdul-Muneer, P.M. Impairment of Pericyte-Endothelium Crosstalk Leads to Blood-Brain Barrier Dysfunction Following Traumatic Brain Injury. *Exp. Neurol.* 2019, 317, 260–270. [CrossRef]
- 148. Nakamura, K.; Ikeuchi, T.; Nara, K.; Rhodes, C.S.; Zhang, P.; Chiba, Y.; Kazuno, S.; Miura, Y.; Ago, T.; Arikawa-Hirasawa, E.; et al. Perlecan Regulates Pericyte Dynamics in the Maintenance and Repair of the Blood-Brain Barrier. J. Cell Biol. 2019, 218, 3506–3525. [CrossRef]
- 149. Larochelle, C.; Uphaus, T.; Broux, B.; Gowing, E.; Paterka, M.; Michel, L.; Dudvarski Stankovic, N.; Bicker, F.; Lemaître, F.; Prat, A.; et al. EGFL7 Reduces CNS Inflammation in Mouse. *Nat. Commun.* **2018**, *9*, 819. [CrossRef]
- 150. Halder, S.K.; Kant, R.; Milner, R. Chronic Mild Hypoxia Promotes Profound Vascular Remodeling in Spinal Cord Blood Vessels, Preferentially in White Matter, via an A5β1 Integrin-Mediated Mechanism. *Angiogenesis* 2018, 21, 251–266. [CrossRef]